Wahlenbergia ramosa

Common Name(s):

Coastal Harebell

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. New Zealand: North and South Islands from East Cape and Taranaki south to the Marlborough Sounds. Most common on the shores of Cook Strait.

Habitat:

Mostly coastal (rarely inland). On rocks, cliffs, grassy places and in open scrub

Features*:

Radicate, perennial herb. Stems 100-500 mm long, 1-2 mm thick, decumbent or ascending or stiffly erect. Mature leaves and bracts pale-green, alternate, sessile, variable, usually obovate-spathulate to oblance olate on lower stem, grading to linear-lance olate in upper stem, 8×2 to 30 \times 10 mm; margins flat and distantly denticulate to strongly undulate, usually with scattered hairs on margins and lamina. Callus teeth 4-5 on each margin. Pedicels slender, 30-100 mm long. Flowers glabrous, 3-5-lobed, pastel lilac at anthesis, fading to off-white or pure white in different populations, 5-20 mm diameter, 4-9 mm long. Corolla rotate with shallow saucer-shaped tube, 1×3 -4 mm; lobes spreading, 5×4 to 8×5 mm, broadly ovate, touching or overlapping in open flower. Style thickened at top, white. Calyx lobes glabrous, 1.5×1.0 to 3.0×1.5 mm, triangular; capsule glabrous 5-7 × 4 mm, ellipsoid to obconic, with protruding apical valves. Bud at anther dehiscence short and rounded. Self-fertile. Seeds 0.5 mm long.



Caption: Baring Head. Nov 2012. **Photographer:** Pat Enright



Caption: Baring Head. Nov 2012. **Photographer:** Pat Enright

Flowering:

Fruiting:

November - April

December - May

Threats:

Not Threatened

*Attribution:

Fact Sheet Prepared by P.J. de Lange 12 June 2007. Description adapted from Petterson (1997).

References and further reading:

Petterson, J.A. 1997: Revision of the genus *Wahlenbergia* (Campanulaceae) in New Zealand. *New Zealand Journal of Botanv 35*: 9-54.

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1365