Celmisia lateralis

Common Name(s):

shrub daisy

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. South Island: North-West Nelson to North Westland (Paparoa Range).

Habitat:

Upper montane to subalpine. In rocky places, grassland, herbfield and fellfield



Caption: Mt Robert. January **Photographer:** John Smith-

Dodsworth

Features*:

Prostrate or scrambling subshrub up to 300×100 mm, with slender branched stems up to 400 mm long; branchlets usually close-set. Leaves numerous, densely imbricate, ascending, incurved at tips but becoming reflexed. Lamina $6.0-12.0 \times 1.0-1.5$ mm; linear, grading into sheath, coriaceous; upper surface glabrous, \pm glandular; lower surface similar; apex acuminate to obtuse; \pm glandular-pubescent; sheath 2-3 mm long, membranous, with a few \pm appressed hairs at base. Scape slender, 40-80 mm long, glandular-pubescent, with or without floccose hairs; bracts similar to leaves, up to 10 mm long. Capitula 10-20 mm diameter; involucral bracts 2-seriate, subulate-lanceolate to narrow-oblong, acute, up to 8 mm long, glandular, hairs \pm floccose. Rayflorets c.10 mm long, white, linear, abruptly expanded towards 3-5-toothed apex; disk-florets tubular, slightly > pappus. Achenes narrow-cylindric, compressed, 2-3 mm long, ribs with long ascending hairs. Pappus-hairs sordid-white, very slender to filiform, finely barbellate, up to c.5 mm long

Flowering: Fruiting:

November - March December - May

Threats:

Not Threatened

*Attribution:

Description adapted from Allan (1961)

References and further reading:

Allan, H.H. 1961: Flora of New Zealand. Vol. I, Government Printer, Wellington.

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 2009 Vol. 11 No. 4 pp. 285-309

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1631