



Celmisia ramulosa var. *tuberculata*

Common Name(s):

mountain daisy

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. South Island: Central Otago (mainly in the east)

Habitat:

Upper montane to subalpine rocky places and fellfield

Features*:

Small shrub or subshrub with rather stout procumbent to ascending branched woody stems up to c.20 mm long; branchlets stiff, rather close-set, more or less 50 mm long. Leaves densely imbricate, erect, finally reflexed. Lamina c.5.0-10.0 × 1.5-2.0 mm, linear-oblong to subulate, coriaceous; upper surface glabrous or nearly so, pale green; lower surface densely clad in soft white tomentum obscuring midrib; margins strongly revolute, studded with minute papillae (these mingled with a few rather stout hairs); apex obtuse, more or less trigonous-cucullate; base passing into broad pale membranous glabrous sheath more or less = lamina, with distinct midrib. Scape c.50 mm long, slender, densely glandular-pubescent; bracts few, distinctly widened toward base, more or less 5 mm long, linear. Capitula 20-25 mm diameter; involucre bracts c.5 mm long, linear-oblong, acute to subacute, densely glandular-pubescent. Ray-florets linear oblong, 15 mm or longer, 2 mm broad, obtuse, thickened and incurved at the tip, disk-florets narrow-tubular, c.5 mm long. Achenes c.3-4 mm long, narrow-cylindric, obscurely ribbed, with a few hairs. Pappus-hairs fine, white to sordid-white, up to c.4 mm long, very minutely barbellate

Flowering:

November - February

Fruiting:

January - May

Threats:

Not Threatened

*Attribution:

Description adapted from Allan (1961)

References and further reading:

Allan, H.H. 1961: Flora of New Zealand. Vol. I, Government Printer, Wellington.

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1637



Caption: Rock and Pillar Range

Photographer: John Barkla



Caption: Rock and Pillar Range

Photographer: John Barkla