Pteris comans

Common Name(s):

coastal brake, netted brake

Current Threat Status (2012):

Not Threatened

Distribution:

Indigenous (see Taxonomic Notes). New Zealand: Kermadec Islands (Raoul, the Meyers Islands and Macauley Island), Three Kings and North Island from North Cape to Bay of Plenty in the east and Waitakere coastline in the west

Habitat:

Coastal in forest especially on the sides of gullies, on banks and in valley heads. A very common offshore island fern

Features*:

Terrestrial ferns. Rhizomes short, erect, scaly. Stipes 0.25-0.6 m long, pale brown, glabrous or scaly at very base. Laminae 0.2-1.8 × 0.15-0.9 m, dark green to yellow-green, 2-3-pinnate at base, ovate, coriaceous, veins reticulate. Pinnae not overlapping; most lower secondary pinnae adnate. Ultimate segments $10-55 \times 5-10$ mm, oblong, apices tapering or bluntly pointed, margins toothed. Sori continuous along pinna margins on a marginal vein, protected by a membranous inrolled pinna margins.

Flowering:

Not applicable - spore

producing

Fruiting:

Not applicable - spore producing

Threats:

Not Threatened

*Attribution:

Fact sheet [repared for NZPCN by P.J. de Lange (17 January 2012). Description adapted from Brownsey & Smith-Dodsworth (2000).

References and further reading:

Allan, H.H. 1961: New Zealand Flora. Vol. I. Wellington, Government Printer.

Brownsey, P.J.; Smith-Dodsworth, J.C. 2000: New Zealand Ferns and Allied Plants. Auckland, David Bateman.

de Lange, P.J.; Norton, D.A.; Courtney, S.P.; Heenan, P.B.; Barkla, J.W.; Cameron, E.K.; Hitchmough, R.; Townsend, A.J. 2009: Threatened and uncommon plants of New Zealand (2008 revision). New Zealand Journal of Botany 47: 61-96.

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=2231



Caption: Motuoruhi, Coromandel Photographer: John Smith-Dodsworth



Caption: Motuoruhi, Coromandel Photographer: John Smith-

Dodsworth