# Abutilon darwinii

#### **Common Name(s):**

Chinese lantern

### **Current Threat Status (2009):**

Exotic

#### **Habitat:**

Terrestrial.

#### Features:

Perennial shrub to c. 3m high, sometimes suckering to form thickets. Stems sparsely hairy with short stellate hairs and scattered short simple hairs, becoming more or less glabrous. Leaves densely clothed in fine stellate hairs below, more or less glabrous to moderately hairy with simple and stellate hairs above. lighter green below, broadly ovate to suborbcular, cordate at base, usually 3-(5) lobed, to 1/4-1/2 radius, crenate or serrate 6-15-(18)cm long; petioles 2-20 cm long; stipules narrow-oblong to lanceolate, 6-12 mm long, deciduous. Flowers axillary, usually solitary or sometimes 2-3 in fascicles; fruiting pedicels up to 20 cm long: calvx shallowly campanulate: calvx teeth much greater than tube, triangular, acuminate, densely clothed in stellate hairs, more or less erect at fruiting, petals yellow to orange or scarlet, often with darker veins, 30-60 mm long. Cells c. 10 per fruit, covered in stellate hairs, and with longer stellate hairs along the ridged back; dorsal apical angle awnless. Seeds brown, hairy, partivulary on the back c. 3mm diameter (Webb et al 1988)

## Flowering:

January, February, March, April

#### For more information, visit:

http://nzpcn.org.nz/flora\_details.asp?ID=2399



**Caption:** Abutilon darwinii Hook.f. × A. pictum **Photographer:** John Smith-Dodsworth



Caption: Abutilon darwinii Hook.f.

× A. pictum

Photographer: John Smith-

Dodsworth