



Pimelea dura

Common Name(s):

Pimelea

Current Threat Status (2012):

Data Deficient

Distribution:

Endemic. New Zealand: South Island (inland Canterbury, in the middle catchments of the Rakaia, South Ashburton and Rangitata Rivers)

Habitat:

Montane to alpine. A species of sparsely vegetated well-drained silty, sandy, or gravelly soils on old moraines and outwash surfaces. Mostly associated with short tussock (*Festuca novae-zelandiae* and *Poa colensoi*) grassland, often growing admixed with small shrubs and herbs or within moss and lichen cushions.

Features*:

A small, much-branched, appressed shrub up to 50 mm high and 100-200 mm across. Short branches radiate from a stout main stem with short sympodial and lateral branchlets, forming small, discrete, open cushions. Young stems brown, sparsely to moderately densely covered in short, grey-white hairs; internodes 0.3-1.0 mm long. Older stems grey-brown, sometimes buried in wind-blown silt or sand. Node buttresses smooth, brown, lunate to elongate, sometimes prominent on leafless stems. Leaves decussate, ascending, loosely imbricate, sessile or on very short (0.1 mm) red petioles. Lamina 3.0-4.0 × 0.8-1.2 mm, pale glaucous-green, elliptic to ovate, or oblong, slightly keeled and adaxially concave; tip obtuse; base cuneate; adaxial surface glabrous (rarely with a few hairs), abaxial surface moderately densely covered with short hairs; stomata on both adaxial and abaxial leaf surfaces. Inflorescences terminal on branchlets, 2-6-flowered; involucre bracts 4, similar in size to adjacent leaves or larger (4.0 × 1.6 mm) partly hiding the flowers. Receptacles sparsely to moderately hairy. Plants gynodioecious. Flowers white or sometimes faintly pink-tinted, on very short (0.1 mm) pedicels, with a sparse cover of short hair outside, inside hairless. Female tube 1.5-2 mm long, ovary portion 1.5 mm, calyx lobes 0.9 × 0.9 mm; hermaphrodite tube 3-4 mm long, ovary portion 1.3 mm, calyx lobes 1.1 × 1.1 mm. Anther dehiscence introrse. Ovary apex bearing a cluster of short hairs at summit. Fruits ovoid, opaque, pinkish white, pink or pale orange 4.5 × 3.1 mm. Seeds 2.2 × 1.2 mm.

Flowering:

September - November

Fruiting:

unknown

Threats:

Burrows (2011) notes that this species is common in the Lake Heron and Lake Clearwater basins and in some reserved areas but that it is otherwise vulnerable to land development especially land being modified by topdressing, over-sowing or that which is heavily stocked with cattle or subject to the spread of wildling pines. He further observes that the ecology of the species is little known. Despite these comments the species was not assessed using the New Zealand Threat Classification Manual (Townsend et al. 2008) and so has no valid threat assessment. At this stage it seems that based on the evidence presented by Burrows (2011) an interim threat assessment of "Data Deficient" is probably the most appropriate.

***Attribution:**

Fact Sheet Prepared for NZPCN by: P.J. de Lange (30 April 2011) adapted from Burrows (2011).

References and further reading:

Burrows, C.J. 2011: Genus *Pimelea* (Thymelaeaceae) in New Zealand 4. The taxonomic treatment of ten endemic abaxially hairy-leaved species. *New Zealand Journal of Botany* 49: 41-106.

de Lange, P.J.; Norton, D.A.; Courtney, S.P.; Heenan, P.B.; Barkla, J.W.; Cameron, E.K.; Hitchmough, R.; Townsend, A.J. 2009: Threatened and uncommon plants of New Zealand (2008 revision). *New Zealand Journal of Botany* 47: 61-96.

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=6676