Xeronema callistemon f. callistemon

Common Name(s):

Poor Knights Lily, Raupo-Taranga

Current Threat Status (2012):

At Risk - Naturally Uncommon

Distribution:

Endemic. Only known from the Poor Knights Islands near Tutukaka, and from Hen (Taranga) Island,near Whangarei. Xeronema belongs to a newly established family of one genus with two species, the Xeronemataceae.

Habitat:

Rhyolite sea cliffs and rock outcrops. Occasionally in forest on rubble or as an epiphyte on pohtukawa (Metrosideros excelsa). These latter occurrences probably stem from fallen plants captured in trees or resprouting on the forest floor.

Features*:

Forming huge colonies 1-4 m across. Leaves green to yellow-green, arising from thick rhizomes, and forming flattened fans, 60-150~x 3-5 cm, Dead leaves long persistent, ultimately detached from the frayed base to form a fibrous mass. Inflorescences on green, thick, leafy stems (peduncle) up to 1 m. Racemes 10-60 cm long, aligned vertically on upper side of peduncle. Flowers crowded, numerous. Flower stalks (pedicels) 1-15 mm long, subtended by an pale green translucent leafy bract, which is obviously shorter than the buds.



Caption: Tatua Peak **Photographer:** Peter de Lange



Caption: Tatua Peak, Aorangi Island **Photographer:** Gillian Crowcroft

Flowers red, tepals 10-15 mm, outer 3 mm wide, inner 1.5 mm. Stamen filaments twice length of tepals, anthers 4 mm long, pollen tangerine-orange. Ovary 4 x 2 mm, oblong-3-angled, covered with copious nectar. Capsule 9-11 mm. Seeds ovate, 1.6 x 1 mm, black.

Flowering: Fruiting:

September to December, peaking in October.

November to January.

Threats:

None. The specie sis liste donly because it is endemic to two very small island archipelago.

*Attribution:

Fact Sheet prepared for NZPCN by P.J. de Lange 1 October 2004. Description based on Moore & Edgar (1970) supplemented by observations obtained from fresh specimens and herbarium material (see also de Lange & Cameron 1999).

References and further reading:

de Lange, P.J.; Cameron, E.K. 1999: The vascular flora of Aorangi Island, Poor Knights Islands, northern New Zealand. *New Zealand Journal of Botany 37*: 433-468.

Moore, L.B.; Edgar, E. 1970: Flora of New Zealand. Vol. II. Wellington, Government Printer.

Oliver, W.R.B. 1926: New Zealand angiosperms. *Transactions and Proceedings of the New Zealand Institute 56*: 1-5.

Oliver, H. 1968. Note on the growing of Xeronema callistemon. Wellington Botanical Society Bulletin 35: 22

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=706