



Piper excelsum subsp. excelsum

Common Name(s):

kawakawa, pepper tree

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. North and South Islands. Common from te Pahi south to about Okarito, North Canterbury and Banks Peninsula.

Habitat:

Coastal to lowland (extending up 500 m a.s.l. in warmer parts of the country). Usually an important understorey species in coastal forest.

Features*:

Small tree to at least 5 m tall; stems erect (occasionally layering), not notably lenticellate, new shoots red-green or green (leaf nerves, petioles and new stems with reddish colouring), taste peppery; pith of axes (including rachis of spike) without a mucilage core. Prophyll a collar to 0.3 (-2.2) mm high. Leaf blades submembranous, orbicular, suborbicular, at vegetative nodes to 100(-120) mm diameter, usually with 5-8 principal nerves, cordate at base, with a very narrow or closed sinus, occasionally basal lobes overlapping, upper surface of blade not bullate; petiole to 40(-60) mm long, c.0.4×as long as blade, the sheath 0.3-1.0(-2) × as long as non-sheathing part, truncate-rounded at apex and not produced there, the non-sheathing part of petiole to 4.0 mm diameter. Inflorescences solitary or 2-3 together on a short (rarely more than 10 mm long) axillary shoot, and (usually solitary) on the adjacent terminal shoot (occasionally this shoot not fertile); reduced leaf at apex of fertile shoot with a glabrous petiole and usually with a green oblong lamina at least 5 mm long, but lamina often ± lacking, especially on terminal fertile shoot. Female inflorescence erect in flowering and remaining so into fruit, peduncle to c. 1.5 cm long, spike to 60(-100) × c.6 mm diameter, with uniseriate usually 5-10-cellular hairs to 0.15 mm long on lower part of bract stalks and sparingly on rachis, these hairs not obvious on the peduncle just below the lowermost bracts; bracts peltate, bract heads 0.40-0.75 mm diameter; flowers at full emergence centred c.1.3 mm apart, emergent part of ovary ovoid; stigmas 3-4(-5), together c. 1.2 mm diameter. Male inflorescence erect, spike to c.110 mm long, proximally c.6 mm diameter, bracts and hairs as in female inflorescence; staminal filaments c. 0.25 mm long, anthers c.1.00 × 0.75 mm wide. Ripe infructescence c.10 mm diameter; fruitlets coalescent, sunken apically about the persistent dark stigmas, exocarp and mesocarp orange; seed oblong to slightly obovoid, apiculate at apex, c.2.0 × 1.5 dark brown, with (3-)4-5(-7) broad longitudinal furrows.

Flowering:

August - November

Fruiting:

Throughout the year

Threats:

Not Threatened

*Attribution:

Fact sheet prepared for NZPCN by P.J. de Lange 30 August 2005. Description based on Gardner (1997).

References and further reading:


de Lange, P.J. 2012: Taxonomic notes on the New Zealand flora: new names in *Piper* (Piperaceae). *New Zealand Journal of Botany* DOI:10.1080/0028825X.2012.708904

Gardner, R.O. 1997: *Macropiper* (Piperaceae) in the south-west Pacific. *New Zealand Journal of Botany* 35: 293-307.

Jaramillo, M.A.; Callejas, R; Davidson, C.; Smith, J.F.; Stevens, A.C.; Tepe, E.J. 2008: A phylogeny of the tropical genus *Piper* using ITS and the chloroplast intron psbJ-petA. *Systematic Botany* 33: 647-660.

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=964



Caption: Lower Hutt. Jul 2013.

Photographer: Jeremy Rolfe



Caption: Cathedral Cove, Coromandel

Photographer: John Sawyer