



Trees and shrubs of Horoeka Street Scenic Reserve, Stokes Valley, 2013



Table of Contents

Introduction	1
<i>Alectryon excelsus</i> subsp. <i>excelsus</i>	2
<i>Alseuosmia pusilla</i>	3
<i>Aristotelia serrata</i>	4
<i>Beilschmiedia tawa</i>	5
<i>Brachyglottis repanda</i>	6
<i>Carpodetus serratus</i>	7
<i>Coprosma areolata</i>	8
<i>Coprosma grandifolia</i>	9
<i>Coprosma lucida</i>	10
<i>Coprosma microcarpa</i>	11
<i>Coprosma rhamnoides</i>	12
<i>Coprosma robusta</i>	13
<i>Coprosma tenuicaulis</i>	14
<i>Cordyline australis</i>	15
<i>Cordyline banksii</i>	16
<i>Dacrycarpus dacrydioides</i>	17
<i>Dacrydium cupressinum</i>	18
<i>Elaeocarpus dentatus</i> var. <i>dentatus</i>	19
<i>Fuchsia excorticata</i>	20
<i>Gaultheria antipoda</i>	21
<i>Geniostoma ligustrifolium</i> var. <i>ligustrifolium</i>	22
<i>Griselinia littoralis</i>	23
<i>Hebe stricta</i> var. <i>atkinsonii</i>	24
<i>Hedycarya arborea</i>	25
<i>Helichrysum lanceolatum</i>	26
<i>Knightia excelsa</i>	27
<i>Kunzea ericoides</i> var. <i>ericoides</i>	28
<i>Laurelia novae-zelandiae</i>	29
<i>Leptecophylla juniperina</i> subsp. <i>juniperina</i>	30
<i>Leptospermum scoparium</i> var. <i>scoparium</i>	31
<i>Leucopogon fasciculatus</i>	32
<i>Lophomyrtus bullata</i>	33
<i>Melicytus ramiflorus</i>	34
<i>Metrosideros robusta</i>	35
<i>Myrsine australis</i>	36
<i>Myrsine salicina</i>	37
<i>Nestegis cunninghamii</i>	38
<i>Nothofagus solandri</i> var. <i>solandri</i>	39

<i>Nothofagus truncata</i>	40
<i>Olearia rani</i> var. <i>colorata</i>	41
<i>Piper excelsum</i> subsp. <i>excelsum</i>	42
<i>Pittosporum divaricatum</i>	43
<i>Pittosporum eugenioides</i>	44
<i>Pittosporum tenuifolium</i>	45
<i>Plagianthus regius</i> subsp. <i>regius</i>	46
<i>Podocarpus cunninghamii</i>	47
<i>Podocarpus totara</i> var. <i>totara</i>	48
<i>Prumnopitys ferruginea</i>	49
<i>Pseudopanax arboreus</i>	50
<i>Pseudopanax crassifolius</i>	51
<i>Pseudowintera axillaris</i>	52
<i>Rhopalostylis sapida</i>	53
<i>Schefflera digitata</i>	54
<i>Syzygium maire</i>	55
<i>Weinmannia racemosa</i>	56
Glossary	57

Made on the New Zealand Plant Conservation Network website – www.nzpcn.org.nz

Copyright

All images used in this book remain copyright of the named photographer. Any reproduction, retransmission, republication, or other use of all or part of this book is expressly prohibited, unless prior written permission has been granted by the New Zealand Plant Conservation Network (info@nzpcn.org.nz). All other rights reserved.

© 2013 New Zealand Plant Conservation Network

Introduction

This book was compiled from information stored on the website of the New Zealand Plant Conservation Network (www.nzpcn.org.nz).

This website was established in 2003 as a repository for information about New Zealand's threatened vascular plants. Since then it has grown into a national database of information about all plants in the New Zealand botanic region including both native and naturalised vascular plants, threatened mosses, liverworts and fungi.

Funding to develop the website was provided by the New Zealand Government's Terrestrial and Freshwater Biodiversity Information System Programme (TFBIS).

The species information used on the website has come from a variety of sources. The indigenous vascular plant text was written largely by Dr Peter de Lange (former Network Vice President). Peter based the descriptions on a wide range of sources including the Flora of NZ Series (Allan 1961, Moore and Edgar 1970 and Webb et al 1987) as well as numerous other taxonomic treatments. For a full bibliography of information sources see the References at the end of this book.

Where no published treatment was available Peter used herbarium specimens and his own knowledge of the flora to prepare species pages. Various other contributors have provided text and additional information to many species pages including botanists such as Mike Thorsen, John Barkla, Cathy Jones, Simon Walls, Nick Singers and many others. The threatened fungi text was written by Eric Mackenzie and Peter Buchanan (Landcare Research).

More than 200 photographers have kindly provided images to illustrate the website and for use in this book especially John Smith-Dodsworth, Jeremy Rolfe, Peter de Lange, Wayne Bennett and Gillian Crowcroft.

The New Zealand Botanic Region

The information on the Network website, from which this book was compiled, is for species that are indigenous to or naturalised within the New Zealand Botanic Region as defined by Allan (1961). The New Zealand botanic region encompasses the Kermadec, Manawatawhi/Three Kings, North, South, Stewart Island/Rakiura, Chatham, Antipodes, Bounties, Snares, Auckland Campbell island/Motu Ihupuku and Macquarie.

About the Network

The Network has more than 800 members worldwide and is New Zealand's largest non-governmental organisation solely devoted to the protection and restoration of New Zealand's indigenous plant life.

The vision of the New Zealand Plant Conservation Network is that '*no indigenous species of plant will become extinct nor be placed at risk of extinction as a result of human action or indifference, and that the rich, diverse and unique plant life of New Zealand will be recognised, cherished and restored*'.

Since it was founded in 2003 the Network has undertaken a range of conservation initiatives in order to achieve its vision.

That work has included:

- Training people in plant conservation
- Publishing plant books, reports and posters
- Raising money for the David Given Threatened Plant Research Trust to pay for plant conservation research scholarships
- Advocacy to raise awareness of the importance of plant life in general and especially New Zealand's status as a Global Centre of Plant Diversity
- Lobbying central and regional government and business to protect indigenous plant life
- Educating people about plant life through the Network website
- Connecting people through the monthly newsletter, the Network conference and the annual general meeting

What is a threatened plant?

The NZ Threatened Plant Committee was formed in 1991 and ever since then it has met at regular intervals to review the status of indigenous vascular plants. It is made up of a small group of botanists that between them have an extensive knowledge of the native plants of New Zealand. This group is chaired by Dr Peter de Lange of the New Zealand Department of Conservation.

This committee applies a set of criteria to each native plant to determine its conservation status. The resulting list of species classified as threatened is published in the NZ Journal of Botany (see for example de Lange et al. 2009). The main threat categories used are: Extinct, Critical, Endangered, Vulnerable, Declining. Other categories used are: Recovering, Relict, Naturally Uncommon, Coloniser, Vagrant and Data Deficient. For vascular plants the threat status used in this book is taken from the 2009 conservation assessment (see de Lange et al 2009).

More recently other committees have been established to review the status of non-vascular plants but their lists are yet to be published.

Alectryon excelsus subsp. *excelsus*

Common Name(s):

New Zealand ash, titoki

Threat Status (2009):

Non Threatened

Distribution:

Endemic. North and South Islands from Te Paki to Banks Peninsula

Habitat:

A widespread coastal to lowland forest tree. Often favouring well drained, fertile, alluvial soils along river banks and associated terraces. It is also a major component of coastal forests, particularly those developed within exposed situations or on basaltic or andesite volcanics. It is a common offshore island tree within the Hauraki Gulf. The large fruits are bird dispersed and so titoki trees often occur as a sparse components of most lowland forest types, throughout the North Island.

Features:

Tree between 10m and 20m tall. Branches stout, erect, all parts invested with fine, velutinous, ferruginous hairs. Bark brown. Adult leaves dark green, matt when mature, imparipinnate, alternate 80-260 mm long. Leaflets 3-7 pairs; lamina 45-105 x 19-40 mm, subcoriaceous, lanceolate, oblong or narrowly-ovate, apex, subacute often acuminate, rarely obtuse; base cuneate, truncate to oblique, upper leaf surface matt; lamina margin entire or deeply serrated 1-4 times near apex. Inflorescences axillary 90-120 mm long, sparingly branched panicles. Flowers bisexual or staminate. Petals absent. Stamens 5-8 in bisexual and 6-10 in staminate flowers, crimson. Stigma ovoid, in staminate flowers ovary tholiform, style absent, in perfect flowers broadly urceolate, style 1.5-2 mm, erect. Fruits sessile, 1-2-lobed, 14-20 x 9-14 mm, pubescent, globular, carina 3-5 mm long on one side. Seed 7-10 x 4-8 mm, subglobose, black, lustrous, sarcotesta fleshy, scarlet, papillose.

Flowering:

October - December (-June)

Fruiting:

November - August

Threats:

Not Threatened

References and further reading:

Cameron, E.K. 1998. Frost resistance in titoki *Alectryon*. Auckland Botanical Society Journal, 53: 15.

Duguid, F. 1961. Flowering in titoki. Wellington Botanical Society Bulletin, 32: 16

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1486



Caption: Carter Scenic Reserve

Photographer: John Sawyer



Caption: *Alectryon excelsus* subsp *excelsus*

Photographer: Peter de Lange

Alseuosmia pusilla

Threat Status (2009):

Non Threatened

Distribution:

Endemic. North Island from Te Moehau and Mt Pirongia south, mainly montane. South Island, in the west only from North-West Nelson to near Okarito

Threats:

Not Threatened


For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1489



Caption: Holdsworth, Tararua Forest Park.

Photographer: Jeremy Rolfe



Caption: Holdsworth, Tararua Forest Park.

Photographer: Jeremy Rolfe

Aristotelia serrata

Common Name(s):

Makomako, wineberry

Threat Status (2009):

Non Threatened

Distribution:

Endemic. North, South and Stewart Islands. Throughout, but less common in drier areas.

Habitat:

Lowland to montane forests. Often forming dense thickets following disturbance.

Features:

Dioecious tree to c. 10 m tall; trunk and branches upright, to 30 cm diam.; bark smooth, grey, spotted with lenticels; branchlets light to dark red, pubescent. Leaves opposite to subopposite; petiole slender, to 50 mm long, greenish often flushed pink; midvein conspicuous above, raised below; secondary veins obvious and raised below giving surface a wrinkled uneven appearance; lamina membranous, 5-12 x 4-8 cm, glabrate (pubescence may persist on veins below), broad-ovate, margin deeply doubly and irregularly sharply serrate, tip acuminate, base cordate to truncate, upper surface light or dark green, undersides pale green, frequently infused with purple or pink. Juvenile leaves larger. Inflorescences conspicuous, axillary, flowers 4-6 mm diam., in panicles 6-10 cm long, on slender pubescent pedicels 5-10 mm long. Sepals 4, ovate, c. 3 mm long, pubescent, pink; petals 4, 3-lobed (often deeply), c. 9 mm long, white to light pink to red. Stamens many, on glandular minutely pubescent disc, not exceeding petals. Ovary 3-4-celled, styles 3-4. Fruit a c. 8-seeded fleshy depressed-ovoid berry, 5 x 4 mm, bright red to black. Seed irregularly angled, ventral surface flattened, circular or broadly elliptic, 1.9-3.1 mm, surface irregular, aril absent.

Flowering:

September-December

Fruiting:

November-January

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1512



Caption: Waikuku, Aorangi

Photographer: John Sawyer



Caption: Waikuku, Aorangi

Photographer: John Sawyer

Beilschmiedia tawa

Common Name(s):

Tawa

Threat Status (2009):

Non Threatened

Distribution:

Endemic. Common throughout the North Island. In the South Island common from Cape Farewell east through the Marlborough Sounds. Extending south of their only in the east where it almost reaches Kaikoura (the southern limit is just north of the main town).

Habitat:

Major canopy dominant in the lowland and lower montane forests of the North Island and northern South Island. May form pure stands but usually occurs in close association with podocarps such as rimu (*Dacrydium cupressinum*).

Features:

Evergreen tree up to 35 m tall. Trunk straight, 1.2-2 m diam., with buttressed base. Bark smooth, dark brown. Branches erect to spreading, slender to moderately robust. Young branchlets, leaves and inflorescences finely pubescent, hairs simple, pale golden. Foliage opposite to sub-opposite, simple, somewhat leathery when mature. Petioles (6-)8(-12) mm. Leaves (30-)40-80(-95) x (8-)11-16(-40) mm, narrowly to broadly lanceolate sometimes elliptic, yellow-green to green, glabrous when mature, undersides glaucous. margins entire, and undulate, apex acute to acuminate. Inflorescences, an erect, axillary panicle up to 100 mm long. Flowers sexually perfect, 2-4 mm diam, pale green, perianth cleft into 6 segments, ovate-oblong, stamens 12. Fruit a pendulous, ellipsoid to ovoid drupe (20-)30(-38) x (9-)12(-18) mm, 1-seeded, pericarp fleshy, dark purple-black when ripe, glaucous or shiny.

Flowering:

(October-) January (- May)

Fruiting:

(December-) January (- March)

Threats:

Not Threatened

References and further reading:

Moorfield, J. C. (2005). *Te aka : Māori-English, English-Māori dictionary and index*. Pearson Longman: Auckland, N.Z.

Landcare Research. Ngā Tipu Whakaoranga - Māori Plant Use Database.
<http://maoriplantuse.landcareresearch.co.nz>



Caption: Flowers ex Hakarimata Range.

Photographer: John Braggins



Caption: Flowers of *Beilschmiedia tawa*

Photographer: Wayne Bennett

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1541

Brachyglottis repanda

Common Name(s):

Rangiora, bushmans toilet paper

Threat Status (2009):

Non Threatened

Distribution:

Endemic. North Island throughout. South Island - north west Nelson to just south of Greymouth in the west, and near Kekerengu in the east. Naturalised on Banks Peninsula, Otago Peninsula, and on Stewart Island at Oban.

Habitat:

Common in coastal, lowland and lower montane shrubland and open forest. Often a pioneer species.

Features:

Shrub to small tree up to 6 m or more tall. Trunk one or more arising from ground, covered in somewhat corky bark. Branches stout, spreading, rather brittle, initially densely clad in fine white to buff tomentum becoming glabrescent with age. Petiole stout, grooved, 80-100 mm long. Leaves leathery, 50-250(-300) X 50-20(-30) mm, dark green to pale green above, undersides clad in fine, appressed vivid white hairs, broad- to ovate-oblong, obtuse to subacute, obliquely cordate to truncate at base, margins distantly dentately lobed to sinuate. Inflorescence a much branched panicle. Capitula 5 mm diam., numerous, without ligules (discoid). Involucral bracts 3 mm long, narrow-oblong to narrow spathulate, margins scarious except at base. Florets 10-12, yellow. Seeds (cypselae) narrowly oblong-elliptic to oblong elliptic, 1-1.8 mm long, ribs 6, rounded, broad. Pappus 2-3 mm, buff-yellow, scabrid.

Flowering:

(July-) August-October
(-November)

Fruiting:

(October-) November-
December (-January)

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1562



Caption: *Brachyglottis repanda*

Photographer: Wayne Bennett



Caption: *Brachyglottis repanda*

Photographer: Wayne Bennett

Carpodetus serratus

Common Name(s):

Putaputaweta, marbleleaf

Threat Status (2009):

Non Threatened

Distribution:

Endemic. Widespread. North, South and Stewart Islands.

Habitat:

Coastal to montane (10-1000 m a.s.l.). Moist broadleaf forest, locally common in beech forest. A frequent component of secondary forest. Streamsides and forest margins.

Features:

Monoecious small tree up to 10 m tall. Trunk slender, bark rough, corky, mottled grey-white, often knobbled due to insect boring. Juvenile plants with distinctive zig-zag branching which is retained to a lesser degree in branchlets of adult. Leaves broad-elliptic to broad-ovate or suborbicular; dark green, marbled; membranous becoming thinly coriaceous; margin serrately toothed; tip acute to obtuse. Juvenile leaves 10-30 mm x 10-20 mm. Adult leaves 40-60 mm x 20-30mm. Petioles c. 10 mm; petioles, peduncles and pedicels pubescent; lenticels prominent. Flowers in panicles at branchlet tips; panicles to 50 x 50 mm; flowers 5-6 mm diam.; calyx lobes c. 1 mm long, triangular-attenuate; petals white, ovate, acute, 3-4 mm long. Stamens 5-6, alternating with petals; filaments short. Stigma capitate, tip dark; ovules many. Fruit an indehiscent subfleshy-fleshy capsule, 4-6 mm diam., black when mature; cupped in remains of calyx. Seeds many per capsule, in 3-5 locules, small, 1-2 mm long; testa reticulate.

Flowering:

November-March

Fruiting:

January-February (though dried fruit present at any time)

Threats:

Not Threatened.

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1605



Caption: Rotoiti Mainland Island, Nelson Lakes National Park

Photographer: John Sawyer



Caption: Rotoiti Mainland Island, Nelson Lakes National Park

Photographer: John Sawyer

Coprosma areolata

Common Name(s):

Thin-leaved coprosma

Threat Status (2009):

Non Threatened

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1701



Caption: *Coprosma areolata*

Photographer: Wayne Bennett



Caption: *Coprosma areolata*

Photographer: Wayne Bennett

Coprosma grandifolia

Common Name(s):

Kanono, manono, large-leaved coprosma, raurekau

Threat Status (2009):

Non Threatened

Distribution:

Endemic. North to South Islands. In the South Island extending to Lake Ianthe in the west and the Marlborough Sounds in the east.

Habitat:

Common in the understorey of forest, and in sheltered shady sites from the coast to montane and cloud forest. In areas of high rainfall can be a major component of shrublands, and within regenerating forest. Often common along the margins of logging tracks and roads.

Flowering:

(March-) April (-June) but may also occasionally flower in September.

Fruiting:


(September-) October-January (-April)

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1717



Caption: Leaf of *Coprosma grandifolia*

Photographer: Wayne Bennett



Caption: *Coprosma grandifolia*

Photographer: Wayne Bennett

Coprosma lucida

Common Name(s):

Karamu, shining karamu

Threat Status (2009):

Non Threatened

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1719



Caption: *Coprosma lucida*

Photographer: Wayne Bennett



Caption: Leaf of *Coprosma lucida*

Photographer: Wayne Bennett

Coprosma microcarpa

Common Name(s):

Small seeded coprosma

Threat Status (2009):

Non Threatened

Threats:

Not Threatened


For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1722



Caption: *Coprosma microcarpa* at Kaitoke. April 2004.

Photographer: Jeremy Rolfe



Caption: Whakapapa, Ruapehu

Photographer: John Smith-Dodsworth

Coprosma rhamnoides

Threat Status (2009):

Non Threatened

Threats:

Not Threatened


For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1731



Caption: *Coprosma rhamnoides*
(female)

Photographer: Wayne Bennett



Caption: Fruit of *Coprosma rhamnoides*

Photographer: Wayne Bennett

Coprosma robusta

Common Name(s):

Karamu, glossy karamu

Threat Status (2009):

Non Threatened

Distribution:

Endemic. North and South Islands. Naturalised on the Chatham Islands within a small area between Waitangi and Owenga.

Habitat:

Common throughout coastal, lowland and lower montane habitats within shrublands and open sites within forest.

Features:

Shrub or small tree up to 6 m tall. Branches numerous, stout, erect to somewhat spreading. Petioles stout, 10-20 mm long. Stipules fused towards base, obtuse, glabrous with one of two prominent, black, glandular denticles. Leaves 70-120 x 30-40-50 mm, leathery, dark green above, paler green beneath, glabrous, elliptic, elliptic-oblong to broad-ovate, acute or obtuse, apex mucronate. Venation reticulated, conspicuous. Male flowers in axillary many-flowered glomerules, corolla conspicuous, lobes triangular, acute, stamens 4-5, prominent. Females in compound clusters on peduncles 10-15 mm. Calyx and corolla much reduced, stigmas prominent. Drupe dark orange (rarely yellow), 8-8 x 4-5 mm, oblong to narrow-ovoid.

Flowering:

(July-) August-September (- November)

Fruiting:

(March-) April-May (-July)

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1733



Caption: Fruit of *Coprosma robusta*

Photographer: Wayne Bennett



Caption: *Coprosma robusta* (Karamu)

Photographer: Wayne Bennett

Coprosma tenuicaulis

Common Name(s):

Swamp coprosma, hukihuki

Threat Status (2009):

Non Threatened

Distribution:

Endemic. North and South Islands from Kaitaia to about Okarito

Habitat:

Lowland (rarely montane) in swamps and boggy ground, poorly drained shrubland and riparian forest.

Features:

Shrub up to c.3 m tall; branches slender, filiramulate, subdivaricate, usually interlacing; branchlets slender, pubescent. Leaves on more or less winged, slender, pubescent petioles 3-5 mm long. Stipules small, triangular, subacute, more or less pubescent, ciliate, terminal tuft conspicuous. Lamina subcoriaceous, glabrous, orbicular-ovate, spathulate, obtuse, minutely apiculate, more or less 8-13 × 8-10 mm. Reticulations of veins evident on both surfaces. Male flowers in fascicles of 3-6; calyx 0; corolla funnelform, lobes more or less = tube, ovate, acute. Female flower solitary or in axillary clusters of 2-4 along short branchlets; calyx-teeth minute; corolla funnelform, lobes ovate, acute, = or > tube. Drupe black, globose, 3-4 mm diameter.

Flowering:

July - October

Fruiting:

March - May

Threats:

Not Threatened

References and further reading:

Cheeseman, T.F. 1885. Description of three new species of *Coprosma*. Transactions of the Royal Society, 18: 315-317

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1739



Caption: *Coprosma tenuicaulis*

Photographer: Wayne Bennett



Caption: *Coprosma tenuicaulis*

Photographer: Wayne Bennett

Cordyline australis

Common Name(s):

Cabbage tree, ti, ti kouka, palm lily

Threat Status (2009):

Non Threatened

Distribution:

Endemic. Common in the North, South and Stewart Islands. Probably naturalised on the Chatham Islands.

Habitat:

Widespread and common from coastal to montane forest. Most commonly encountered on alluvial terraces within riparian forest.

Features:

Tree up to 20 m tall, trunk stout, 1.5-2 m diam, many-branched above (prior to flowering, trunk slender and solitary, branching happens after the first flowering). Bark corky, persistent, fissured, pale to dark grey. Leaves numerous (0.2-)0.3-1(-1.5) x (0.2)-0.3(-0.6) m, dark to light green, narrowly lanceolate to lanceolate, erect to erecto-patent, scarcely inclined to droop, midrib indistinct. Petiole indistinct, short. Inflorescence a panicle. Peduncle stout, fleshy 40 mm or more in diam., panicle of numerous flowers, (0.6-)1(-1.8) x).3-0.6(-0.8) m, branching to third or fourth order, these well spaced, basal bracts green and leaf-like, ultimate racemes 100-200 mm long, 20 mm diam., bearing well-spaced to somewhat crowded, almost sessile to sessile flowers and axes. Flowers sweetly perfumed, perianth 5-6 mm diam., white, tepals free almost to base, reflexed. Stamens about same length as tepals. Stigma short, trifid.

Flowering:

(September-) October-
December (-January)

Fruiting:

(December-)
January-March

Threats:

Populations have been decimated from some parts of the country due to a mysterious illness linked to a Myoplast Like Organism (MLO) which is believed to cause the syndrome known as Sudden Decline. Plants stricken with this illness suddenly, and rapidly, wilt, with the leaves failing off still green. If the bark is peeled off the base of the tree near the soil line blackened or rotten spots are typically present. Once stricken with Sudden Decline there is no cure and the trees can die within days. Recently there has been some evidence to suggest the severity of Sudden Decline is lessening.

References and further reading:

Beever, R. et al. 1996. Sudden decline of cabbage tree. NZ Journal of Ecology, 20(1): 53-68

Duguid, F. 1976. *Cordyline australis* at Lake Kopureherehe. Wellington Botanical Society Bulletin, 39: 46-47

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1744



Caption: Awhitu Regional Park,

Auckland region

Photographer: John Sawyer



Caption: *Cordyline australis*

Photographer: Wayne Bennett

Cordyline banksii

Common Name(s):

Ti ngahere, Cabbage tree, Ti rakau

Threat Status (2009):

Non Threatened

Distribution:

Endemic. Common throughout the North Island, In the South widespread through the northern half, extending in the west to about Haast with occasional as unsubstantiated reports of it from the coastal portion of Fiordland.

Habitat:

Common in coastal, lowland, and lower montane forests. Occasionally extending into subalpine habitats in the South Island. Often found in shrublands where it is sympatric with, and often hybridises with *Cordyline pumilio*. Tolerant of a wide range of situations.

Features:

Shrub or small tree up to 4 m tall. Stems (1-)4(- many) 100-150 mm diam., arising from ground level, subequal, sparingly branched. Leaves numerous, 1-2 x 0.4-0.8 m, lanceolate (somewhat "paddle-shaped") broad about middle and drooping from there, narrowed above base into a long, narrow, channeled petiole. Midrib flat, prominent for entire leaf length. Inflorescence a panicle. Peduncle stout, fleshy, 30-40 x 200 mm. Panicle 1-2 m, often smaller, broadly pyramidal, openly branched to third order, lower bracts green and leaf-like. Ultimate racemes 150-300 mm, 200 mm or more in diam., bearing, numerous, well spaced sessile flowers in axes. Flowers sweetly perfumed, perianths 10 mm long, white; tepals fused near base, rather open. Stamens same length as tepals. Stigma shortly trifid. Fruit 4-5 mm diam., globose, white, bluish-white, or blue. Seeds 2 mm diam., black, glossy, 2 sides flat the other convex.

Flowering:

November - January

Fruiting:

February - April

Threats:

Common and not threatened. Does not seem so susceptible to Sudden Decline as *C. australis* has proved to be.

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1745



Caption: Ruahine Range

Photographer: John Sawyer



Caption: *Cordyline banksii*

Photographer: Wayne Bennett

Dacrycarpus dacrydioides

Common Name(s):

Kahikatea, white pine

Threat Status (2009):

Non Threatened

Distribution:

Endemic. North, South and Stewart Islands

Habitat:

Lowland forest, formerly dominant on frequently flooded, and/or poorly drained alluvial soils. Occasionally extends into lower montane forest. Once the dominant tree of a distinct swamp forest type all but extinct in the North Island - the best examples remain on the West Coast of the South Island.

Features:

Stout, dioecious, cohort-forming conifer, 50 (-65) m. tall. Trunk 1(-2) m diam., often fluted and buttressed. Bark grey to dark-grey, falling in thick, sinuous flakes. Wood white, odourless. Trunks bare for 3/4 of length, subadults with a distinctive columnar growth habit, branches arising from 1/3 to 1/2 of trunk length. Branchlets slender, drooping. Leaves of juveniles subdistichous, subpatent, narrow-linear, subfalcate, acuminate, decurrent, 3-7 x 0.5-1mm red, wine-red, dark-green to green.; of subadults less than or equal to 4 mm., dark green or red; those of adults 1-2 mm., imbricating, appressed, keel, subtrigonous, lanceolate-subulate to acuminate with broader base, brown-green or glaucous. Male cones terminal, oblong, 10 mm. Pollen pale yellow. Ovule, terminal, solitary glaucescent. Receptacle fleshy, oblong, compressed, warty, 2.5-6.5 mm., yellow to orange-red. Seed broadly obovate to circular (4-)4.5-6 mm diam., purple-black, thickly covered in glaucous bloom.

Flowering:

October - January

Fruiting:

February - April

Threats:

Not Threatened, although as a forest-type it has been greatly reduced through widespread logging. Very few intact examples of kahikatea-dominated forest remain in the North Island.

References and further reading:

Gardner, R. 2001. Notes towards an excursion Flora. Rimu and kahikatea (Podocarpaceae). Auckland Botanical Society Journal, 56: 74-75

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=2099



Caption: Fruit.

Photographer: © John Braggins



Caption: *Dacrycarpus dacrydioides*

Photographer: Wayne Bennett

Dacrydium cupressinum

Common Name(s):

Rimu, red pine

Threat Status (2009):

Non Threatened

Distribution:

Endemic. North, South and Stewart Islands from North Cape south. Uncommon in large parts of the eastern South Island. Facultatively extinct on Banks Peninsula, where one natural tree is all that remains. Rimu is the type of the genus *Dacrydium*.

Habitat:

Lowland to montane forest - occasionally ascending to subalpine scrub.

Features*:

Dioecious conifer 35(-60) m tall. Adult trees with trunk bare of branches for 3/4 of length. Trunk stout, 1.5-2 m diam., bark dark brown, falling off in large thick flakes. Wood dark red. Branches in juveniles numerous, slender, branchlets pendulous. Adult branches few, spreading, branchlets slender, pendulous. Leaves dark green, bronze-green, red-green or orange, imbricate, those of juveniles 4-7(-10) mm., 0.5-1 mm wide, keeled, acute, linear-subulate, subfalcate, decurrent; those of subadults ascending, incurved 4-6 mm., rhomboid; of adults similar but appressed, 2-3 mm., rigid, subacute, trigonous. Male and Female "cones" first appear on subadults. Male cones (strobili) solitary or paired, terminal 5-10 mm., oblong. Pollen yellow. Ovules solitary, terminal on up-curved branchlets. Receptacle a fleshy red or deep-orange cup 1-2 mm long. Seed oblong or elliptic-oblong, compressed in section, 3-3.8(-4) mm long, semi-glossy, dark-brown.

Fruiting:

Flowering:

December -
March

Fruits take a year or more to mature and co-occur with young female cones, they are most frequently seen between February and May.

Threats:

Not Threatened, although as a forest-type it has been greatly reduced through widespread logging. Very few intact examples of rimu-dominated forest remain in the North Island.

*Attribution:

P.J. de Lange 3 February 2006. Description adapted from Allan (1961), Webb & Simpson (2001), fresh material and herbarium specimens.

References and further reading:

Allan, H.H. 1961: Flora of New Zealand. Wellington, Government Printer.

Gardner, R. 2001. Notes towards an excursion Flora. Rimu and kahikatea (Podocarpaceae). Auckland Botanical Society Journal, 56: 74-75

Kirk, T. 1889: The Forest Flora of New Zealand. Wellington, Government Printer.

Webb, C.J.; Simpson, M.J.A. 2001: Seeds of New Zealand Gymnosperms and Dicotyledons. Christchurch, Manuka Press.

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=2100



Caption: Pihanga, Tongariro National Park

Photographer: John Sawyer



Caption: Pihanga, Tongariro National Park

Photographer: John Sawyer

Elaeocarpus dentatus var. *dentatus*

Common Name(s):

Hinau

Threat Status (2009):

Non Threatened

Distribution:

Endemic. North, and South Islands

Habitat:

Common tree of mainly coastal and lowland forest though occasionally extending into montane forest.

Features:

Tree up to 20 m tall (usually less), with broad spreading crown. Trunk 1 m diam., bark grey. Branches erect then spreading, branchlets silky hairy when young. Petioles stout, 20-25 mm long. Leaves leathery, (50-)100-120 x 20-30 mm, narrow- to obovate-oblong, broad-obovate, oblanceolate, apex obtuse or abruptly acuminate, dark green and glabrescent above, off-white, silky-hairy below; margins somewhat sinuate, recurved, serrate to subentire. Inflorescence a raceme 100-180 mm long, 8-12(-20)-flowered. Pedicels 10 mm long, silky-hairy. Flowers drooping, (8-)12(-15) mm diam., sepals lanceolate-oblong, 6 mm long, petals white, obovate-cuneate, 3-5-lobed, c. 10 mm long. Stamens 10-20. Fruit a fleshy, ovoid purple-black 12-18 x 9 mm, drupe. Endocarp deeply furrowed and wrinkled.

Flowering:

October - February

Fruiting:

December - May(-June)

Threats:

Not Threatened.

References and further reading:

Allan, H.H. 1961: Flora of New Zealand. Vol. I. Wellington, Government Printer.

Druce, A.P. 1993: Indigenous vascular plants of New Zealand. Ninth Revision. Unpublished Checklist held at Landcare Research, Lincoln, New Zealand.

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1830



Caption: Flowers of *Elaeocarpus dentatus*

Photographer: Wayne Bennett



Caption: Flowers of *Elaeocarpus dentatus*

Photographer: Wayne Bennett

Fuchsia excorticata

Common Name(s):

Kotukutuku, Tree fuchsia

Threat Status (2009):

Non Threatened

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1901



Caption: Rotoiti Mainland Island,
Nelson Lakes National Park

Photographer: John Sawyer



Caption: Rotoiti Mainland Island,
Nelson Lakes National Park

Photographer: John Sawyer

Gaultheria antipoda

Common Name(s):

Bush snowberry, fools beech

Threat Status (2009):

Non Threatened

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1912



Caption: *Gaultheria antipoda*

Photographer: Wayne Bennett



Caption: *Gaultheria antipoda*

Photographer: Wayne Bennett

Geniostoma ligustrifolium var. *ligustrifolium*

Common Name(s):

Hangehange

Threat Status (2009):

Non Threatened

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1923



Caption: Puketi Forest, Northland

Photographer: Dean Baigent-Mercer



Caption: Puketi Forest, Northland

Photographer: Dean Baigent-Mercer

Griselinia littoralis

Common Name(s):

Broadleaf, kapuka, papauma

Threat Status (2009):


Non Threatened

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1957



Caption: Mt Frith, Rimutaka

Photographer: John Sawyer



Caption: Coromandel, October

Photographer: John Smith-Dodsworth

Hebe stricta var. *atkinsonii*

Common Name(s):

Koromiko

Threat Status (2009):

Non Threatened

Distribution:

Endemic. In the North Island common from the Manawatu Gorge south to Wellington. In the South Present in the Marlborough Sounds south to about Rarangi.

Habitat:

Common in successional habitats from coastal areas to lower montane habitats.

Features:

Shrub (1-)2 m tall. Branchlets finely pubescent. Stem internodes longer than stem diameter. Leaf bud without sinus. Leaves, spreading, 50-100 (-110) mm, dull yellow-green (not glossy), narrow-elliptic to linear-lanceolate, somewhat leathery, apex acute (rarely acuminate), leaf margin usually entire, occasionally toothed. Inflorescence lateral, racemose, about length of leaves, sometimes drooping, with the exception of ciliolate bracts and calyx, the remaining inflorescence structures glabrous. Flowers sweetly (sometimes over powerfully so) scented, lilac, mauve or white. Corolla tube 6 mm, exceeding calyx, narrow, cylindric, lobes rounded. Capsules < 5 mm long, glabrous, usually erect.

Flowering:

(July-) August (-October)
but flowering can also
occur sporadically
throughout the year

Fruiting:

(September-) November (-
January) but seed capsules
may be found throughout
the year

Threats:

Not Threatened

References and further reading:

Bayly M. and Kellow A. (2006). An Illustrated Guide to New Zealand Hebes. Te Papa Press: Wellington

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=2010



Caption: Stokes Valley. Feb 2010.

Photographer: Jeremy Rolfe



Caption: Putangirua Pinnacles,
Palliser Bay. Aug 2010.

Photographer: Jeremy Rolfe

Hedycarya arborea

Common Name(s):

Porokaiwhiri, Pigeonwood

Threat Status (2009):

Non Threatened

Distribution:

Endemic. Three Kings, North and South Islands. In the South island uncommon in the east south of Kaikoura reaching its southern limit on that coastline on Banks Peninsula, it is more ranging in the west reaching northern Fiordland at least.

Habitat:

A common forest tree of coastal and lowland forest, extending into montane areas in the warmer parts of the North Island

Features:

Tree up to 12 m. tall; trunk up to 0.5m dbh, clear of branches for first few metres; bark dark grey to brown-grey, firm (not flaking) finely tessellated. Branches numerous, upright to spreading; branchlets finely brown-pubescent at tips. Leaves coriaceous, glabrous except for midrib and main veins and petioles, adaxially dark green, glossy or glaucescent, abaxially similar but paler and dull; petioles 10-15-20(-35)mm long; lamina 40-120(-180) × 25-30(-50-60) mm, elliptic-obovate, oblanceolate to lanceolate, cuneately narrowed to base, obtuse to subacute or acute, margins distantly serrate (with occasional subentire leaves) or toothed. Inflorescence a branched raceme; peduncles and pedicels slender, pubescent. Male with perianth c.10 mm diameter, pubescent, stamens numerous, anthers sessile. Female with perianth c.6 mm diameter; carpels up to 20. Drupe 1-seeded, ovoid, 10-15(-16) mm long, red or orange-red up to 10 per branch. Endocarp 9-14 mm long, elliptic to obovate, rarely circular, brown to grey-brown, surface ± smooth, usually with a few irregular bumps and/or longitudinal ridges. Description adapted from Allan (1961) and Webb & Simpson (2001).

Flowering:

December - February

Fruiting:

March - June

Threats:

Not Threatened

References and further reading:

Allan, H.H. 1961: Flora of New Zealand. Vol. I, Wellington, Government Printer.

de Lange, P.J.; Cameron, E.K. 1999: The Vascular Flora of Aorangi Island, Poor Knights Islands, Northern New Zealand. New Zealand Journal of Botany 37: 433-468.

de Lange, P.J.; Murray, B.G. 2002: Contributions to a chromosome atlas of the New Zealand flora – 37. Miscellaneous families. New Zealand Journal of Botany 40: 1-24.

Webb, C.J.; Simpson, M.J.A. 2001: Seeds of New Zealand Gymnosperms and Dicotyledons. Christchurch, Manuka Press.

Wright, A. E. 1984: Beilschmiedia Nees (Lauraceae) in New Zealand. New Zealand Journal of Botany 22: 109-125.

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=805



Caption: *Hedycarya arborea* (Porokaiwhiri)

Photographer: Wayne Bennett



Caption: Fruit of *Hedycarya arborea*

Photographer: Wayne Bennett

Helichrysum lanceolatum

Threat Status (2009):

Non Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=809



Caption: Leitz's Gully, Otago

Photographer: Jesse Bythell



Caption: Otago Peninsula

Photographer: John Barkla

Knightia excelsa

Common Name(s):

Rewarewa, NZ honeysuckle

Threat Status (2009):

Non Threatened

Distribution:

Endemic monotypic genus. North and South Islands. Common in the North Island, but confined to the Marlborough Sounds in the South Island.

Habitat:

A common tree of coastal, lowland and lower montane shrubland, secondary regrowth, and on occasion mature forest. Frost-tender when young so generally scarce from cooler, frost-prone habitats - nevertheless it can be very common in suitable sites on the Central Volcanic Plateau of the North Island.

Features:

Tall tree with columnar (fastigiate) growth-form up to 30 m tall. Trunk up to 1 m diam. Bark dark brown. Branches erect, fastigiate, at first angled, clad in red-brown (rust-coloured), velutinous, tomentum. Juvenile leaves yellow-green, 150-300(-400) x 10-15 mm, narrowly linear-lanceolate, sometimes forked 2,3 or 4 times, margins acutely serrated. Adult leaves dark green, 100-150(-200) x 25-40 mm, broad lanceolate to narrow-oblong or oblong, sometimes obovate, occasionally forked, rigid, bluntly and coarsely serrated, covered in deciduous velutinous red-brown pubescence. Inflorescence a stout raceme up to 100(-180) mm x 60 mm, densely flowered. Pedicels and perianth clad in red-brown, velutinous tomentum. Flowers sexually perfect. Perianth 4, exterior covered in red-brown tomentum, interior dark crimson, segments at first cylindric and fused, soon separating and curling spirally. Stamens 4, filaments crimson, short, anthers long, linear, rich golden-yellow. Ovar sessile. Style long, crimson, long persistent. Fruits, follicles 30-40 mm long, 2-valved, woody, pubescent; valves tapering to persistent style. Seeds 10 mm, apex terminated by 15 mm long wing.

Flowering:

(September-) October-December

Fruiting:

October-January (fruit takes a year to mature, so fruit and flowers may co-occur)

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1383



Caption: Awhitu Regional Park, Auckland region

Photographer: John Sawyer



Caption: Rangitoto Island

Photographer: John Barkla

Kunzea ericoides var. *ericoides*

Common Name(s):

Manuoea, Titira, Atitira, Manuka-Rauriki, Kanuka

Threat Status (2009):

Non Threatened

Distribution:

Endemic. As circumscribed here *K. ericoides* var. *ericoides* is endemic to the South Island, where it is common from North West Nelson and the Marlborough Sounds south to the upper Buller River. From here it is common along the northern margin of the Buller to the upper Wairau River, from where it extends along the southern Richmond Range to Rarangi. Outliers occur in the east south of Rarangi in pockets to Kaikoura and the coastal portion of the north Canterbury foothills, and in the west around Karamea, the Lower Buller Gorge, and the upper Ahaura River. Outside this area there are a number of distinctive New Zealand variants which may warrant formal description. An allied complex of species and possibly unnamed species occurs in Australia.

Habitat:

Coastal to lowland shrubland, regenerating forest and forest margins, also present in montane forest, ultramafic shrubland and very occasionally present in subalpine shrubland.

Features:

Shrub or tree (2-)10(-20) m tall. Usually with a single trunk. Trunk slender, erect, often multi-trunked from base. Branches numerous, slender, and pendulous, branchlets slender, brittle. Bark loose, flaking readily into tabular, fibrous shards, typically with much secondary peeling; secondary peels often inrolling, like wood shavings. Branchlets glabrescent to glabrous, hairs if present (20x magnification) are erect, sparse, and short (like stubble). Leaves bright green, linear to linear-filiform (6-)8(-12) x (0.8-)1(-1.3) mm. Inflorescences corymbiform racemes, (1-)8(-20)-flowered. Flowers (4-)6(-8) mm diam., faintly to strongly scented. Petals 5(-6), white. Stamens (9-)20(-32), antipetalous (1-)3, antisepalous variable. Ovary 5 locular, stigma broad, capitate. Capsule long persistent, grey, obconic, sepals persistent. Seeds numerous, rather fine, orange-yellow.

Flowering:

(October)-November-January
(-February)

Fruiting:

(November-)December
(-March)

Threats:

Not threatened, though some stands are at risk from clearance for farmland or through felling for firewood.


For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=885



Caption: *Kunzea ericoides* - tree showing weeping branches characteristic of this species

Photographer: Peter de Lange



Caption: Marahau

Photographer: Peter de Lange

Laurelia novae-zelandiae

Common Name(s):

Pukatea

Threat Status (2009):

Non Threatened

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=901



Caption: *Laurelia novae-zelandiae*

Photographer: John Smith-Dodsworth



Caption: Lake Rotokare, Taranaki. Jun 2012.

Photographer: Colin Ogle

Leptecophylla juniperina subsp. *juniperina*

Common Name(s):

Prickly Mingimingi, Mingimingi

Threat Status (2009):

Non Threatened

Distribution:

Indigenous. New Zealand (North, South and Stewart Islands), also Australia (Tasmania only)

Habitat:

Coastal to montane, in scrub and forest.

Features:

Dioecious, compact or tall shrubs 0.4-2.0 m tall. Bark firm, fibrous, weakly tessellated and furrowed, dark grey brown or dark brown. Stems grey, brown or grey brown; branchlets usually brown but occasionally yellow-brown or red-brown, rounded, scabrous or puberulent. Leaves spreading or occasionally reflexed, dark green, bronze-green or green, narrowly ovate, 4.0-18.0 × 1.0-2.1 mm, apex acute, tip pungent, 0.4-1.6 mm long, margin flat, glabrous or ciliolate toward apex; upper surface glabrous or puberulent at base, lower surface with intervenal papillae and 5 veins; petiole erect 0.6-1.7 mm long, appressed to stem, glabrous or puberulent on the upper surface. Flowers sickly sweet fragrant, solitary, terminal and axillary on erect or recurved pedicels 2-5 mm long in males, 1.3-3.0 mm in females; bracts ovate, 0.5-0.9 × 0.6-1.4 mm, obtuse, glabrous, margin usually ciliolate at the apex; bracteoles and sepals ovate or elliptic, obtuse, glabrous; bracteoles 8-24 per flower, imbricate, 1.2-2.4 × 1.1-2.0 mm; sepals 1.7-3.1 × 1.1-2.3 mm. Corolla tube white or cream, campanulate, exceeding the calyx, 1.5-2.8 mm long in males, 1.6-2.8 mm long in females, usually glabrous; lobes white or cream, shorter than tube 1.1-2.3 mm long, apex acute, glabrous or with short, sparse hairs. Anthers of male flowers 1.1-2.0 mm long, half-exserted; filaments 0.2-0.5 mm long, slightly exserted and visible between the lobes. Ovary ± spherical, 0.5-1.0 × 0.6-1.3 mm, glabrous, 4-6-celled; style straight, glabrous, attenuate from the ovary, 1.0-1.8 mm long in males, 0.9-1.5 mm long in females; stigma 0.1-0.2 mm tall; nectary 0.3-0.7 mm tall, of distinct scales or weakly adherent scales separating with pressure, margin toothed or rounded and occasionally with hairs. Drupe fleshy, white, pink, pinkish white or red, slightly flattened, spherical, 4-7 × 5-9 mm. Endocarp transversely elliptic, terete, 2.0-3.5 × 3.0-5.0 mm, longitudinally ridged, 5-6 of these distinctly more prominent; apex rounded to a broad flat or concave end; base rounded but narrower than apex, with a small hollow. Surface brown to dark brown, fading with age to light brown or fawn. Internally 4-6-celled, usually with 2-3 of these filled

Flowering:

Throughout the year

Fruiting:

Throughout the year

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=907



Caption: Coal Island

Photographer: John Barkla



Caption: Leptecophylla juniperina
subsp. juniperina

Photographer: Wayne Bennett

Leptospermum scoparium var. *scoparium*

Common Name(s):

Manuka, tea tree, kahikatoa

Threat Status (2009):

Non Threatened

Distribution:

Indigenous to New Zealand and Australia. Most Australian forms of *L. scoparium* do not match the range seen in New Zealand. However, plants from Tasmania are very similar to, if not identical with some South Island forms, differing mainly by their wider leaf base, and longer, more pungent leaf apex. Manuka was also collected once from Rarotonga by Thomas Cheeseman in the 1800s. It has not been found there since, and is assumed to have been a failed introduction. Further study using DNA sequencing is underway to resolve the status of *L. scoparium* forms both here and in Australia.

Habitat:

Abundant from coastal situations to low alpine habitats.

Features:

Decumbent shrub, subshrub, shrub, or small tree up to 5 m in height and in decumbent forms 2-4 m across. Bark light grey to charcoal grey, peeling in long papery flakes, these curling with age. Wood red. Branches numerous erect, spreading or decumbent, arising from base, sometimes sprouting adventitious roots and/or layering on contact with soil. Young branches, young leaves and flower buds densely to sparingly clad in long silky, white hairs. Leaves leathery, pale to dark green, glabrescent to glabrous, linear-filiform, narrowly lanceolate, lanceolate, oblanceolate, to elliptic or obovate (5-)10-15(-20) x 1-2-5(-8) mm, invariably apex drawn out into a long stiff, pungent point, midrib usually distinct sometimes obscure, leaf margin finely crenate, veins simple, scarcely branched. Flowers solitary in leaf axils, (8-)10-20(-25) mm diam. Receptacle dark red, crimson or pink. Petals white, sometimes flushed pink or dark red. Stamens numerous.

Flowering:

Throughout the year

Fruiting:

The capsules are long persistent so invariably mature plants always possess at least some capsules.

Threats:

Not threatened, though some stands are at risk from clearance for farmland or through felling for firewood.

References and further reading:


Gardner, R. 2002. Notes towards an excursion Flora .Manuka *Leptospermum scoparium* myrtaceae. Auckland Botanical Society Journal, 57: 147-149

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=2302



Photographer: © John Braggins



Caption: Flowers of *Leptospermum scoparium* var. *scoparium*

Photographer: Wayne Bennett

Leucopogon fasciculatus

Common Name(s):

Mingimingi, tall mingimingi

Threat Status (2009):

Non Threatened

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=925



Caption: Flowers of *Leucopogon fasciculatus*

Photographer: Wayne Bennett



Caption: *Leucopogon fasciculatus* (Mingimingi)

Photographer: Wayne Bennett

Lophomyrtus bullata

Common Name(s):

Ramarama, bubble leaf

Threat Status (2009):

Non Threatened

Distribution:

Endemic. North and South Islands. Scarce in the South Island where it ranges to about North Canterbury and Greymouth

Habitat:

Coastal to montane forest and shrubland. Often a locally conspicuous component of the understorey of lowland Podocarp riparian forest. *Lophomyrtus bullata* also occasionally grows on in suitable sites in slope forest, and in wetter areas is sometimes a common component of regenerating shrubland in cut over forest. Where it meets with rohutu (*Lophomyrtus obcordata*) the hybrid L. ×ralphii is often commonly found. Sometimes *Lophomyrtus xralphii* is locally dominant occurring in places where ramarama is scarce or has seemingly died out.

Features:

Shrub or tree up to 6 m tall or more. Trunk slender, up to 0.2 m diameter. Bark reddish, fibrous, flaking in small irregular shards, underbark pink. Branches numerous, erect, compactly branched, Branchlets initially 4-angled becoming terete with age, rather brittle, finely hairy, hairs ± persistent. Leaves opposite, coriaceous, finely hirsute when young (hairs somewhat stiffly erect to sericeous, appressed, caducous), maturing glabrous, surface minutely glandular-punctate, oil glands colourless, leaf lamina and petiole decurrent with branchlet; petiole 2-5(-10) mm long, rather brittle; leaf lamina 15-30-(50) × 10-15-(40) mm, broadly ovate to suborbicular, bullate, apex obtuse or acute and then often minutely apiculate, adaxially dark green to yellow green, mottled and/or spotted with red, maroon or purple-black circular blemishes, abaxially pink or red-tinged. Flowers 4-merous, 12-14 mm diameter, borne in axillary, solitary monads, on slender, 12-14(-18) mm long, hirsute pedicels. Hypanthium subturbinate, not extending beyond ovary summit, calyx lobes 4, 1.5-2.2 mm long, persistent, spreading, elliptic-oblong, obtuse to subacute. Petals 8-10 × 6-9 mm, suborbicular, white, margins entire to slightly irregular, ciliate, oil glands colourless. Stamens 80-100(-200 or more), free, in 4 (or more) weakly defined whorls, filaments 8-12 mm long, anthers cream, dorsifixated, latrorse. Ovary inferior, 2-3-locular, ovules numerous, in a single row on each linear placenta. Style 10-12 mm long, slender, white, stigma capitate, scarcely dilated. Fruit a broadly ovoid, dark red or black 6-8 mm long berry. Seeds numerous, reniform, 2.7-5.5 mm diameter, testa dark brown, glossy ± smooth, very hard. Seed description modified from Webb & Simpson (2001).

Flowering:

November - March

Fruiting:

January - June

Threats:

Not Threatened


For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1388



Caption: *Lophomyrtus bullata*

Photographer: Wayne Bennett



Caption: *Lophomyrtus bullata*

Photographer: Wayne Bennett

Melicytus ramiflorus

Common Name(s):

mahoe, whitey wood

Threat Status (2009):

Non Threatened

Distribution:

Endemic subspecies. Three other subspecies occur, one endemic to Norfolk (probably a different species), one to Fiji and one to Samoa. In addition forms from Raoul Island (Kermadec Islands Group) and the Three Kings and eastern Northland may warrant formal recognition. Research into this variation is in progress.

Habitat:

Abundant small tree of coastal, lowland, and lower montane forests throughout the country.

Features:

Shrub or small tree up to 15 m tall. Trunk 1 or more, 0.6-0.8 m diam, typically much branched from near base. Wood soft, white. Bark greyish-white, underbark bright green. Branchlets numerous, twiggy, rather brittle. Petioles 20 mm or more long. Leaves, firmly fleshy, 50-150 x 30-50 mm, light or dark green, lanceolate-oblong to elliptic oblong, apex acute to acuminate (rarely obtuse), leaf margins coarsely serrated (very rarely subentire, or irregularly coarsely toothed). Inflorescence 2-10 flowered fascicles arising from branchlets or leaf axils. Flowers 3-4 mm diam., female or inconstant male (flowers types on separate plants) borne on slender pedicels 5-10 mm long. Bracts subtending flowers, calyx lobes minute, petals greenish-yellow, yellow (rarely cream), lanceolate, apex obtuse. Anthers sessile, stigma 4-6-lobed. Fruit a violet, dark blue or purple berry, 4-5 mm diam., obovoid to globose. Seeds 3-6 per berry.

Flowering:

November - February

Fruiting:

November - March

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=973



Caption: Flower buds. Western Hutt hills. Aug 2013.

Photographer: Jeremy Rolfe



Caption: Carter Scenic Reserve, Wairarapa

Photographer: John Sawyer

Metrosideros robusta

Common Name(s):

Northern rata

Threat Status (2009):

Non Threatened

Distribution:

Endemic. New Zealand: Three Kings Islands, North Island (formerly widespread from Te Paki south to Wellington, now scarce over large parts of this range, and apparently absent from the Hawkes Bay). South Island (abundant from Nelson west and south to Greymouth, from there locally common to about Hokitika, reaching a southern limit just south of Lake Mahinapua. In the east recently recorded from one site near Okiwi Bay, western Marlborough Sounds - though this site is unusual and may not be natural).

Habitat:

Coastal and Lowland forest occasionally extending to montane forest in some parts of the country. Once the co-dominant emergent tree of a distinctive vegetation type called rimu (*Dacrydium cupressinum*)/rata forest.

Features:

Stout tree 25-40 m tall, often starting life as epiphyte, so basal trunk is hollow, and composed of interlocking roots. Trunk 2-3(-4) m diam. Bark firm, persistent, grey-brown, brown or rarely pale yellow, tessellated, shallowly furrowed, somewhat corky. Branchlets numerous, very twiggly (broom-like), puberulent with rust-brown hairs when young. Leaves (excl. water shoots) 25-50(-65) x (10-)15-25(-30) mm, leathery, dark-green, elliptic, ovate-oblong, to rhomboidal, apex obtuse, distinctly notched. Young growth pink, finely covered in rust-brown hairs, becoming glabrescent with age (hairs long persistent on midrib and leaf base). Water shoots - variable shape and size, glabrescent, pale green or yellow-green, delicate and wilting if detached from tree. Inflorescence a broad, terminal corymbiform, cymose, cluster of numerous flowers apically dominated by a temporarily dormant vegetative bud, which recommences growth following flowering. Pedicels 5-8 mm long. Hypanthia obconic, 9 mm long, sepals broad-triangular, petals shedding early, 2 x 3 mm, oblong, dark red, pink, orange or yellow, stamens numerous (25)-30-40 mm long, anthers versatile, pollen dark yellow to orange. Pistil similar length, stigma capitate. Ovary fused to hypanthium, ovules numerous. Capsules oblong 6-9 mm, distinctly raised above sepals and hypanthial rim. Seeds 2.5-5.5 mm, narrowly elliptic to linear, often twisted with apices usually curved or hooked.

Flowering:

(October-) November-January
(-February)

Fruiting:

(December-)January
(-March)

Threats:

Northern rata is most at risk from possum (*Trichosurus vulpecula*) browse. Possums can seriously damage and kill trees, and have, in some situations been directly responsible for the regional loss of northern rata. The species remains common over large parts of range, a situation being improved by the efforts of people encouraged by the national coordination of Project Crimson. Another threat to northern rata comes from hybridization with pohutukawa (*Metrosideros excelsa*) which has now become established well south of its presumed natural southern limits. Ideally people should be discouraged from planting pohutukawa in places it is not natural to, especially when this borders habitats containing northern or southern rata (*Metrosideros umbellata*).


References and further reading:

Allan, H.H. 1961: Flora of New Zealand. Vol. I. Wellington, Government Printer.

Beddie, A.D. 1953. Root behaviour in *Metrosideros*. Wellington Botanical Society Bulletin, 26: 2-6


Report on Northern rata dieback - Minginui faces by Gordon Hosking (DOC Conservation Advisory Science Notes, No. 66, 1994)

Sawyer, J.W.D., Mckessar, K. 2007. Northern rata (*Metrosideros robusta*): a species in decline? Wellington Botanical Society Bulletin, 50: 48-55



Caption: Roots girdling trunk of rimu. Tararua Forest Park. Nov 2012.

Photographer: Jeremy Rolfe



Caption: *Metrosideros robusta*

Photographer: Wayne Bennett

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=977

Myrsine australis

Common Name(s):

Red mapou, red matipo, mapau, red maple

Threat Status (2009):

Non Threatened

Distribution:

Endemic. Three Kings, North, South and Stewart Islands.

Habitat:

Common tree of regenerating and mature forest in coastal to montane situations. Often common on northern offshore islands.

Features:

Shrub or small tree up 6 m tall. Trunk stout, 0.2-0.6 m diam. Bark dark black or purple-black, red on younger branches. Branchlets numerous erect to spreading, very leafy. Petioles stout, fleshy, 5 mm long, often red or green mottled red. Leaves 30-60 x 15-25 mm, dark green to yellow-green variously mottled or blotched with red, or purple spots, leathery, glabrous except for finely pubescent mid vein, obovate-oblong to broad-elliptic, apex obtuse, margins entire, strongly undulate, rarely flat. Inflorescence a fascicle, usually numerous and crowded, produced along branchlets and in leaf axils. Fixed female and inconstant male flowers on different plants, 1.5-2.5 mm diam., white, cream or pale green. Pedicels short, stout, dark red or purple-black. Calyx-lobes 4, sometimes heavily reduced, long persistent. Petals 4, lanceolate, obtuse, free, revolute. Fruit a 1-seeded drupe, 2-3 mm diam., purple-black to black when mature.

Flowering:

August - January

Fruiting:


September - May

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1007



Caption: Male flowers. Rimutaka Forest Park.

Photographer: Jeremy Rolfe



Caption: Male flowers. Rimutaka Forest Park.

Photographer: Jeremy Rolfe

Myrsine salicina

Common Name(s):

Toro

Threat Status (2009):

Non Threatened

Distribution:

Endemic. North, South Islands from Te Paki to about Hokitika

Habitat:

Coastal to montane in forest (rarely shrubland along riversides). On occasion *Myrsine salicina* may form a major part of forest canopy along stream sides.

Features:

Small diffuse to moderately densely branched conical, gynodioecious tree up to 10 m tall. Branches stout, upright, glabrous. Bark firm (not flaking) dark red, maroon-red to almost black. Leaves, fleshy-coriaceous, adaxially dark green, yellow-green or pale pinkish-green, usually blemished with maroon spots, abaxially pink to wine-red or pale green, blemising on adaxial surface apparent on abaxial surface, margins entire, flat or very slightly recurved, midrib deeply impressed adaxially, prominent ridged abaxially (side veins not evident when fresh); petioles 10-14 mm long, fleshy, stout, flattened. Lamina 70-180 × 20-30 mm, narrow-elliptic, narrow-oblong, to linear-oblong, apex obtuse, base attenuate to cuneately narrowed (gradually tapering to base). Inflorescences in ± dense 10-15(-20)-flowered fascicles.

Pistillate flowers; greenish yellow to cream with maroon spotting or wine-red with purple-black spotting; calyx 1.3-1.9 mm, tube 0.2-0.6 mm, lobes 4-5, 0.7-1.0 x 0.6-0.8 mm, oblong to ± triangular, apex acute to subacute, margins minutely ciliolate; corolla 2.8-4.2 mm, tube 0.2-0.3 mm, lobes 4-5(-6), 2.0-2.4 x 1.0 mm, elliptic, apex acute. Antherodes malformed, 0.82-1.10 x 0.5-0.6 mm, apiculus strongly recurved; pollen absent. Ovary 1 x 1 mm. Stigma 0.30-0.48 mm high, spreading, outer parts appressed to ovary ± 2.5 mm diameter.

Bisexual flowers with the same colouration; calyx 1.6-2.0 mm, tube 0.4-0.7 mm, lobes 4-5, 0.7-1.1 x 0.6-0.9 mm, oblong, apex acute, margins minutely ciliolate. Corolla 3.0-4.2 mm, tube 0.3-0.6 mm, lobes 4-5, 2.6-2.8 x 1.0-1.4 mm, elliptic, apex acute. Anthers 1.1-1.8 x 0.8-1.2 mm, apiculus upright; pollen white. Ovary 0.7-0.9 x 0.8-1.0 mm. Stigma 0.8-0.85 mm high, upright. Drupe (5-)8-9 mm long, obovoid, flesh red to orange (rarely maroon), on pedicels 8-10 mm long. Endocarp 5.0-6.7 × 3.5-4.5 mm, obovate to broadly obovate, dull, buff to buff brown, orange-brown or henna, bearing 1(-2) seeds.

Flowering:

August - January

Fruiting:

September - May

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1011



Caption: Ripe fruit, Whanganui Inlet, North West Nelson

Photographer: Simon Walls



Caption: Pinehaven. Jan 2005.

Photographer: Jeremy Rolfe

Nestegis cunninghamii

Common Name(s):

Black maire

Threat Status (2009):

Non Threatened

Distribution:

Endemic. North, South, Hauturu (Little Barrier) and Aotea (Great Barrier) Islands. Scarce north of Auckland and often uncommon over wide parts of its former lowland range due to it being preferentially logged by early settlers. In the North Island Black Maire is now most commonly seen in the main Axial Ranges and Central Volcanic Plateau. In the South Island extending to Napenape in the east and near Greymouth in the west.

Habitat:

Widespread in coastal to montane forest. Often prominent in riparian Podocarp forest and on the Podocarp forests developed on the ignimbrite and pumice country of the Central Volcanic Plateau. As a rule Black Maire seems to prefer more frost prone habitats than White Maire (*Nestegis lanceolata*) though both species often grow together. Black maire is also common host for white mistletoe (*Tupeia antarctica*) in the Central Volcanic Plateau, Hawkes Bay and Wairarapa.

Features:

Stout gynodioecious spreading tree up to 25 m tall usually forming a broadly domed canopy; trunk up to 1·5m diameter, sometimes with several arising from base, usually straight and arching, sometimes twisted; bark firm (not flaking), grey-brown to dark brown, tessellated. Branches spreading, branchlets minutely pubescent. Leaves glabrous, coriaceous, dark green to brown-green above, ± dull, paler beneath, margins plane, entire with prominent raised midrib and side veins, borne on rigidly stout petioles 9-10(-15) mm long; lamina of juveniles 100-300 × 5-10 mm, narrow-linear, apex acute, base cuneately narrowed or attenuate; adult lamina 150-400(-600) × 20-40 mm, lanceolate to ovate- or elliptic-lanceolate, obtuse or subacute, coriaceous, weakly bullate and somewhat rough to touch; midrib and side veins impressed above (prominent below). Inflorescence a stout 8-12(-20)-flowered raceme 10-25 mm long; rhachis and pedicels densely pubescent. Male flowers with 2 large exserted anthers, ovary mostly rudimentary (occasionally functional); female flower with 2 sessile barren anthers, ovary with large 2-lobed stigma. Drupe 15-20 mm long, ovoid, flesh red, orange-red to purple-black; endocarp 10-18 × 6-9 mm, dull, pale orange-yellow, narrowly elliptic, elliptic to broadly elliptic, terete (sometimes weakly compressed). Seed purple-brown. Description adapted from Allan (1961) and Webb & Simpson (2001).

Flowering:

October - November

Fruiting:

December - April

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1032



Caption: *Nestegis cunninghamii* - close up of upper leaf surface

Photographer: Peter de Lange



Caption: *Nestegis cunninghamii* - leaf underside showing distinctive venation

Photographer: Peter de Lange

Nothofagus solandri var. *solandri*

Common Name(s):

Black beech

Threat Status (2009):

Non Threatened

Distribution:

Endemic. North, South Islands. Very rare north of the Central Volcanic Plateau and East Cape. Little Barrier Island appears to be the current northern limit.

Habitat:

Lowland to montane forest. At times the canopy dominant and forming its own distinctive forest type.

Features:

Tree up to 25 m tall, wood pinkish to yellowish when fresh, often with darker patches. Bark rough, furrowed, charcoal black. Trunk 1 m or more diam. Branches numerous, stout, spreading, "pagodaform". Petioles 1-2 mm. Leaves 10-15 x 5-10 mm, narrow- to elliptic-oblong, obtuse, obliquely cuneate at base, apex often apiculate, leathery, glabrous, dark green above, undersides clad in greyish white tomentum, venation distinct on both surfaces. Domatia absent. Male inflorescences 1-4 per branchlet, on short sparsely pubescent peduncles; flowers 1-2, sessile. Perianth broadly-campanulate, 2 x 3 mm, shallowly and obtusely 4-5-lobed; stamens 8-17, anthers 2-3 mm long, dark red. Female inflorescences ovoid, 1-2 per branchlet, pubescent to pilose hairy, sessile, flowers 1-3. Lateral flowers trimerous, terminal ones dimerous; stigmas clavate. Cupule 6-7 mm, glabrous or pubescent, 3-partite. Nuts up to 7 mm long, wings broad at base, narrowed to apex.

Flowering:

September - December

Fruiting:

November - April

Threats:


Not Threatened

References and further reading:

Anonymous. 1957. Construction of key for the genus *Nothofagus*. Auckland Botanical Society Journal, 14: 2-3


For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1038



Caption: Lower Hutt. Jul 2013.

Photographer: Jeremy Rolfe



Caption: in cultivation

Photographer: Jesse Bythell

Nothofagus truncata

Common Name(s):

Hard beech

Threat Status (2009):

Non Threatened

Threats:

Not Threatened

References and further reading:

Anonymous. 1946. Note on *Nothofagus truncata*. Auckland Botanical Society Journal, 3: 5-6

Anonymous. 1957. Construction of key for the genus *Nothofagus*. Auckland Botanical Society Journal, 14: 2-3


Cameron, E.K. 1997. Distribution of beech in the Waitakere Ranges. Auckland Botanical Society Journal, 52: 68-72.

Greenwood, R.M. 1959. Hard beech in the Tararuas. Wellington Botanical Society Bulletin, 31: 15-18

Rogan, D. 1999. New northern limit for hard beech *Nothofagus truncata*. Auckland Botanical Society Journal, 54: 1

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1039



Caption: Maidstone Park, Upper Hutt. May 2013.

Photographer: Jeremy Rolfe



Caption: Saplings in understorey, Maidstone Park, Upper Hutt.

Photographer: Jeremy Rolfe

Olearia rani var. colorata

Common Name(s):

Heketara

Threat Status (2009):

Non Threatened

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1059



Caption: Wellington. Nov 1993.

Photographer: Jeremy Rolfe



Caption: Puccinia atkinsonii rust
on *Olearia rani var. colorata*.
Rotoehu. Oct 2007.

Photographer: John Hobbs

Piper excelsum subsp. *excelsum*

Common Name(s):

kawakawa, pepper tree

Threat Status (2009):

Non Threatened

Distribution:

Endemic. North and South Islands. Common from te Paki south to about Okarito, North Canterbury and Banks Peninsula.

Habitat:

Coastal to lowland (extending up 500 m a.s.l. in warmer parts of the country). Usually an important understorey species in coastal forest.

Features:

Small tree to at least 5 m tall; stems erect (occasionally layering), not notably lenticellate, new shoots red-green or green (leaf nerves, petioles and new sterns with reddish colouring), taste peppery; pith of axes (including rachis of spike) without a mucilage core. Prophyll a collar to 0.3 (-2.2) mm high. Leaf blades submembranous, orbicular, suborbicular, at vegetative nodes to 100(-120) mm diameter, usually with 5-8 principal nerves, cordate at base, with a very narrow or closed sinus, occasionally basal lobes overlapping, upper surface of blade not bullate; petiole to 40(-60) mm long, c.0.4× as long as blade, the sheath 0.3-1.0(-2)× as long as non-sheathing part, truncate-rounded at apex and not produced there, the non-sheathing part of petiole to 4.0 mm diameter. Inflorescences solitary or 2-3 together on a short (rarely more than 10 mm long) axillary shoot, and (usually solitary) on the adjacent terminal shoot (occasionally this shoot not fertile); reduced leaf at apex of fertile shoot with a glabrous petiole and usually with a green oblong lamina at least 5 mm long, but lamina often ± lacking, especially on terminal fertile shoot. Female inflorescence erect in flowering and remaining so into fruit, peduncle to c. 1.5 cm long, spike to 60(-100) × c.6 mm diameter, with uniseriate usually 5-10-cellular hairs to 0.15 mm long on lower part of bract stalks and sparingly on rachis, these hairs not obvious on the peduncle just below the lowermost bracts; bracts peltate, bract heads 0.40-0.75 mm diameter; flowers at full emergence centred c.1.3 mm apart, emergent part of ovary ovoid; stigmas 3-4(-5), together c. 1.2 mm diameter. Male inflorescence erect, spike to c.110 mm long, proximally c.6 mm diameter, bracts and hairs as in female inflorescence; staminal filaments c. 0.25 mm long, anthers c.1.00 × 0.75 mm wide. Ripe infructescence c.10 mm diameter; fruitlets coalescent, sunken apically about the persistent dark stigmas, exocarp and mesocarp orange; seed oblong to slightly obovoid, apiculate at apex, c.2.0 × 1.5 dark brown, with (3-)4-5(-7) broad longitudinal furrows.

Flowering:

August - November

Fruiting:

Throughout the year

Threats:

Not Threatened

References and further reading:

de Lange, P.J. 2012: Taxonomic notes on the New Zealand flora: new names in *Piper* (Piperaceae). *New Zealand Journal of Botany* DOI:10.1080/0028825X.2012.708904

Gardner, R.O. 1997: *Macropiper* (Piperaceae) in the south-west Pacific. *New Zealand Journal of Botany* 35: 293-307.

Jaramillo, M.A.; Callejas, R; Davidson, C.; Smith, J.F.; Stevens, A.C.; Tepe, E.J. 2008: A phylogeny of the tropical genus *Piper* using ITS and the chloroplast intron psbJ-petA. *Systematic Botany* 33: 647-660.



Caption: Male inflorescence.
Tamahunga, Omaha.

Photographer: Jeremy Rolfe



Caption: Male inflorescence.
Tamahunga, Omaha.

Photographer: Jeremy Rolfe

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=964

Pittosporum divaricatum

Threat Status (2009):


Non Threatened

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1134



Caption: Sugarloaf, Canterbury
(leaf detail)

Photographer: Jesse Bythell



Caption: Cobb valley, November

Photographer: John Smith-Dodsworth

Pittosporum eugenioides

Common Name(s):

Tarata, lemonwood

Threat Status (2009):

Non Threatened

Distribution:

Endemic. Common in the North and South Islands.

Habitat:

Common tree of regenerating and mature forest in coastal to montane situations.

Features:

Gynodioecious tree up to 12 m tall but usually much less. Trunk 0.6-1 m diam, stout, clad in persistent pale-grey bark, branches numerous, erect then spreading. Leaf buds sticky, resinous. Leaves borne on slender petioles 10-20 mm long, alternate, 50-100(-150) x 25-40 mm, yellow-green, green, more or less blotched and mottled with paler green or yellow-green (sometimes white), somewhat leathery, glossy, smelling strongly when crushed of ivy or resin, elliptic to elliptic-oblong, apex acute to subacute; leaf margin undulate (very rarely not so), midrib pale green. Inflorescences terminal, numerous, subcorymbose compound umbels. Flowers pale yellow to yellow, very fragrant. Peduncles 10-20 mm, pedicels 5 mm, both sparsely hairy. Sepals 2 mm, ovate to narrow-ovate, pale caducous. Petals 5, 5-7 mm long, narrow-oblong. Capsules 2-valved (rarely 3), 5-6 mm, ovoid to elliptic, caducous, seeds immersed in dark yellow viscid pulp, whole structure covered in long persistent papery endocarp.

Flowering:

October - December

Fruiting:

October - January

Threats:

Not Threatened

References and further reading:

Cooper, R.C. 1956: The Australian and New Zealand species of *Pittosporum*. Annals of the Missouri Botanical Garden 43: 87-188

Gardner, R. 1999. Notes towards an excursion Flora. *Pittosporum eugenioides* as a wild plant. Auckland Botanical Society Journal, 54, 1


For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1135



Caption: Masterton

Photographer: John Barkla



Caption: Maidstone Park, Upper Hutt.

Photographer: Jeremy Rolfe

Pittosporum tenuifolium

Common Name(s):

Kohukohu, black matipo

Threat Status (2009):

Non Threatened

Distribution:

Endemic and widespread throughout country.

Habitat:

A small tree of coastal to montane shrubland and forested habitats. Preferring successional habitats.

Features:

Shrub or small gynodioecious tree up to 10 m tall (usually much less). Trunk 0.3-0.4(-0.6) m diam., stout, clad in dark grey-black or brown persistent bark. Branches numerous, erect then spreading. Branchlets and young leaves pubescent, hairs pale yellow or cream. Petioles short, somewhat fleshy. Leaves alternate, (10-)30(-70) x (5-)10(-20) mm, leathery, pale-green to dark green above, lighter below, oblong, oblong-ovate or elliptic-obovate, apex obtuse to acute, rarely acuminate, margins entire, often undulose. Flowers solitary or in axillary cymes, rather fragrant, especially at night. Pedicels stout, pale green, fleshy, bracts entire, lanceolate, caducous. Sepals narrowly ovate-oblong, subacute to obtuse, silky hairy. Petals 12 mm long, lanceolate, dark red, black (rarely yellow or white). Capsules 2-valved (rarely 3), subglobose, valves woody, black when mature, long persistent. Seeds immersed in sticky, red or yellow viscid pulp.

Flowering:

October - November (-December)

Fruiting:

January - March

Threats:

Not Threatened

References and further reading:

Allan, H.H. 1961: Flora of New Zealand. Vol. I. Wellington, Government Printer

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1139



Caption: Pittosporum tenuifolium in flower Dunedin

Photographer: John Barkla



Caption: Quail Island

Photographer: John Barkla

Plagianthus regius subsp. *regius*

Common Name(s):

Manatu, ribbonwood, lowland ribbonwood

Threat Status (2009):

Non Threatened

Distribution:

Endemic. New Zealand: North, South and Stewart Islands

Habitat:

Coastal to lower montane. Often a prominent tree in lowland alluvial forest.

Flowering:

September - November

Threats:


Not Threatened

References and further reading:

Wilcox, M.D. 2002. Lowland ribbonwood *Plagianthus regius* at Clevedon. Auckland Botanical Society Journal, 57: 144-146

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1142



Caption: Bark, Dunedin

Photographer: John Barkla



Caption: Woodhaugh, Dunedin

Photographer: John Barkla

Podocarpus cunninghamii

Common Name(s):

Mountain totara, Hall's totara, thin-barked totara, totara-kiri-kotukutuku

Threat Status (2009):

Non Threatened

Distribution:

Endemic. North, South and Stewart Islands.

Habitat:

Lowland, montane to lower subalpine forest (but notable more common in montane forest).

Features:

Robust dioecious conifer up to 20 m tall. Trunk stout, 1-1.5 m diam., clad in papery, thin, freely flaking reddish-grey bark. Trunk without branches at base, branches slender, erect, spreading or somewhat drooping. Leaf bud significantly broader than the diam., of the branchlet, surrounded by caducous, papery, ovate bracts. Leaves yellow-green, green, or brownish-green, erect, leathery; juvenile 25-50 x 4-5 mm, adults 20-30 x 3-4 mm., narrow-linear to linear-lanceolate, acute to acuminate, apex very pungent, mid-vein distinct. Male cones (strobili) axillary, 10-25 mm, solitary or up to 5 on a common peduncle. Female branchlets axillary, ovules solitary or paired. Receptacle of 2-4 scales, irregularly elliptic-oblong to obovate-oblong, maturing as a red, swollen, succulent, sweet tasting "fruit" this surmounted by a 1(-2) elliptic, elliptic-oblong or ovate-oblong, (5-)6.5-8.5 mm long, grey nut brown or dark brown (green when fresh) seed.

Fruiting:

Flowering:

(August-) October (- December)

Fruits take a year or so to ripen, and may be found throughout the year, usually peaking at about the same time that cones are produced. They are most frequently seen between April and May

Threats:

Not Threatened

References and further reading:

Allan, H.H. 1961: Flora of New Zealand. Wellington, Government Printer.

Connor, H.E.; Edgar, E. 1987: Name changes and Nomina Nova IV. New Zealand Journal of Botany 25:

de Lange, P.J.; Rolfe, J.R. 2010: New Zealand Indigenous Vascular Plant Checklist. Wellington, New Zealand Plant Conservation Network. 164pp.

Gardner, R. 1990. Totara and Halls totara. Auckland Botanical Society Journal, 45: 27-28.

Kirk, T. 1889: The Forest Flora of New Zealand. Wellington, Government Printer.

Molloy, B.P.J. 1985: The continuing saga of native conifer nomenclature. DSIR Botany Division Newsletter 102: 26-27.

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1174



Caption: Seeds of *Podocarpus cunninghamii*

Photographer: Wayne Bennett



Caption: *Podocarpus cunninghamii*

Photographer: Wayne Bennett

Podocarpus totara var. *totara*

Common Name(s):

Totara

Threat Status (2009):

Non Threatened

Distribution:

Endemic. Common throughout most of the North and South Islands. Present but extremely scarce on Stewart Island (Freshwater River).

Habitat:

Widespread and at times abundant tree of lowland, montane and lower subalpine forest. May also form a vegetation type in which it is the dominant species.

Features:

Robust dioecious conifer up to 30 m tall. Trunk stout, 2-3 m diam., clad in thick, corky, furrowed and somewhat stringy reddish-grey bark. Trunk without branches at base, branches stout, erect to spreading. Leaf bud narrower than or the same diam., as branchlet, surrounded by caducous, papery, narrowly lanceolate bracts. Leaves brownish-green, erect, leathery; juvenile 20 x 1-2 mm, adults 15-30 x 3-4 mm., linear-lanceolate, acute, apex pungent, mid-vein distinct to obscure. Male cones (strobili) axillary 10-15 mm, solitary or in 4s. Female branchlets axillary, ovules solitary or paired, receptacle of 2-4 scales, acute and free at tips, maturing as a red, swollen, succulent, sweet tasting "fruit" this surmounted by a 1(-2) broadly elliptic, ovoid-oblong 3-6 mm, semi-glossy, buff, grey nut brown, henna or dark brown (green to glaucous-green) when fresh, seed.

Fruiting:

Flowering:
(August-) October (- December)

Fruits take a year or so to ripen, and may be found throughout the year, usually peaking at about the same time that cones are produced. They are most frequently seen between April and May

Threats:

Not Threatened, though as a vegetation type it is all but extinct throughout most of its former range.

References and further reading:

Gardner, R. 1990. Totara and Halls totara. Auckland Botanical Society Journal, 45:27-28.

Moorfield, J. C. (2005). Te aka : Maori-English, English-Maori dictionary and index. Pearson Longman: Auckland, N.Z.

Landcare Research. Nga Tipu Whakaoranga - Maori Plant Use Database.
<http://maoriplantuse.landcareresearch.co.nz/WebForms/default.aspx>

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1176



Caption: *Podocarpus totara* var. *totara* at Pokemokemoke

Photographer: Wayne Bennett



Caption: Seeds of *Podocarpus totara* var. *totara*

Photographer: Wayne Bennett

Prumnopitys ferruginea

Common Name(s):

Miro, brown pine

Threat Status (2009):

Non Threatened

Distribution:

Endemic. North, South and Stewart Islands.

Habitat:

Common tree of lowland to montane forest.

Features:

Stout tree up to 25 m tall. Trunk 1-1.5 m diam., in adults clear of branches for 2/3 of length. Bark thick, grey. Falling in thick, sinuous flakes. Leaves feathery, dark green, green to bronze-green, distichous, erect, narrow-linear, acute, falcate to subfalcate, acute to subacute, mid vein distinct, margins recurved, juveniles up to 30 mm long, those of adults 15-25 x 2-3 mm. Male cones (strobili) solitary, axillary, 5-15 mm long. Ovules solitary (rarely paired), on short branchlets 10 or less mm. long. Fruit a broadly oblong to sub-spherical red, pink-red fleshy drupe up to 20 mm long - fleshy, oily, smelling and tasting strongly of terpenes. Stone elliptic to broadly elliptic 11-17 mm long, dark brown to black-brown.

Flowering:

June - August
- October

Fruiting:

Fruits take 12-18 months to mature.
Ripe fruits are mainly found from
November - April



Caption: Ruahine Range

Photographer: John Sawyer



Caption: Seeds of *Prumnopitys ferruginea*

Photographer: Wayne Bennett

Threats:

Not Threatened, although as a forest-type it has been greatly reduced through widespread logging. Very few intact examples of miro-dominated forest remain in the country.

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=794

Pseudopanax arboreus

Common Name(s):

Fivefinger, five finger, whauwhaupaku

Threat Status (2009):

Non Threatened

Distribution:

Endemic. Widespread (though rare in Central Otago). North and South Islands

Habitat:

Coastal to montane (10-750 m a.s.l.). Moist broadleaf forest. Frequently epiphytic. A frequent component of secondary forest. Streamsides and forest margins.

Features:

Us. Dioecious. Small multi-branched tree to 8 m tall, branches and branchlets brittle. Leaves alternate, leaflets 5-7 (us. 5), palmate. Petioles c. 15-20 cm long, sheathing branchlet at base. Petiolules c. 3-5 cm long, pale green. Leaflets obovate-oblong to oblong-cuneate, thinly coriaceous, coarsely serrate-dentate, acute or acuminate to obtuse; midveins and main lateral veins obvious above and below; terminal lamina 10-20 x 4-7 cm. Inflorescence and panicle, terminal, compound; flowers usually unisexual; 8-20 primary rays (branchlets), up to 10 cm long; 15-20 secondary rays; umbellules with 10-15 flowers in each. Calyx truncate or obscurely 5-toothed; flowers c. 5 mm diam., sweet-scented; petals 5, white to pink flushed, ovate to triangular, acute; stamens 5, obvious, filaments c. = petals; ovary 2-loculed, each containing 1(-2) ovules; style branches 2, spreading. Fruit fleshy, 5-8 mm diam., style branches retained on an apical disc, very dark purple, laterally compressed. Seeds 2(-3) per fruit, wrinkled, 3-6 mm long.

Flowering:

June to August

Fruiting:


August to February

Threats:

Not Threatened. In places the petiolules of *Pseudopanax arboreus* (and other fleshy-leaved *Pseudopanax* species) are a conspicuous element of possum (*Trichosurus vulpecula*) diet and the forest floor can become littered with discarded leaflets.

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1194



Caption: *Pseudopanax arboreus*

Photographer: Wayne Bennett



Caption: Flowers of *Pseudopanax arboreus*

Photographer: Wayne Bennett

Pseudopanax crassifolius

Common Name(s):

Horoeka, lancewood

Threat Status (2009):

Non Threatened

Distribution:

Endemic. North, South and Stewart Islands. Widespread and common

Habitat:

Lowland to montane forest. Sealevel to c. 750 m a.s.l.

Features:

Bushy topped tree to 15 m tall, branchlets fleshy, trunk us. unbranched in lower part, to 50 cm diam., distinctly ridged when young, bark dark becoming paler with age, wood tough. Leaves alternate; leaflets 1-3 in seedling, palmate, sessile or subsessile on very short petiolule, submembranous coarsely toothed, absent from juvenile and adult. Juvenile leaves dark green, narrow-linear, deflexed, to 1 m long, coriaceous, midrib pale cream-yellow, raised, margins distantly sharply toothed, distal margin of tooth perpendicular to midvein, not swollen. Adult leaves shorter, 10-20 x 2-3 cm, dark green, very occ. trifoliate (probably due to hybridisation with oither species), narrow elliptic-cuneate to lanceolate or linear-obovate, acute or obtuse, margins entire to sunuate or coarsely serrate, subsessile or on petioles to 10 mm long, petiole base expanded around stem. Inflorescence a terminal umbel, irregularly compound; primary rays (branchlets) 5-10, c. 6 cm long; umbellules sometimes racemously arranged. Ovary 5-loculed, each containing 1 ovule; style branches 5, connate, tips sometimes free. Fruit fleshy, subglobose, 4-5 mm diam., style branches retained on an apical disc, dark purple when ripe. Seeds 4-5 per fruit, easily separated, broadly ovate, grooved, 2.2-3.5(-5.5) mm long.

Flowering:

January-April

Fruiting:


January-April

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1196



Caption: *Pseudopanax crassifolius*

Photographer: Wayne Bennett



Caption: Seeds of *Pseudopanax crassifolius*

Photographer: Wayne Bennett

Pseudowintera axillaris

Common Name(s):

Lowland horopito, lowland pepper tree

Threat Status (2009):

Non Threatened

Distribution:

Endemic. North and South Islands. Scarce north of Auckland, extending to near Westport on the west of the South Island

Habitat:

Lowland to montane forest. From near sea level.

Features:

Shrub or small tree up to 7 m tall; trunks and branches upright; bark dark; branchlets dark. Plants glabrous. Petiole slender, 2 cm long, dark reddish brown. Leaves alternate, pungent, 6-10 x 3-6 cm, upper surface glossy, dark green to yellowish-green, without blemishes or blotches; midvein conspicuous, pale; lamina elliptic-oblong to elliptic-obovate to ovate-oblong to lanceolate rhomboid, margin undulate, obtuse to subacute, coriaceous; undersides pale to glaucous but not white; midvein pale. Inflorescences axillary, flowers bisexual, c. 1 cm diam., in fascicles of 1-5-(10), on slender pedicels 5-10 mm long. Calyx cupule margins sub-entire to shallowly lobed. Corolla comprised of 5-(6) free petals, these 5-6 mm long, narrow-oblong to narrow-obovate, greenish yellow, apex obtuse. Carpels 1-6, stigma apical. Stamens 6-20. Fruit a 3-6-seeded fleshyglobose to subglobose berry, 5-6 mm diam., orange to orange-red. Seed 3-angled, obovate to elliptic, 2.5-3.9 mm, surface irregular (showing striping under very high magnification).

Flowering:

September-December

Fruiting:

October-January (some fruit may be retained until June)

Threats:

Not Threatened. Unpalatable to browsers

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=787



Caption: Western Hutt hills. Aug 2013.

Photographer: Jeremy Rolfe



Caption: Herepai Ridge, Tararua Forest Park.

Photographer: Jeremy Rolfe

Rhopalostylis sapida

Common Name(s):

Nikau palm

Threat Status (2009):

Non Threatened

Distribution:

Endemic. North Island, South Island from Marlborough Sounds and Nelson south to Okarito in the west and Banks Peninsula in the east. Also on Chatham and Pitt Islands. However Chatham Islands plants have a distinct juvenile form, larger fruits, and thicker indumentum on the fronds.

Habitat:

Primarily a species of coastal to lowland forest in the warmer parts of New Zealand.

Features:

Trunk up to 15 m, stout, covered in grey-green leaf scars, otherwise green. Crownshaft 0.6(-1) m long, dark green, smooth, bulging. Fronds up to 3 m long; leaflets to 1 m, closely set (sometimes over lapping), ascending. Spathes c.300 x 150 mm., between pink and yellow, caducous. Inflorescence shortly stalked, with many branches, 200-400 mm long. Flowers sessile, unisexual, tightly packed, lilac to pink. Males in pairs, caducous, stamens 6. Females solitary, with minute staminodes, ovary 1-locular, stigmas terminal, recurved, persistent. Fruit c.10 x 7 mm, elliptic-oblong, flesh red.

Flowering:

November - April

Fruiting:

February - November

Threats:

Not Threatened

References and further reading:

Esler, A.E. 1969. Leaf fall and flowering of nikau. Wellington Botanical Society Bulletin, 36: 19-22

Greenwood, R.M. 1969. Notes on growth of young nikau plants. Wellington Botanical Society Bulletin, 36: 22-23


For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1259



Caption: Northland

Photographer: John Barkla



Caption: Rhopalostylis sapida

Photographer: Pat Enright

Schefflera digitata

Common Name(s):

Patete, pate, seven-finger

Threat Status (2009):

Non Threatened

Distribution:

Endemic. Widespread. North, South and Stewart Islands.

Habitat:

Lowland to montane forest (sealevel to 1000 m a.s.l.).

Features:

Dioecious(?) small tree to 8 m. Trunk irregularly branched; bark greenish, finely ridged and with scattered prominent lenticels. Petioles terete, to 25 cm long, sheathing branchlet, reddish. Petiolules to 2 cm, reddish. Leaves alternate, palmate, with (3)-10 leaflets (us. 7), upper surface evenly green in adult, underside pale, shiny, purplish in juvenile. Terminal leaflet to 20 cm long; lateral leaflets decreasing in size; obovate-cuneate, tip acuminate to obtuse; margins sharply serrate in adult, irregularly lobed to pinnatifid in juvenile. Inflorescence a panicle, axillary (occ. cauline), branches many, spreading, to 35 cm; bracts and bactlets small. Umbels many, up to 10 flowers in each; peduncles subsessile to 10 mm long, pedicels shorter. Flowers greenish cream, c. 7 mm diam. Petals 5(-6), acute. Stamens 5, filaments c. = petals. Style branches 5 (or more), connate below forming an irregular disc. Fruit subglobose,c. 3.5 mm diam., fleshy, dark purple when ripe, containing (5-)7-10(-11) seeds. Seed 2-2.5 mm.

Flowering:

February-March

Fruiting:

February-March

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1281



Caption: Waipoua Forest, Northland

Photographer: John Sawyer



Caption: Schefflera digitata (Patete)

Photographer: Wayne Bennett

Syzygium maire

Common Name(s):

swamp maire, maire tawake, waiwaka

Threat Status (2009):

Non Threatened

Distribution:

Endemic. North and South Island from Te Paki south to Rarangi (near Blenheim). Now often scarce or absent over large parts of its former range due to the clearance of swamp forest.

Habitat:

Mostly found in coastal and lowland riparian forest in waterlogged ground, on the margins of swamps and streamsides. Also found in some of montane forest and cloud forest of Northland (e.g., Tutamoe) and the western Waikato (Pirongia, Taumatatotara and Tawarau) where high rainfall and poor drainage provide ideal conditions for this tree to establish on hill slopes, tablelands and with karst landscapes.

Features:

Glabrous tree to c.16 m high. Trunk up to 0.8 m dbh, solitary or with several arising from base, often with knees and where the root plate is exposed frequently bearing pneumatophores. Bark smooth, pinkish grey, grey-brown or white, flaking in soft or brittle, irregular shards. branches numerous, spreading, branchlets numerous, spreading, 4-angled. Leaves opposite, subcoriaceous, adaxially yellow-green to green, glossy often bearing small galls and leaf blisters, midrib impressed, side veins slightly impressed scarcely evident when viewed from above; abaxial surface pale green, midrib prominently raised, side veins evident when fresh or dried; margins entire, sinuate or undulate; petioles 5-10 mm long, slender, brittle. Lamina 15-60 × 10-25 mm, usually elliptic, sometimes broadly elliptic. Inflorescences in cymose 5-30-flowered clusters, up to 100 mm diameter. Pseudopedicels slender. Hypanthium 2-3 mm long at anthesis, obconic; calyx lobes very short and broad, persistent on fruit. Petals 2-3 mm diameter, orbicular, white, forming calyptrum in bud, caducous. Stamens numerous, 5-12(-18) mm long, white, in 6-8 (or more) indistinct whorls, filaments 4.5-17.5 mm long, white, anthers basifix, pollen white. Style 5-18 mm long, distinctly broader than stamens and tapering, cream to yellow-green. Ovary adnate to base of hypanthium. Fruit 10-15 mm diameter, subglobose, broad-ellipsoid or elliptic-ovoid, flesh deep crimson, glossy. Seed 1, 6-11 mm long, obovate, testa dull, very hard, covered in fibres, striped pale orange-yellow and pale brown, brown or grey-brown.

Flowering:

November - July

Fruiting:

January - December

Threats:

Not Threatened. However, many populations now qualify as "Living Dead" as they persist (and are in slow terminal decline) as remnants within partially drained farmland (previously riparian forest). In some parts of its range it is listed as regionally threatened, e.g., Auckland and Wellington.

References and further reading:

Cameron, E.K., Cutting, M. 1995. Maire tawake at Browns bay Auckland. Auckland Botanical Society Journal, 50: 66-70.

Webb, C.J.; Simpson, M.J.A. 2011: Seeds of New Zealand Gymnosperms and Dicotyledons. Christchurch, Manuka Press.

Webb, C. J.; Sykes, W. R.; Garnock-Jones, P. J. 1988: *Flora of New Zealand. Vol. IV. Naturalised Pteridophytes, Gymnosperms, Dicotyledons*. Christchurch, New Zealand, Botany Division, D.S.I.R.



Caption: *Syzygium maire*

Photographer: Wayne Bennett



Caption: Flower of *Syzygium maire*

Photographer: Wayne Bennett

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1313

Weinmannia racemosa

Common Name(s):

kamahi

Threat Status (2009):

Non Threatened

Distribution:

Endemic. North, South, Stewart Island. The exact northern limits of *Weinmannia racemosa* are uncertain but probably lie somewhere along the Manukau Harbour and Hunua Ranges across the Kamai Range. North of here the distinction between *Weinmannia racemosa* and *W. silvicola* is often confused. This needs further study.

Habitat:

Coastal to subalpine. A widespread and common tree of disturbed habitats in coastal and lowland to montane forest, often becoming locally dominant in higher altitude montane forest in the higher ranges of the North Island and western South Island.

Features:

Tree up to 28 m tall often forming a narrowly domed canopy (but this will vary according to local conditions). Trunk up to 1.2 m diameter. Branches numerous, erect to spreading, Foliage heterophyllous, with distinct seedling, juvenile and adult leaves (reversion shoots common). Stipules caducous, 3-6 mm long, lanceolate, finely pubescent, yellow-green to pinkish. Seedling and juvenile leaves membranous to subcoriaceous, 10-60 × 10-30 mm; lamina simple to 3-lobed or 3-foliate, ovate-elliptic to elliptic or lanceolate, apices subacute to acute, margins serrate to incised-serrate; adult leaves coriaceous, on petioles up to 20 cm long, lamina 30-100 × 20-40 mm lamina simple, elliptic, ovate-elliptic to broad-ovate, apices obtuse to subacute, margins rather coarsely, bluntly serrate. Inflorescences in racemose; racemes 60-1140 mm long, rachises and pedicels finely, pilose-pubescent; pedicels 2-4 mm long, clustered, ascending to spreading. Sepals 1.0-1.5 mm. long, ovate, persistent; petals 4(-5), 2-3 mm long, ovate-oblong, white, cream or pale pink; stamens 8-10, exserted, filaments up 10 mm long, white or pinkish white, anthers 0.2-0.3 mm diameter, cream; nectaries 8, red; ovary, narrowly ovoid 0.8 mm diameter, covered in appressed hairs, carpels 2, free almost to base. Styles 3-4 mm long, pale pink, persistent; stigma 0.2-0.4 mm, pink or pale pink, punctate. Fruit a pubescent, broadly cylindrical capsule 4.0-5.8 × 2.7-3.1 mm, initially greyish drying honey-brown or dark brown. Seeds numerous, 1.0-1.5 mm long, narrowly elliptic to elliptic-oblong, orange-brown, apices bearing dense hair tufts, otherwise glabrous.

Flowering:

July - January

Fruiting:

October - May

Threats:


Not Threatened

References and further reading:

McKenzie, R. 1960. The distributional overlap of *Weinmannia silvicola* and *Weinmannia racemosa*. Auckland Botanical Society Journal, 17: 7-8

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1368



Caption: Tararua Range. Feb 2008.

Photographer: Jeremy Rolfe



Caption: Mt Pirongia

Photographer: Gillian Crowcroft

Definitions of botanical terms

A glossary has been provided below with definitions for many of the botanical terms used in the species descriptions.

Glossary

Term	Definition
Abaxial	Facing away from the stem of a plant (especially denoting the lower surface of a leaf).
Acerose	Narrow with a sharp stiff point.
Achene	A simple, dry, one-seeded (one-celled) fruit
Acicular	Needle-shaped.
Acidic	Having a low pH, opposite of basic or alkaline.
Acroscopic	Pointing towards, or on the side of, the apex
Acuminate	Gradually tapered to a point. Sharply pointed.
Acute	Pointed or sharp, tapering to a point with straight sides.
Adnate	Fusion of unlike parts, e.g. stamens fused to petals.
Adventive	A plant that grows in the wild in New Zealand but which was introduced to the country by humans.
Agglutinated	Stuck together.
Allelopath	An organism that releases compounds that are toxic to other species.
Allelopathy	The release by an organism of compounds that are toxic to other species.
Alternate	Attached singly at each node but changing from one side of a stem to the other.
Alveolate	Honeycombed with ridged partitions.
Amplexicaul	clasping or surrounding the stem
Anamorph	Asexual fruiting stage, usually of an ascomycete fungus.
Anastomosing	Rejoining after branching, as in some leaf veins.
Annual	A plant that completes its complete life cycle within the space of a year
Annual evergreen	Plants that lose their over-wintering leaves rapidly in the first half of the growing season. Annual evergreens never present a leafless appearance, but are closer in a functional sense to a deciduous plant than they are to multi-annual evergreens.
Annulus	Line of thickened cells that governs the release of spores from a sporangium
Anterior	Towards the front.
Anther	The pollen-bearing portion of the stamen.
Antheridium	Male reproductive organ formed on the prothallus of a fern
Anthesis	When the flower is fully developed and functioning. The time of pollination or bloom.
Apex	Tip; the point furthest from the point of attachment.
Apices	Plural of apex. Tip, the point furthest from the point of attachment
Apiculate	Bearing a short slender and flexible point.
Apiculus	A small, slender point.
Apomixis	A form of reproduction whereby seed is formed without the usual mode of sexual fusion
Appressed	Pressed against another organ or surface.
Aquatic	Growing, or living in, or frequenting water. Applied to plants and animals and their habitats. Opposite of terrestrial (land living).
Archegonium	Female reproductive organ of a fern formed on the prothallus
Arcuate	Curved into an arch.
Aril	An often fleshy appendage on the outside of a seed.
Artificial thinning	Selectively removing vegetation to create gaps to facilitate natural invasion of native plants, or to plant later successional plants.
Ascending	Growing obliquely upward.
Asexual	Vegetative reproduction, lacking sexual involvement by sperm or egg cells
Attenuate	Narrowing gradually
Auricle	A small, ear-shaped appendage.
Auriculate	Bearing a small, ear-shaped appendage.
Autogamous	Self-fertilising flowers.
Autotrophic	Of or relating to organisms (as green plants) that can make complex organic nutritive compounds from simple inorganic sources by photosynthesis
awn	A stiff or bristle like projection often from the tip or back of an organ
Axil	The upper angle between the leaf and the stem.
Axis	The longitudinal supporting structure around which organs are borne, e.g., a stem bearing leaves.
Barbellate	Barbed, having or covered with protective barbs or quills or spines or thorns or setae
Basal	At the base.
Basiscopic	Pointing towards the base
Beak	A prominent extension of an organ
Bifid	Deeply split into two lobes.
Bifurcate	Divided into two.

Term	Definition
Biosecurity	Preventing, eradicating, controlling and managing risks posed by pests and diseases.
Biotic	Pertaining to the living parts of the environment
Bipinnate	With each primary pinna divided to the midrib into a secondary pinna
Biserrate	Doubly serrate.
Blade	The flattened part of a leaf.
Blunt	Not pointed at the ends
Bog	A quagmire covered with specialised plants including sphagnum moss, grasses, sedges, rushes, sundews, umbrella ferns and other plants; has wet, spongy ground, a marsh-plant community on wet, very acid peat. Fed only by rainfall.
Bottleneck	A genetic term; refers to the fact that in smaller populations there could be lower genetic variability
Brachyblasts	Short shoots
Bract	A reduced leaf or leaf-like structure at the base of a flower.
Bracteate	Bearing bracts: leaves or leaf-like structure reduced at the base of a flower.
Bracteolate	With small bracts.
Bracteole	A small bract.
Bracteoles	Bracts directly below the flower
Brevideciduous	Brief (1 month or less) loss of most leaves from the canopy just before flowering or during flushing of a new cohort of leaves.
Bryophyte	Plant group including mosses, liverworts and hornworts
Bryophytes	Plant group including mosses, liverworts and hornworts
Bulbil	A bud produced vegetatively on the stem or frond that is capable of breaking off and growing into a new plant
Bullate	With rounded projections covering the surface as if blistered
Caespitose	Growing in dense tufts
Calli	Circular, warty, stalked thickenings commonly found on the lip (labellum) of the orchid (plural of callus).
Callose	Hardened or thickened.
Callus	Stalked thickening on the lip (labellum) of an orchid.
Calyx	The group of sepals, or outer floral leaves, of a flower
Campanulate	Bell-shaped.
Canaliculate	With longitudinal channels or grooves.
Canopy	The uppermost cover formed by the branches and leaves of trees or the spread of bushes, shrubs and ground covers.
Canopy closure	Stage where canopies of shrub and tree species meet.
Canopy manipulation	Selectively removing vegetation to create gaps to facilitate natural invasion of native plants, or to plant later successional plants.
Capillary	Hair-like
Capitula	Plural of capitulum: A dense head-like inflorescence of many flowers as occurs in most Asteraceae (daisies)
Capitulum	A dense head-like inflorescence of many flowers as occurs in most Asteraceae (daisies)
Capsule	A dry fruit formed from two or more fused carpels that splits open when ripe.
Carbon sinks	Carbon locked away, or sequestered e.g. by trees
Carpel	One unit of the female part of a flower that consists of a basal seed-bearing ovary joined to a receptive stigma by a stalk-like style.
Cauda	Tail-like appendage. (pl. caudae; adj. caudate)
Caudex	The axis of a woody plant, esp. a palm or tree fern, comprising the stem and root.
Cauline	Belonging to the stem, as in cauline leaves emerging from the stem.
Cerise	Bright or deep red.
Chartaceous	Having a papery texture.
Chlorophyll	The green pigment of plants.
Chlorotic	Lacking chlorophyll, therefore yellowish, suffering from chlorosis.
Cilia	Short small hair-like structures on a cell or microorganism
Ciliate	With small hairs (cilia).
Ciliolate	Diminutive of ciliate, i.e., having very small hairs
Cladode	Flattened stem with the function of a leaf
Cladodes	Usually flattened, photosynthetically active branches, these may be leaf-like (e.g., <i>Phyllocladus</i>) or branch-like (e.g., <i>Carmichaelia</i>)
Clavate	Club-shaped, gradually widening towards apex.
Cleft	Having indentations that extend about halfway to the center, as in certain leaves.
Cleistogamous	Flowers that self-fertilise without opening.
Coherent	Sticking together of like parts.
Column	Stamen and stigmas fused to form a single organ.

Term	Definition
Columnar	Shaped like a column
Composite	many small flowers tightly packed together e.g., daisy flowers.
Compound	Composed of several similar parts (cf simple)
Concave	Curved inward.
Conecolorous	Of the same colour.
Conical	Cone-shaped.
Connate	Fusion of like parts.
Conspecific	Individuals of the same species.
Cordate	Heart-shaped with the notch at the base.
Coriaceous	Leather-like; thick, tough, and somewhat rigid.
Corolla	The whorl of petals of a flower.
Corymb	Modified raceme where stalks of lower flowers are elongated to same level as the upper flowers.
Cosmopolitan	A species or other taxonomic group that is distributed widely throughout the world.
Costa	The midrib
Crenate	With rounded teeth (bluntly toothed) along the margin.
Crisped	Margin tightly wavy or crinkled, curled or wavy.
Cristate	With a crest.
Crown	The growing point of an upright rhizome or trunk. This usually produces a tuft or ring of fronds.
Crura	The two small projections at the mouth of a utricle in Carex
Cucullate	Hood-shaped.
Culm	The erect stem of a grass.
Cuneate	Wedge-shaped.
Cupular	Cup-shaped.
Cuttings	Stems and/or leaves taken from plants for propagation
Cyathium	A cup-like structure that surrounds the inflorescence in Euphorbia
Cyme	Inflorescence at the terminus of a branch and where new flowering branches emerge laterally below the flower.
Cytotrace	Populations (or infraspecific taxa) that differ in chromosome number or chromosome morphology, e.g., Nematoceras trilobum agg. has two cytotypes, a diploid and a tetraploid (in which the chromosomes are doubled).
Cytotype	Populations (or infraspecific taxa) that differ in chromosome number or chromosome morphology, e.g., Nematoceras trilobum agg. has two cytotypes, a diploid and a tetraploid (in which the chromosomes are doubled).
Deciduous	Marked leaflessness in winter, and greater than 90% leaves lost by beginning of spring flush.
Decrescent	Diminishing.
Decumbent	With a prostrate or curved base and an erect or ascending tip.
Decurrent	Attached by a broadened base.
Decurved	Curved downward.
Deflexed	Bent abruptly downward.
Dehiscence	The time of opening at maturity to release the contents, e.g., a capsule releasing the seeds.
Dehiscent	Splitting open at maturity to release contents (of a fruit).
Deltoid	Shaped broadly like an equilateral triangle.
Dentate	Toothed along the margin with the teeth pointing outward, not forward.
Denticles	minute teeth
Denticulate	having a very finely toothed margin
Dichotomous	Divided into two equal branches.
Digitiform	Finger-like.
Dioecious	Having male and female flowers on separate plants of the same species.
Diploid	With two complete sets of chromosomes in each cell.
Disarticulating	Separating at a joint.
Discoid	Disc-shaped.
Disjunct	A species or other taxonomic group that occupies areas that are widely separated and scattered and therefore have a discontinuous distribution.
Distal	Toward the apex, away from the point of attachment (cf. proximal).
Distichous	In two rows on opposite sides of the axis.
Divaricating	Branching at a very wide angle with stiff intertwined stems.
Domatia	small structures on the lower surface of a leaf in some woody dicotyledons, located in the axils of the primary veins and usually consisting of depressions partly enclosed by leaf tissue or hairs.

Term	Definition
Dorsal	Of the back or outer surface relative to the axis. (cf. ventral)
Drupe	A stone fruit, the seed enclosed in a bony covering (endocarp) which is surrounded by a + fleshy layer (mesocarp)
Early successional species	Plants which are able to colonise an open area after disturbance but which are often temporary and are replaced by taller plants in time and shaded out.
Echinate	having sharply pointed spines or bristles.
Ecological district	A characteristic landscape and biological community defined in the PNA (Protected Natural Area) programme.
Ecological restoration	Attempt to reinstate original (pre-disturbance) state of a habitat, plant community or ecosystem.
Ecosourced	Plants sourced from seed collected from similar naturally growing plants in the area of the planting site.
Ecosourcing	Using native plants grown from locally grown seeds. Eco-sourced plants help to preserve the ecological distinctiveness of an area, and ecosourced plants fare better and are adapted to survive in the local conditions.
E glandular	Without glands.
Elaiosome	Fleshy, oil-rich structure attached to seed that attracts ants which act as dispersers.
Ellipsoid	Elliptic in long section and circular in cross-section.
Elliptic	Broadest at the middle
Emarginate	With a notch at the apex.
Emarginated	Having a shallow notch at the tip, as in some petals and leaves.
Emergent	In an aquatic sense - wetland herbs that are rooted in the substrate below water level, but carry leaves and stems above the water level e.g. rushes and raupo. Found on the shallow margins of lakes, ponds and waterways. In a forest sense - tree that is appearing above the surrounding canopy.
Emergent marginals	An aquatic plant having most of its structure above water. Other aquatic plants are submerged or floating.
Endemic	Unique or confined to a place or region, found naturally nowhere else.
Endophyte	An endosymbiont (usually a bacterium or fungus) that lives within a plant for at least part of its life without causing any apparent disease.
Endophytes	Endosymbionts (usually bacteria or fungi) that live within plants for at least part of their lives without causing any apparent disease.
Endosperm	The nutritive tissue of a seed, consisting of carbohydrates, proteins, and lipids.
Enrichment planting	Returning to a revegetation site and creating gaps, or filling existing gaps, with different plants of plants, usually later successional plants which may not have survived being planted in the first phases of the project.
Ensiform	Sword shaped
Entire	Smooth. Without teeth, notches or divisions.
Entomophilous	Pollinated by insects.
Epicalyx	Calyx-like structure outside, but close to, the true calyx.
Epigeal	Growing on or close to the ground or emerging from the ground after germination (often used for cotyledons).
Epiphyte	A plant that grows upon another plant but is not parasitic and does not draw nourishment from it.
Epiphytic	Growing upon another plant but not parasitic and not drawing nourishment from it
Erose	Irregularly toothed, as if gnawed.
Estuarine	Pertaining to the meeting of freshwater and seawater wetlands.
Ethnobotany	The study of people's classification, management and use of plants.
Eusporangia	Sporangia that arise from groups of epidermal cells
Evanescent	Lasting a very short time or running a short distance.
Ex situ	Away from the place of natural occurrence.
Ex-situ	Maintenance of plants as live specimens or propagules in cultivation as insurance against the loss of wild populations and as source for material for translocation.
Excurrent	Having the axis prolonged to form an undivided main stem or trunk (as in conifers).
Extravaginal	Outside an enclosing sheath
Falcate	Hooked or curved like a sickle.
Fastigiate	Branches erect and close to central axis.
Fen	A type of wet land that accumulates peat deposits. Fens are less acidic than bogs, deriving most of their water from groundwater rich in calcium and magnesium.
Ferruginous	Rust-like (a colour term)
Fertile frond	Fronds that bear sporangia.
Filamentous	Resembling a filament.
Filiform	Thread like, resembling a filament.
Filiramulate	Branching at a very wide angle with stiff intertwined stems.
Fimbriae	Plural of fimbria: Fringe. A fimbria is composed of many fimbillae (individual hair-like structures).
fimbriate	With fringes.
Flabellate	Fan shaped.
Flaccid	Limp, not rigid, flabby.
Flange	A projecting rim.

Term	Definition
Flexuose	With curves or bends.
Floccose	Having tufts of soft woolly hairs
Floret	A small flower, usually one of a cluster - the head of a daisy for example.
Foliaceous	Leaf-like.
Foliolate	Having leaflets.
Founder effect	When a small number of plants (and therefore their genes) from a larger population are selected some genetic information is lost.
Frond	A leaf, the complete leaf of a fern including the stipe and lamina
Fulvous	Orange-yellow.
Funneliform	Funnel-shaped.
Fusiform	Broadest near the middle and tapering toward both ends.
Galea	Helmet- or hood-shaped.
Galeate	Shaped like a helmet or hood.
Gametophyte	A plant that produces sperm and egg cells and in which sexual reproduction takes place - in ferns this is known as the prothallus
Gene pool	The mixture of all genes and gene variations of a group or population.
Genetic diversity	The variety of genes in a plants or populations.
Genetic variation	Differences displayed by individuals within a plant which may be favoured or eliminated by selection.
geniculate	abrubtly bent
Genus	A taxonomic rank of closely related forms that is further subdivided in to species (plural = genera). In a scientific name (e.g., <i>Sicyos australis</i>), the first word is the genus, the second the species.
Gibbous	Swollen or enlarged on one side, as in a gibbous moon.
Glabrescent	Lacking hair or a similar growth or tending to become hairless
Glabrous	Without or devoid of hairs, smooth.
Gland	A structure that secretes a sticky or oily substance.
Glandular	A structure that secretes a sticky or oily substance.
Glaucoous	Covered with a fine, waxy, removable powder that imparts a white or bluish cast to the surface.
Gley	A soil prone to seasonal inundation.
Globose	Globe-shaped.
Glume	One of two bracts at the base of a grass spikelet.
Groundwater	Groundwater is the water beneath the surface that can be collected with wells, tunnels, or drainage galleries, or that flows naturally to the earth's surface via seeps or springs. Groundwater is the water that is pumped by wells and flows out through springs.
Gymnosperm	Plants in the class Gymnospermae that have seeds which are not enclosed in an ovary.
Gynodioecious	A species population containing plants that produce bisexual (perfect) flowers, and plants that produce only female (pistillate) flowers.
Gynoecium	The female reproductive organs of a flower; the pistil or pistils considered as a group. Means literally "womans house" i.e., the overall structure that contains the female sex organs
Hastate	Spear like. Shaped like an arrowhead, but with basal lobes pointing outward rather than downward.
Haustorium	The absorbing organ of a parasite or hemiparasite
Hemi-parasite	Obtains water and nutrients from the roots of other plants but also manufactures food through photosynthesis.
Hemi-parasitic	Obtaining water and nutrients from the roots of other plants then manufacturing food through photosynthesis.
Herbarium	The place where collections of dried/pressed plants are kept.
Hermaphrodite	Having both male and female sexual characteristics and organs.
Heteroblastic	Exhibiting differences in leaf shapes or forms in juvenile and adult phases of the plant.
Heteroblasty	The state of being heteroblastic (i.e., exhibiting differences in leaf shapes or forms in juvenile and adult phases of the plant).
Hirsute	Hairy.
Hyaline	Membranous, thin and translucent.
Hybrid	An individual that is the offspring of a cross between two different varieties or species.
Hybridise	Breeding with a member of a different plant or type.
Hydrophyte	A plant species adapted to growing in or on water or in wet situations. Aquatic or semi-aquatic.
Hymenium	The fertile, spore-bearing layer of a fruitbody.
Hypanthium	A ring-like, cup-shaped, or tubular structure of a flower on which the sepals, petals, and stamens are borne.
Imbricate	Overlapping.
imbricating	Overlapping.
Imparipinnate	Odd-pinnate, a leaf shape; pinnate with a single leaflet at the apex.
In-situ	On site conservation relating to the maintenance of plants in the wild.
Inbreeding	Genetic similarity in offspring of closely related individuals.

Term	Definition
Incoherent	Not sticking together.
Incursion	Entrance of a pest into an area where it is not present
Indumentum	A covering of fine hairs (or sometimes scales)
Indusia	Plural of indusium, a membrane covering a sorus of a fern
Indusium	A thin tissue that covers the sorus in many ferns. Plural: indusia.
Inflorescence	The arrangement of flowers on the stem. A flower head.
Infundibuliform	Funnel-like.
Interkeel	The space between the keel and the leaf blade
Internode	The part of an axis between two nodes; the section of the stem between leaves.
Internodes	Part of a stem between two nodes.
Intramarginal	Within or near the margin.
Invulneral bracts	The scales surrounding the flower head or capitula.
Involute	A group of bracts surrounding a flower head.
Involute	With margins rolled inward toward the upper side.
Irritable	Responding to touch.
Jugate	Paired.
Juvenile	A plant of non-reproducing size.
Keel	A prominent or obvious longitudinal ridge (as in a boat).
Labellar	Pertaining to the labellum: a lip; in orchid flowers referring to the middle petal which usually differs in size, shape or ornamentation from the two lateral petals.
Labellum	A lip; in orchid flowers referring to the highly modified middle petal which usually differs in size, shape or ornamentation from the two lateral petals.
Lacinia	A jagged lobe.
Laciniæ	Jagged lobes.
Lacinate	Cut into narrow, irregular lobes or segments.
Lacustrine	Of or having to do with a lake, of, relating to, or formed in lakes, growing or living in lakes.
Lamina	The expanded flattened portion or blade of a leaf, fern frond or petal.
Lanceolate	Lance-shaped; of a leaf several times longer than wide with greatest width about one third from the base, tapering gradually to apex and more rapidly to base
Lateral	On or at the side.
Lax	With parts open and spreading, not compact.
Laxly	With parts open and spreading, not compact
Leaflet	One section of a compound leaf.
Lemma	The lower of two bracts enclosing the flower in grasses.
Lenticillate	Bark that is covered in fine lenticels (breathing pores)
Ligulate	Strap-like, tongue-shaped
Ligule	The membrane between the leaf and the stem of a grass; the "petal" of a ray floret in a composite inflorescence
Linear	Long and narrow with more or less parallel sides.
Littoral	Occurring at the border of land and sea (or lake). On or pertaining to the shore. The shallow sunlit waters near the shore to the depth at which rooted plants stop growing.
Lobe	A recognisable, but not separated, rounded division or segment of a leaf or pinna. Used to describe ferns and leaves in <i>Cotula</i> and <i>Leptinella</i> .
Lobed	Part of a leaf (or other organ), often rounded, formed by incisions to about halfway to the midrib.
Lobule	A small lobe or sub-division of a lobe
Lustrous	Glossy, shiny.
Lycophytes	Seedless vascular plants that belong to the phylum Lycophyta (characterised by microphylls -primitive leaves found in ancient plants).
Lyrate	Pinnatifid or pinnatisect terminal lobe much larger than lower lobes.
Maculate	Blotched or spotted.
Mangrove	Coastal wetland dominated by Manawa or mangrove <i>Avicennia marina</i> var. <i>resiifera</i> . Northern New Zealand only, salt marsh replaces it further south.
Margin	The edge or border of a leaf
Marine	Pertaining to the sea and saltwater systems.
Marsh	A tract of wet land principally inhabited by partially-submerged herbaceous vegetation. Has fewer woody plants than swampier habitats.
Mealy	Dry, powdery, crumbly.
Median	In the middle.
Membranous	Very thin, like a membrane.
Mid-lobe	The middle part into which a leaf is divided.
Midrib	The central or principal vein of a leaf or pinna of a fern.
Mire	Synonymous with any peat-accumulating wetland. Term covers bogs and peaty swamps, fens, carr, moor, muskeg and peatland. Term excludes marsh which is non-peat forming.

Term	Definition
Molecular techniques	Where proteins and genes are used to investigate plant relationships
Monitoring	Recording of quantitative data over time to document changes in condition or state of species or ecosystems.
monoecious	Having male and female flowers on the same plant of the same species.
Montane	Land between 300 and 800 metres above sea level.
Mucronate	Tipped with a short, sharp, point.
Mucronulate	Having a very small mucro; diminutive of mucronate.
Multi-annual evergreen	Overlapping annual cohorts of leaves always present.
Multifid	Cleft into many lobes or segments
Multiseptate	With many septa.
Mycorrhiza	A symbiotic relationship between a fungus and a plant.
Mycorrhizal associations	Symbiotic association between fungi and plant roots which assists plant health by allowing increased ability for uptake of nutrients and promote plant growth.
Napiform	A long swollen but tapering root – like a parsnip, or carrot.
Native naturalised	Naturally occurring in New Zealand (i.e., not introduced accidentally or deliberately by humans). Referring to plants that have escaped from cultivation (including gardens or forest plantations) and can now reproduce in the wild (without human assistance)
Nectary	Organ that produces nectar.
Nerve	Prominent vein or rib.
Nerves	Strands of conducting and usually strengthening tissue in a leaves or similar structures
Net veins	Veins that repeatedly divide and re-unite.
Net venation	Feather-like or hand-like venation on a leaf.
Nival	Growing at high altitudes. From Latin: nivalis, snowy etc. from nix, nivis, snow.
Node	The point at which leaves, branches or roots arise on a stem.
Ob-	Prefix meaning inverted, in reverse direction.
Obcordate	Heart shaped with the notch at the apex.
Oblanceolate	Tapering and widest towards the apex or inversely lanceolate.
Oblique	Slanting; of a leaf, larger on one side of the midrib than the other, in other words asymmetrical.
Oblong	Rectangular.
Obovate	Roughly elliptical or reverse egg shaped and widest near the apex (i.e., the terminal half broader than the basal half).
Obtuse	Blunt or rounded at the apex, with the sides meeting at an angle greater than 90°.
Operculate	With a small lid.
Opposite	A pair of organs attached at nodes in pairs on either side of a stem or axis.
Orbicular	Almost or approximately circular.
Outbreeding depression	A reduction in vigor of offspring from distant parents. It can occur when a locally adapted population is moved and mixed with plants adapted to different conditions.
Outer canopy deciduous	Marked reduction in leaf number in the outer canopy in exposed high light environments over winter.
Oval	Planar, shaped like a flattened circle, symmetrical about both the long and the short axis; about twice as long as broad, tapering equally both to the tip and the base. Synonymous with elliptical.
Ovary	Part of a flower containing the ovules and later the seeds.
Ovate	Egg-shaped and widest at base.
Ovoid	Oval; egg-shaped, with rounded base and apex.
Pakihi	A term which in its strict sense refers to open clears within forest dominated by low scrub and rushes. However, more usually used to refer natural and induced wetlands and their associated shrublands. A vernacular most frequently used in the West Coast for impoverished soils and their associated peats, left after forest has been cleared
Palea	The small upper bract enclosing the flower of a grass
Palmettately	Radiating from a point, as fingers radiating from the palm of a hand.
Palmatifid	Deeply divided into several lobes arising from more or less the same level.
Palmatisect	Intermediate between palmate and palmatifid, i.e. the segments are not fully separated at the base; often more or less digitate.
Palustrine	Pertaining to wet or marshy habitats. Term covers mires and marshes
Pandurate	Fiddle-shaped.
Panicle	Highly branched (multiple raceme).
Papilla	A short rounded projection.
Papillae	A soft, fleshy projection, usually small and nipple-like.
Papillate	With short rounded projections.
Papillose	Warty, with short rounded projections or gland-dotted
Parallel venation	Veins are parallel along leaf.

Term	Definition
Parasite	An organism that derives all its nourishment from its host.
Patent	Spreading or expanded, e.g., spreading petals.
Peat	A mass of partially carbonised plant tissue formed by partial decomposition in water of various plants and especially of mosses of the genus <i>Sphagnum</i> , widely found in many parts of the world, varying in consistency from a turf to a slime used as a fertiliser, as stable litter, as a fuel, and for making charcoal. Partially carbonized vegetable matter saturated with water; can be used as a fuel when dried. A type of soil deriving from dead organic material situated in a wet area, where the reduced amount of oxygen available in the wet conditions results in the organic material not decomposing as much as it usually would do so in the presence of more oxygen. Used in growing media. Represents an important carbon sink –drainage of peat releases large amounts of carbon (CO ₂) to the atmosphere.
Pedicel	The stalk of a single flower in an inflorescence or fruit (either in a cluster or existing singularly).
Peduncle	The stalk of a solitary flower or the main stalk of an inflorescence or flower cluster.
Pedunculate	Describing fruits, which are borne on a stalk (a peduncle).
Pellucid	Transparent.
Peltate	Shield-like, with the stalk attached well inside the margin
Pendent	Hanging down from its support
Pendulous	Hanging or drooping.
Penicillate	With a tuft of hairs at the end, like a brush.
Perennial	A plant lasting for three seasons or more
Perianth	A collective term for the calyx (sepals or tepals) and corolla (petals) of the flower, especially when these are indistinguishable
Petal	Part of flower inside the sepals; usually coloured.
Petiolate	Having a petiole.
Petiole	Leaf stalk.
Phloem	The vascular tissue in land plants that is primarily responsible for the distribution of sugars and nutrients manufactured in a shoot.
Photopoint	A monitoring technique where repeat photos are taken of the same scene from the same point over a period of time in order to quantify changes.
Pilose	Bearing long, soft hairs.
Pinna	A segment of a divided lamina that is classified as primary, secondary or tertiary according to the degree of dissection of the lamina.
Pinnae	Divisions of a pinnate leaf
Pinnate	With leaflets arranged regularly in two rows on either side of a stalk as in a feather; the lamina on a fern is divided into separate pinnae
Pinnatifid	Pinnately lobed, cleft more than halfway to the midrib. Not cleft all the way to the rachis.
Pinnatisect	Pinnately divided almost to midrib but segments still confluent.
Pioneer	Plant species are hardy species that should be planted first to establish a good canopy cover that restricts weed growth and promotes natural regeneration. In natural ecosystems these are the first plants to arrive and grow on a site.
Pistil	The female reproductive organ of a flower, consisting of an ovary, style, and stigma.
Pistillate	A flower with one or more pistils, but no stamens.
Plano-convex	Flat on one side, convex on the other.
Plumose	Feathery.
Podzol	Infertile, acidic soil, strongly leached to form a whitish-grey subsoil underlain by a layer enriched in iron, aluminium and organic matter; usually under forest in a wet temperate climate.
Pole	A subcanopy size individual with a long thin trunk and foliage tuft of a potential canopy tree.
Pollinia	Compact masses of orchid pollen.
Population enhancement	Increasing a population for a specific biological purpose, e.g., when a species is already present in an area but extra individuals are added to address a sex imbalance.
Porrect	Extending forward.
Procumbent	Lying and flat along the ground but not rooting
Propagate	To reproduce a plant by sexual (i.e., from seed) or asexual (e.g., from cuttings) means.
Prostrate	A general term for lying flat along the ground. This includes procumbent (that is lying and flat along the ground but not rooting) and decumbent (with a prostrate or curved base and an erect or ascending tip).
Provenance	The place of origin (of a plant that is in cultivation).
Proximal	Toward the base or point of attachment (cf. distal).
Pseudobulb	Thickened surface stem; usually looking like a bulb.
Pseudoterminal	Falsely terminal – as in a bud which appears to occupy a terminal position but does not
Puberulent	Minutely clad in short, soft hairs
Pubescence	Covering of soft, fine hairs
Pubescent	Covered in short, soft hairs.
Pungent	Ending in a stiff sharp point
Pustule	Small blister-like elevation.

Term	Definition
Quadrata	Square, rectangular.
Raceme	An unbranched, elongated inflorescence with pedicellate flowers maturing from the bottom upward i.e., flowers attached to the main stem by short stalks.
Rachis	the axis of an inflorescence or of a compound leaf
Ray	An outer ring of strap-like florets in the head of Asteraceae (daisy) flowers.
Re-introduction	Translocating wild or cultivated individuals to sites where the taxon has been known to occur in the past, but from which it has disappeared.
Recurved	Curved backward.
Reflexed	Bent back on itself
Reniform	Kidney shaped.
Repand	With a slightly wavy margin.
Replum	The outer structure of a pod in which the valves have dehisced (persists after the opening of the fruit)
Restiad	Area dominated by rush-like plants (collectively known as restiads) of the family Restionaceae. Includes Chatham Island and North Island Sporodanthus and oioi (<i>Apodasmia similis</i>)
Retorse	Pointing backward.
Retuse	A shallow notch at the rounded or blunt apex of a leaf.
Rhizoid	Any of various slender filaments that function as roots in mosses and ferns and fungi.
Rhizomatous	With underground creeping stems.
Rhizome	An underground stem (usually spreading horizontally or creeping) or short and erect.
Rhombic	Diamond-shaped.
Rhomboid	Diamond shaped, nearly rhombic.
Riparian	Relating to or living or located on the bank of a natural watercourse (as a river) or sometimes of a lake or a tidewater.
Riparian margin	Refers to the edges of streams, rivers, lakes or other waterways.
Riparian plants	Refers to plants found growing near the edges of streams, rivers or other waterways.
Riparian zone	A strip of land next to streams, rivers, and lakes where there is a transition from terrestrial (land vegetation) to aquatic (water) vegetation. Also known as "berm".
Riverine	Pertaining to rivers, streams and such like flowing water systems.
Rootstock	A short, erect, underground stem.
Rosette	A radiating cluster of leaves.
Rostellum	In orchids, a modified stigma that prevents self-fertilisation.
Rosulate	A dense radiating cluster of leaves.
Rugose	Wrinkled.
Rugulose	Having small wrinkles.
Runcinate	Sharply pinnatifid or cleft, the segments directed downward.
Runner	A trailing stem that roots at the nodes.
Rupestral	Growing on rocks.
Rushes	A group of distinctive wetland plants. They have solid stems (grasses have hollow stems), true rushes <i>Juncus</i> sp. have rounded leaves.
Sagittate	Shaped like the head of an arrow; narrow and pointed but gradually enlarged at base into two straight lobes directed downwards; may refer only to the base of a leaf with such lobes; cf. hastate.
Salt marsh	A coastal wetland, with specialized salt tolerant plants (halophytes).
Sapling	A juvenile tree that has reached the stage of 1 or 2 main stems but is still in the shrub layer.
Saprophyte	A plant lacking chlorophyll and living on dead organic matter.
Saprophytic	Lacking chlorophyll and living on dead organic matter.
Sarcotesta	The fleshy, often highly coloured outer layer of the seed coat in some species, e.g., titoki (<i>Alectryon excelsus</i>).
Scabrid	Roughened or rough with delicate and irregular projections.
Scale	Any thin, flat, membranous structure.
Scape	A leafless flower stem.
Scutiform	Shield-shaped.
Sedges	A group of grass-like or rush-like herbaceous plants belonging to the family Cyperaceae. Many species are found in wetlands some are forest floor plants. Leaves are usually angular. Hence the saying "rushes are round and sedges have edges".
Seedling	A newly germinated plant.
Self-sustaining	Able to sustain itself, or replace itself, independently of management i.e. regenerate naturally
Self-thinning	Natural tree death in a crowded, even-aged forest or shrubland.
Semi-deciduous	Partial leaflessness in winter, and greater than 50% leaves lost by the beginning of spring flush.
Sepal	Outer part of flower; usually green.
Serrate	Sharply toothed with teeth pointing forwards towards apex.
Serrulate	Finely serrate, i.e., finely toothed with asymmetrical teeth pointing forward; like the cutting edge of a saw.

Term	Definition
Sessile	Attached by the base without a stalk or stem.
Seta	The stalk of a fruiting moss capsule
Sheath	A portion of an organ that surrounds (at least partly) another organ (e.g., the tubular envelope enclosing the stem in grasses and sedges).
Silicles	The flattened usually circular capsule – compared with the narrow, elongated fruit (silique) – containing the seed/seeds. A term used almost exclusively for plants within the cabbage family (Brassicaceae)
Silique	A capsule, usually 2-celled, with 2 valves falling away from a frame (replum) bearing
Simple	Of one part; undivided (cf compound).
Sinuate	With a wavy margin.
Sinus	The space or recess between lobes; in hebes a gap between the margins of two leaves of an opposite pair that may be present in the bud before the pair of leaves separate.
Sorus	A cluster of two or more sporangia on the margin or underside of the lamina of a fern, sometimes protected by an indusium.
Spathulate	Spatula or spoon-shaped, a rounded blade tapering gradually to the base.
Spheroidal	Almost spherical but elliptic in cross section.
Spicate	Arranged in a spike.
Spike	Flowers attached to main stem without stalks.
Spikelet	Collection of individual grass florets borne at the end of the smallest branch of the inflorescence.
Sporangia	Plural of sporangium. Structures in which spores are produced.
Sporangium	Structure in which spores are produced.
Spore	A single-celled reproductive unit similar in function to that of the seed in a flowering plant.
sporophyte	The spore producing plant in ferns that is usually the visible part.
Stamen	The male reproductive organ of a flower where pollen is produced. Consists of an anther and its stalk.
Stamens	The male, pollen bearing organ of a flower.
Standing water	Where water lies above the soil surface for much of the year.
Stellate	Irregularly branched or star shaped.
Stigma	Female part of the flower that is receptive to pollen, usually found at or near the tip (apical end) of the style where deposited pollen enters the pistil.
Stipe	The stalk of a frond.
Stipitate	Borne on a stipe or stalk.
Stipulate	A leaf with stipules.
Stipule	A scale-like or leaf-like appendage at the base of a petiole, usually paired.
Stolon	A stem which creeps along the ground, or even underground.
Stoloniferous	Producing stolons
Stramineous	Chaffy, like straw or straw-colored.
Stria	A fine line or groove.
Striae	Fine lines or grooves.
Striate	Fine longitudinal lines or minute ridges
Style	The elongated part of the flower between the ovary and the stigma.
Sub-	A prefix meaning under, somewhat or almost.
Subglabrous	Very slightly, but persistently, hairy.
Suborbicular	Slightly rounded in outline
Substrate	The surface upon which an orchid grows.
Subtended	Immediately beneath, occupying a position immediately beneath a structure, i.e., flower subtended by bract
Subulate	Slender and tapering to a point.
Succession	Progressive replacement of one species or plant community type by another in an ecosystem.
Successional	Referring to species, plant communities or habitats that tend to be progressively replaced by another.
Succulent	Fleshy and juicy.
Summer-green	Used in New Zealand to indicate herbs or sub-shrubs that die down to a root stock or rhizomatous network.
Supplementary planting	Returning to a revegetation site and creating gaps, or filling existing gaps, with different plants of plants, usually later successional plants which may not have survived being planted in the first phases of the project.
Surface water	Water present above the substrate or soil surface.
Surveillance	Regular survey for pests inside operational and managed areas e.g. nurseries, standout areas on parks.
Survey	Collection of observations on the spatial distribution or presence or absence of species using standardised procedures.
Sustainable Land Management	The use of farming practices which are sustainable both financially and environmentally including management of nutrient runoff, waste disposal or stock effluent, reducing impacts of nutrients on waterways, preventing erosion and soil loss, and protecting native forest and wetland habitats from stock damage.
Swamp	Low land that is seasonally flooded; has more woody plants than a marsh and better drainage than a bog. They are more fertile and less acidic than bogs because inflowing water brings silt, clay and organic matter. Typical swamp plants include raupo, purei and harakeke (flax). Zonation and succession often leads through manuka to kahikatea swamp forest as soil builds up and drainage improves.

Term	Definition
Symbiote	An organism that has an association with organisms of another species whereby the metabolic dependence of the two associates is mutual.
Symbiotic	The relation between two different species of organisms that are interdependent; each gains benefits from the other (see also symbiosis).
Sympatric	Occupying the same geographical region.
Synangia	Structures made up of fused sporangia
Synonym	A botanical name that also applies to the same taxon.
Systematics	The study of taxonomy, phylogenetics, and taxogenetics.
Tabular	Shaped like a rectangular tablet.
Taxa	Taxonomic groups. Used to refer to a group at any level e.g., genus, species or subspecies.
Taxon	A taxonomic group. Used to refer to a group at any level e.g., genus, species or subspecies.
Taxonomy	The process or science of classifying, naming, and describing organisms
Tepal	An individual member of the perianth.
Terete	Cylindrical and tapering.
Terminal	At the tip or apex.
Ternatifid	Leaflets In threes,
Tetrad	A group of four.
Tomentum	A hairy covering of short closely matted hairs.
Translocation	The movement of living organisms from one area to another.
Trifid	Divided into three.
Trifoliate	Having three leaflets.
Trigonus	Three–angled
Tripinnate	With each secondary pinna divided to the midrib into tertiary pinnae
Triquetrous	Triangular in cross section and acutely angled.
Truncate	With the apex or base squared at the end as if cut off.
Tuberculate	Bearing small swellings.
Tubular	Tube-shaped.
turbinate	Top-shaped.
Turgid	Distended through internal pressure
Type locality	The place or source where a holotype or type specimen was found for a species.
Ultramafic	A type of dark, usually igneous, rock that is chemically dominated by magnesium and iron-rich minerals, the partially metamorphosed form of which is serpentinite.
Umbel	Umbrella like; the flower stalks arise from one point at the stem.
Undulate	Wavy edged.
Undulose	Wavy edged.
Unitubular	A tube partitioned once – literally one tube (compare – multitubular – many tubes)
Utricle	A thin loose cover enveloping some fruits (eg., Carex, Uncinia)
Valvate	Opening by valves.
Vascular plant	A plant that possesses specialised conducting tissue (xylem and phloem). This includes flowering plants, conifers and ferns but excludes mosses, algae, lichens and liverworts.
Velutinous	Thickly covered with delicate hairs; velvety.
Ventral	Of the front or inner (adaxial) surface relative to the axis. (cf. dorsal)
Vermiform	Worm-shaped.
Vernicose	Glossy, literally as if varnished, e.g., Hebe vernicosa has leaves that appear as if varnished
Verrucose	Having small rounded warts.
Verticillium	A fungus disease that will cause wilting and death.
Villous	Covered with long, soft, fine hairs.
Water table	The level at which water stays in a soil profile. The zone of saturation at the highest average depth during the wettest season.
Wetland	A site that regularly has areas of open water for part or all of the year, or has a water table within 10 cm of the surface for at least 3 months of the year. Wetland ecosystems support a range of plant and animal species adapted to a aquatic or semi-aquatic environment.
Whipcord	A shrub in which the leaves are reduced to scales that are close-set and pressed against the stem.
Whorl	A ring of branches or leaves arising at the same level around the stem of a plant.