



Flora of NZ - Native plant identification study-guide - Angiosperms 1/2



Table of Contents

Introduction	1
<i>Alectryon excelsus</i> subsp. <i>excelsus</i>	2
<i>Apodasmia similis</i>	3
<i>Aristotelia serrata</i>	4
<i>Beilschmiedia tarairi</i>	5
<i>Brachyglottis repanda</i>	6
<i>Carmichaelia australis</i>	7
<i>Coprosma repens</i>	8
<i>Coprosma robusta</i>	9
<i>Cordyline australis</i>	10
<i>Corokia cotoneaster</i>	11
<i>Corynocarpus laevigatus</i>	12
<i>Dodonaea viscosa</i>	13
<i>Dysoxylum spectabile</i>	14
<i>Entelea arborescens</i>	15
<i>Geniostoma ligustrifolium</i> var. <i>ligustrifolium</i>	16
<i>Griselinia littoralis</i>	17
<i>Hebe stricta</i> var. <i>stricta</i>	18
<i>Hoheria populnea</i>	19
<i>Knightia excelsa</i>	20
<i>Kunzea ericoides</i>	21
Glossary	22

Made on the New Zealand Plant Conservation Network website – www.nzpcn.org.nz

Copyright

All images used in this book remain copyright of the named photographer. Any reproduction, retransmission, republication, or other use of all or part of this book is expressly prohibited, unless prior written permission has been granted by the New Zealand Plant Conservation Network (info@nzpcn.org.nz). All other rights reserved.

Introduction

This book was compiled from information stored on the website of the New Zealand Plant Conservation Network (www.nzpcn.org.nz).

This website was established in 2003 as a repository for information about New Zealand's threatened vascular plants. Since then it has grown into a national database of information about all plants in the New Zealand botanic region including both native and naturalised vascular plants, threatened mosses, liverworts and fungi.

Funding to develop the website was provided by the New Zealand Government's Terrestrial and Freshwater Biodiversity Information System Programme (TFBIS).

The species information used on the website has come from a variety of sources. The indigenous vascular plant text was written largely by Dr Peter de Lange (former Network Vice President). Peter based the descriptions on a wide range of sources including the Flora of NZ Series (Allan 1961, Moore and Edgar 1970 and Webb et al 1987) as well as numerous other taxonomic treatments. For a full bibliography of information sources see the References at the end of this book.

Where no published treatment was available Peter used herbarium specimens and his own knowledge of the flora to prepare species pages. Various other contributors have provided text and additional information to many species pages including botanists such as Mike Thorsen, John Barkla, Cathy Jones, Simon Walls, Nick Singers and many others. The threatened fungi text was written by Eric Mackenzie and Peter Buchanan (Landcare Research).

More than 200 photographers have kindly provided images to illustrate the website and for use in this book especially John Smith-Dodsworth, Jeremy Rolfe, Peter de Lange, Wayne Bennett and Gillian Crowcroft.

The New Zealand Botanic Region

The information on the Network website, from which this book was compiled, is for species that are indigenous to or naturalised within the New Zealand Botanic Region as defined by Allan (1961). The New Zealand botanic region encompasses the Kermadec, Manawatawhi/Three Kings, North, South, Stewart Island/Rakiura, Chatham, Antipodes, Bounties, Snares, Auckland Campbell island/Motu Ihupuku and Macquarie.

About the Network

The Network has more than 800 members worldwide and is New Zealand's largest non-governmental organisation solely devoted to the protection and restoration of New Zealand's indigenous plant life.

The vision of the New Zealand Plant Conservation Network is that '*no indigenous species of plant will become extinct nor be placed at risk of extinction as a result of human action or indifference, and that the rich, diverse and unique plant life of New Zealand will be recognised, cherished and restored*'.

Since it was founded in 2003 the Network has undertaken a range of conservation initiatives in order to achieve its vision.

That work has included:

- Training people in plant conservation
- Publishing plant books, reports and posters
- Raising money for the David Given Threatened Plant Research Trust to pay for plant conservation research scholarships
- Advocacy to raise awareness of the importance of plant life in general and especially New Zealand's status as a Global Centre of Plant Diversity
- Lobbying central and regional government and business to protect indigenous plant life
- Educating people about plant life through the Network website
- Connecting people through the monthly newsletter, the Network conference and the annual general meeting

What is a threatened plant?

The NZ Threatened Plant Committee was formed in 1991 and ever since then it has met at regular intervals to review the status of indigenous vascular plants. It is made up of a small group of botanists that between them have an extensive knowledge of the native plants of New Zealand. This group is chaired by Dr Peter de Lange of the New Zealand Department of Conservation.

This committee applies a set of criteria to each native plant to determine its conservation status. The resulting list of species classified as threatened is published in the NZ Journal of Botany (see for example de Lange et al. 2009). The main threat categories used are: Extinct, Critical, Endangered, Vulnerable, Declining. Other categories used are: Recovering, Relict, Naturally Uncommon, Coloniser, Vagrant and Data Deficient. For vascular plants the threat status used in this book is taken from the 2009 conservation assessment (see de Lange et al 2009).

More recently other committees have been established to review the status of non-vascular plants but their lists are yet to be published.

Alectryon excelsus subsp. *excelsus*

Common Name(s):

New Zealand ash, titoki

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. North and South Islands from Te Pahi to Banks Peninsula

Habitat:

A widespread coastal to lowland forest tree. Often favouring well drained, fertile, alluvial soils along river banks and associated terraces. It is also a major component of coastal forests, particularly those developed within exposed situations or on basaltic or andesite volcanics. It is a common offshore island tree within the Hauraki Gulf. The large fruits are bird dispersed and so titoki trees often occur as a sparse components of most lowland forest types, throughout the North Island.

Features:

Tree between 10m and 20m tall. Branches stout, erect, all parts invested with fine, velutinous, ferruginous hairs. Bark brown. Adult leaves dark green, matt when mature, imparipinnate, alternate 80-260 mm long. Leaflets 3-7 pairs; lamina 45-105 x 19-40 mm, subcoriaceous, lanceolate, oblong or narrowly-ovate, apex, subacute often acuminate, rarely obtuse; base cuneate, truncate to oblique, upper leaf surface matt; lamina margin entire or deeply serrated 1-4 times near apex. Inflorescences axillary 90-120 mm long, sparingly branched panicles. Flowers bisexual or staminate. Petals absent. Stamens 5-8 in bisexual and 6-10 in staminate flowers, crimson. Stigma ovoid, in staminate flowers ovary tholiform, style absent, in perfect flowers broadly urceolate, style 1.5-2 mm, erect. Fruits sessile, 1-2-lobed, 14-20 x 9-14 mm, pubescent, globular, carina 3-5 mm long on one side. Seed 7-10 x 4-8 mm, subglobose, black, lustrous, sarcotesta fleshy, scarlet, papillose.

Flowering:

October - December (-June)

Fruiting:

November - August

Threats:

Not Threatened

References and further reading:

Cameron, E.K. 1998. Frost resistance in titoki *Alectryon*. Auckland Botanical Society Journal, 53: 15.

Duguid, F. 1961. Flowering in titoki. Wellington Botanical Society Bulletin, 32: 16

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 2009 Vol. 11 No. 4 pp. 285-309

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1486



Caption: Algies Bay, Auckland
Photographer: John Sawyer



Caption: Carter Scenic Reserve
Photographer: John Sawyer

Apodasmia similis

Common Name(s):

jointed wire rush, oiioi

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. Three Kings, North, South, Stewart and Chatham Islands.

Habitat:

Mostly coastal in estuaries, saltmarshes, dunes and sandy flats and hollows. Occasionally inland in gumland scrub, along lake margins, fringing peat bogs or surrounding hot springs.

Features*:

Dioecious, rush-like perennial herb. Rhizomes 3-7 mm diameter, covered in closely sheathing, imbricating, dark brown scales, 10-20 mm long, each enclosing a tuft of coarse brown hairs. Culms numerous, 0.5-2.6 x 1.5-2.5(-3.0) mm, densely packed, erect, sometimes with upper third decurved to more or less pendulous, simple, terete, glaucous, grey-green, yellow-green or red-green. Leaves reduced to bract-like sheaths, these dark brown or maroon-black, regularly spaced at 70-90 mm intervals at the base of the culm, 10-60 mm apart higher up; margins entire. Male inflorescences, paniculate or fascicled, bearing numerous stalked spikelets; upper floral bracts ovate-lanceolate, mucronate, red-brown to maroon, margins membranous; tepals 6-4 more or less completely hyaline, the outer longer, brownish, the inner shorter, paler; stamens 3; ovary rudimentary. Female inflorescences fascicled, spikelets more or less sessile; upper floral bracts ovate, mucronate, > tepals; tepals 6, the outer keeled, lanceolate, acuminate, inner flat, smaller, more or less hyaline, more obtuse, mucronate; styles 3, united to midway, bright red to orange-red; staminodes 0. Fruit c.1 x 0.5 mm, triquetrous, indehiscent. Seed c.1 x 0.4 mm, oblong-elliptical, golden-brown, surface reticulate, both ends apiculate, one end dark brown, the other, almost white.

Flowering:

October - December

Fruiting:

December - March

Threats:

Not Threatened

*Attribution:

Description adapted from Edgar and Moore (1970).

References and further reading:

Briggs, B.G. & Johnson, L.A.S. (1998) New genera and species of Australian Restionaceae (Poales). *Telopea* 7: 345-373. http://www.rbgsyd.nsw.gov.au/__data/assets/pdf_file/0004/73237/Tel7Bri345.pdf

Moore, L.B.; Edgar, E. 1970: *Flora of New Zealand*. Vol. I. Government Printer, Wellington.

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 2009 Vol. 11 No. 4 pp. 285-309

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=2052



Caption: *Apodasmia similis*
Photographer: Bec Stanley



Caption: *Apodasmia similis*
Photographer: Bec Stanley

Aristotelia serrata

Common Name(s):

Makomako, wineberry

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. North, South and Stewart Islands. Throughout, but less common in drier areas.

Habitat:

Lowland to montane forests. Often forming dense thickets following disturbance.

Features*:

Dioecious tree to c. 10 m tall; trunk and branches upright, to 30 cm diam.; bark smooth, grey, spotted with lenticels; branchlets light to dark red, pubescent. Leaves opposite to subopposite; petiole slender, to 50 mm long, greenish often flushed pink; midvein conspicuous above, raised below; secondary veins obvious and raised below giving surface a wrinkled uneven appearance; lamina membranous, 5-12 x 4-8 cm, glabrate (pubescence may persist on veins below), broad-ovate, margin deeply doubly and irregularly sharply serrate, tip acuminate, base cordate to truncate, upper surface light or dark green, undersides pale green, frequently infused with purple or pink. Juvenile leaves larger. Inflorescences conspicuous, axillary, flowers 4-6 mm diam., in panicles 6-10 cm long, on slender pubescent pedicels 5-10 mm long. Sepals 4, ovate, c. 3 mm long, pubescent, pink; petals 4, 3-lobed (often deeply), c. 9 mm long, white to light pink to red. Stamens many, on glandular minutely pubescent disc, not exceeding petals. Ovary 3-4-celled, styles 3-4. Fruit a c. 8-seeded fleshy depressed-obovoid berry, 5 x 4 mm, bright red to black. Seed irregularly angled, ventral surface flattened, circular or broadly elliptic, 1.9-3.1 mm, surface irregular, aril absent.

Flowering:

September-December

Fruiting:

November-January

*Attribution:

Description adapted from Allan (1961), Heenan and de Lange (2006), Eagle (2000) and Webb and Simpson (2001).

References and further reading:

Allan, H.H. 1961. Flora of New Zealand. Government Printer, Wellington

Heenan, P.B, de Lange, P.J. 2006. *Pseudowintera insperata* (Winteraceae), an overlooked and rare new species from northern New Zealand. NZ J. Botany 44: 89-98

Eagle, A. 2000. Eagle's complete trees and shrubs of NZ. Te Papa Press, Wellington

Webb, C.J. & Simpson, M.J.A. 2001. Seeds of NZ gymnosperms and dicotyledons. Manuka Press, Christchurch.

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1512



Caption: Flowering wineberry

Photographer: Jane Gosden



Caption: Waikuku, Aorangi

Photographer: John Sawyer

Beilschmiedia tarairi

Common Name(s):

Taraire

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. Confined to the North Island where it most common north of Auckland and Thames. However it also occurs in scattered pockets in the west south of Port Waikato to the Kawhia Harbour, inland at Pukemokemoke (near Tauhei), and in the east it occurs very locally from the eastern end of Papatea Bay to East Cape.

Habitat:

Common canopy forming tree in lowland and lower montane forests north of Auckland. Often associated with kauri (*Agathis australis*), and pohutukawa (*Metrosideros excelsa*), and on basalt rocks and soils puriri (*Vitex lucens*).

Features:

Evergreen tree up to 22 m tall, with very broad canopy crown. Trunk to 1 m diam. Bark smooth, dark brown. Branches stout, spreading. Branchlets, young leaves, petioles and young inflorescences densely clad in reddish brown tomentum. Foliage closely alternate, erectopate, simple, leathery. Petioles (8-)10(-12) mm. Leaves (36-)50-72 (-85) x (26-)34-48(-56) mm, wide-elliptic to wide-obovate, dark green and glabrous above, bullate, glaucous below, with stout veins covered in reddish brown tomentum, margins entire,, apex rounded, retuse and mucronate. Inflorescence and erect, axillary panicle up to 100 mm long. Flowers sexually perfect, 3-5 mm diam., greenish, often partially clothed in dense reddish-brown tomentum, perianth cleft into 6, stamens 12. Fruit an erect, ellipsoid to ovoid drupe (28-)30(-35) x (14-)16(-18) mm, 1-seeded, pericarp fleshy, dark purple when ripe, covered in waxy glaucous bloom.

Flowering:

(September-) November (-December)

Fruiting:

March - November

Threats:

Not Threatened

References and further reading:

Wilcox, M.D. 2001. Establishment of forest monitoring plots in Kirks Bush Papakura with special reference to Taraire (*Beilschmiedia tarairi*). *Auckland Botanical Society Journal*, 56: 76-79.

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 2009 Vol. 11 No. 4 pp. 285-309

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1540



Caption: Wenderholm

Photographer: John Barkla



Caption: Taraire leaves

Photographer: DoC

Brachyglottis repanda

Common Name(s):

Rangiora, bushmans toilet paper

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. North Island throughout. South Island - north west Nelson to just south of Greymouth in the west, and near Kekerengu in the east. Naturalised on Banks Peninsula, Otago Peninsula, and on Stewart Island at Oban.

Habitat:

Common in coastal, lowland and lower montane shrubland and open forest. Often a pioneer species.

Features:

Shrub to small tree up to 6 m or more tall. Trunk one or more arising from ground, covered in somewhat corky bark. Branches stout, spreading, rather brittle, initially densely clad in fine white to buff tomentum becoming glabrescent with age. Petiole stout, grooved, 80-100 mm long. Leaves leathery, 50-250(-300) X 50-20(-30) mm, dark green to pale green above, undersides clad in fine, appressed vivid white hairs, broad- to ovate-oblong, obtuse to subacute, obliquely cordate to truncate at base, margins distantly dentately lobed to sinuate. Inflorescence a much branched panicle. Capitula 5 mm diam., numerous, without ligules (discoïd). Involucral bracts 3 mm long, narrow-oblong to narrow spatulate, margins scarious except at base. Florets 10-12, yellow. Seeds (cypsela) narrowly oblong-elliptic to oblong elliptic, 1-1.8 mm long, ribs 6, rounded, broad. Pappus 2-3 mm, buff-yellow, scabrid.

Flowering:

(July-) August-October
(-November)

Fruiting:

(October-) November-
December (-January)

Threats:

Not Threatened

References and further reading:

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 2009 Vol. 11 No. 4 pp. 285-309

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1562



Caption: *Brachyglottis repanda*
Photographer: Wayne Bennett



Caption: *Brachyglottis repanda*
Photographer: Wayne Bennett

Carmichaelia australis

Common Name(s):

Common Broom

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. New Zealand: North and South Islands (except southern South Island)

Habitat:

Coastal to montane, on river terraces, stream banks, colluvium, rock outcrops, talus and fan toe slopes, among tussock grassland and grey scrub, on the edge and margins of dense bush, forest, and in swamps

Features*:

Shrub, 2-8 × 2-5 m. Branches up to 100 mm diameter, ascending and spreading. Cladodes 30.0-200.0 × 1.5-8.0 mm, ascending or spreading, linear, striate, weakly plano-convex to strongly flattened and compressed, green, yellow-green, or brown-green, glabrous to sparsely hairy, apex obtuse to subacute; leaf nodes 4-15. Leaves 1-3-foliolate, present on seedlings and rarely on adults, terminal leaflet larger; lamina 3.0-22.0 × 1.3-16.0 mm, obovate to oblong, fleshy, green, sometimes with dark mottling, surfaces glabrous to moderately hairy, apex emarginate, base cuneate; petiole 1-10 mm long, glabrous to moderately hairy, green; petiolule < 0.25 mm long, glabrous or sparsely hairy, light green. Leaves on cladodes reduced to scales, < 0.5 mm long, broad-triangular, glabrous, apex subacute, margin hairy. Stipules c.1.0 × c.1.5 mm, free, broad-triangular, upper surface glabrous, lower surface glabrous or glabrescent, apex subacute, margin hairy. Inflorescence a raceme, 1-3 per node, each with 4-15 flowers. Peduncle 1-10 mm long, glabrous to moderately hairy, green, occasionally flushed red. Pedicel 2-4 mm long, glabrous to sparsely hairy, pale green. Calyx 1.3-1.7 × 1.7-2.2 mm, campanulate, green, outer surface glabrous to moderately hairy. Calyx lobes 0.2-0.4 mm long, triangular, inner surface glabrous, appressed to corolla or rarely weakly spreading, apex acute. Standard 4.0-6.0 × 5.0-6.5 mm, obovate, patent, positioned in central part of keel, keeled, apex retuse; inner surface white, sometimes with a purple blotch, purple-veined; outer surface white, green at base, purple-veined; claw c.1 mm long, pale green. Wings 4.0-5.0 × 1.7-2.2 mm, oblong, longer than keel, apex obtuse; inner and outer surfaces white, purple-veined; auricle rounded, white; claw c. 1.5 mm long, pale green. Keel 3.2-4.2 × c.1.5 mm, apex obtuse; distal part of inner and outer surfaces purple, proximal part white; auricle < 0.5 mm long, rounded, pale green; claw c.1.5 mm long, pale green. Stamens 3-4 mm long. Pistil c.4 mm long, exserted, glabrous. Pods 6.5-15 × 2.0-5.5 mm, oblong, broad-oblong, elliptic, broad-elliptic, or sometimes orbicular, laterally compressed, erect or spreading, brown, grey, or black, valves dehiscent; beak 0.5-2.5 mm long, in a central apical position, stout, pungent-tipped. Seeds 1-5 per pod, 2.2-4.0 × 1.7-2.8 mm, oblong to reniform, orange, red, green, or yellow-green and often with black mottling, usually persistent on replum.

Flowering:

October - February

Fruiting:

November - May

Threats:

Not Threatened

*Attribution:

Description from Heenan (1996)

References and further reading:

Heenan, P.B. 1996: A taxonomic revision of *Carmichaelia* (Fabaceae - Galegeae) in New Zealand (part II). *New Zealand Journal of Botany* 34: 157-177. For full synonymy see this paper.

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 2009 Vol. 11 No. 4 pp. 285-309

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1596



Caption: *Carmichaelia australis*
Photographer: Wayne Bennett



Caption: *Carmichaelia australis*
Photographer: Wayne Bennett

Coprosma repens

Common Name(s):

taupata, looking glass plant, mirror plant

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. Three Kings, North and South Islands as far south as Greymouth in the west and Rarangi in the east but now extensively naturalised throughout the South Island, Stewart and Chatham Islands. Also naturalised on Norfolk Island and in Hawaii, in Australia, California and South Africa.

Habitat:

Coastal (rarely inland: Kaitaia – Awanui River, Huntly Basin and in the Manawatu – especially the upper Rangitikei River). A common species of rock stacks, islets, islands coastal cliffs, talus slopes and boulder field. Also a common component of petrel scrub on northern offshore islands, and in coastal forest where it often forms the main understorey and rarely is co-dominant in the canopy. Frequently associated with other coastal *Coprosma*, especially *C. crassifolia*, *C. macrocarpa* subsp. *macrocarpa* and subsp. *minor*, *C. rhamnoides*, *C. neglecta*, and members of the *C. acerosa* complex. Hybrids between *C. repens* and *C. acerosa* are common and are known as *C. xkirkii*, less frequently hybrids between it and *C. crassifolia* are found (*C. xbuchananii*) and with both *C. rhamnoides* and *C. neglecta*.

Features*:

Dioecious (rarely monoecious) shrub or small tree up to 8 m tall, prostrate and widely spreading in exposed sites, shrubb to arborescent in more sheltered situations; branches firm and more or less pliant when young becoming more brittle with age, bark dark to light brown, underbark green; branchlets initially pubescent with short patent hairs, becoming glabrous with age. Leaves on fleshy glabrous, slender to stout petioles 8-16 mm long. Stipule shortly sheathing, margin finely pubescent, otherwise outer surface pubescent, inner more or less glabrous, broad-deltoid, subacute to subtruncate; denticles up to 4 either side of a single large, dark black apical denticle, conspicuous, central one prominent. Lamina thick, subfleshy, coriaceous, 5-90 × 4-60 mm, dark glossy green above, paler and dull below; broad-oblong, elliptic-oblong, broadly ovate-oblong to suborbicular, rounded to truncate, usually apiculate (slightly emarginate to retuse on Three Kings and northern Hauraki Gulf Islands), apiculus caducous, cuneately narrowed to base; margins plane to slightly recurved (very occasionally inrolled). Vein reticulations evident above and especially below. Flowers in compound clusters on branched peduncles. Male flowers 3-20 per cluster; calyx-teeth minute; corolla funnelform, lobes 4-5, acute, about = tube. Female flowers usually 3 per cluster; calyx-teeth short, obtuse; corolla subfunnelform, c.5 mm long, lobes acute or obtuse, < tube; stigmas stout (Perfect flowers occasional (though with pollen often aborted or malformed) through out range but especially common on the northern offshore islands). Drupe orange-red, red (rarely yellow), obovoid often slightly compressed, 8-12 × 8-10 mm

Flowering:

June - February

Fruiting:

July - June

Threats:

Not Threatened

*Attribution:

Description based on Allan (1961) though supplemented with additional measurements and observations taken from herbarium specimens and wild plants.

References and further reading:

Allan, H.H. 1961: Flora of New Zealand. Vol. I, Government Printer, Wellington.

Dawson, J.W. 1961. *Coprosma*. The Spike (or Victoria University College Review). Victoria University of Wellington Student's Association.

Gordon, H.D. 1959. Sex ratio in *Coprosma repens* (rubiaceae). Wellington Botanical Society Bulletin, 31: 11

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 11: 285-309

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1730



Caption: Awhitu Peninsula, Auckland region

Photographer: John Sawyer



Caption: Coprosma repens

Photographer: Wayne Bennett

Coprosma robusta

Common Name(s):

karamu, glossy karamu

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. North and South Islands. Naturalised on the Chatham Islands within a small area between Waitangi and Owenga.

Habitat:

Common throughout coastal, lowland and lower montane habitats within shrublands and open sites within forest.

Features:

Shrub or small tree up to 6 m tall. Branches numerous, stout, erect to somewhat spreading. Petioles stout, 10-20 mm long. Stipules fused towards base, obtuse, glabrous with one of two prominent, black, glandular denticles. Leaves 70-120 x 30-40-50 mm, leathery, dark green above, paler green beneath, glabrous, elliptic, elliptic-oblong to broad-ovate, acute or obtuse, apex mucronate. Venation reticulated, conspicuous. Male flowers in axillary many-flowered glomerules, corolla conspicuous, lobes triangular, acute, stamens 4-5, prominent. Females in compound clusters on peduncles 10-15 mm. Calyx and corolla much reduced, stigmas prominent. Drupe dark orange (rarely yellow), 8-8 x 4-5 mm, oblong to narrow-ovoid.

Flowering:

(July-) August-September (-November)

Fruiting:

(March-) April-May (-July)

Threats:

Not Threatened

References and further reading:

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285-309

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1733



Caption: Fruit of *Coprosma robusta*

Photographer: Wayne Bennett



Caption: *Coprosma robusta* (Karamu)

Photographer: Wayne Bennett

Cordyline australis

Common Name(s):

cabbage tree, ti, ti kouka, palm lily

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. Common in the North, South and Stewart Islands. Probably naturalised on the Chatham Islands.

Habitat:

Widespread and common from coastal to montane forest. Most commonly encountered on alluvial terraces within riparian forest.

Features:

Tree up to 20 m tall, trunk stout, 1.5-2 m diam, many-branched above (prior to flowering, trunk slender and solitary, branching happens after the first flowering). Bark corky, persistent, fissured, pale to dark grey. Leaves numerous (0.2-)0.3-1(-1.5) x (0.2)-0.3(-0.6) m, dark to light green, narrowly lanceolate to lanceolate, erect to erecto-patent, scarcely inclined to droop, midrib indistinct. Petiole indistinct, short. Inflorescence a panicle. Peduncle stout, fleshy 40 mm or more in diam., panicle of numerous flowers, (0.6-)1(-1.8) x .3-0.6(-0.8) m, branching to third or fourth order, these well spaced, basal bracts green and leaf-like, ultimate racemes 100-200 mm long, 20 mm diam., bearing well-spaced to somewhat crowded, almost sessile to sessile flowers and axes. Flowers sweetly perfumed, perianth 5-6 mm diam., white, tepals free almost to base, reflexed. Stamens about same length as tepals. Stigma short, trifid.

Flowering:

(September-) October-
December (-January)

Fruiting:

(December-)
January-March

Threats:

Populations have been decimated from some parts of the country due to a mysterious illness linked to a Myoplast Like Organism (MLO) which is believed to cause the syndrome known as Sudden Decline. Plants stricken with this illness suddenly, and rapidly, wilt, with the leaves failing off still green. If the bark is peeled off the base of the tree near the soil line blackened or rotten spots are typically present. Once stricken with Sudden Decline there is no cure and the trees can die within days. Recently there has been some evidence to suggest the severity of Sudden Decline is lessening.

References and further reading:

Beever, R. et al. 1996. Sudden decline of cabbage tree. *NZ Journal of Ecology*, 20(1): 53-68

Duguid, F. 1976. *Cordyline australis* at Lake Kopureherehe. *Wellington Botanical Society Bulletin*, 39: 46-47

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285-309

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1744



Caption: Awhitu Regional Park, Auckland region

Photographer: John Sawyer



Caption: *Cordyline australis*

Photographer: Wayne Bennett

Corokia cotoneaster

Common Name(s):

Korokio, wire-netting bush

Current Threat Status (2012):

Not Threatened

Threats:

Not Threatened

References and further reading:

Moorfield, J. C. (2005). Te aka : Maori-English, English-Maori dictionary and index. Pearson Longman: Auckland, N.Z.

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 11: 285-309

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1756



Caption: Upper Oreti River, Southland

Photographer: Jesse Bythell



Caption: Fuit, Greenstone Valley

Photographer: John Barkla

Corynocarpus laevigatus

Common Name(s):

Karaka, kopi

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. Exact indigenous distribution uncertain due to its widespread historic planting by Maori. Common from Raoul and the Three Kings Islands, throughout the North and South Islands to Banks Peninsula and Okarito. Also on the Chatham Islands. Most botanists accept it as native only to the northern half of the North Island. It is probably naturalised from deliberate Polynesian plantings on Raoul and the Chatham Islands.

Habitat:

Common in mainly coastal situations, often a major component of coastal forest, rarely dominant. Occasionally found inland, and then often in association with Maori cultural deposits.

Features*:

Leafy canopy tree up to 15 m tall. Trunk stout up to 1 m diam., Bark grey. Branches stout, erect to spreading. Petioles 10-15 mm long. Leaves dark green above paler beneath, thick, leathery, (50-)100-150 (-200) x (30-)50-70 mm, glossy, elliptic to obovate-oblong, margins recurved. Inflorescence a stout, erect panicle up to 200 mm long, peduncles and pedicels short, somewhat fleshy, pale green. Flowers 4-5 mm diam., greenish-cream to off-white or pale yellow. Sepals suborbicular, petals 5, obovate-spathulate, alternating with 5 subpetaloid staminodes. Fruit an ellipsoid to ovoid drupe 25-40(-46) mm long, flesh pale yellow to orange. Endocarp a fibrous reticulum surrounding a smoother, harder papery layer beneath. This structure enclosing a single seed (kernel).

Flowering:

August - November

Fruiting:

January - April

Threats:

Abundant and not threatened. Often naturalising in suitable habitats.

*Attribution:

Fact Sheet prepared for NZPCN by P.J. de Lange (1 September 2004). Description based on Allan (1961).

References and further reading:

Allan, H.H. 1961: Flora of New Zealand. Vol. 1. Wellington, Government Printer.

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 11: 285-309

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1762



Caption: Seedlings. Lake Westmere, Whanganui. Feb 2013.
Photographer: Colin Ogle



Caption: Seedlings. Lake Westmere, Whanganui. Feb 2013.
Photographer: Colin Ogle

Dodonaea viscosa

Common Name(s):

akeake

Current Threat Status (2012):

Not Threatened

Threats:

Not Threatened

References and further reading:

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285-309

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=2109



Caption: Coromandel, November
Photographer: John Smith-Dodsworth



Caption: Coromandel, November
Photographer: John Smith-Dodsworth

Dysoxylum spectabile

Common Name(s):

kohekohe, New Zealand mahogany

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. North and South Islands. In the South Island not extending much beyond the Marlborough Sounds, reaching a southern limit near the Hurunui River (Napenape).

Habitat:

Common and sometimes dominant or co-dominant tree of coastal to lowland forest.

Features:

Tree up to 15 m tall usually with abroad, spreading canopy. Trunk up to 1 m diam., branches stout, erect then spreading. Bark pale brown, under bark green. Leaves compound, imparipinnate, alternate on pulvinate petioles up to 40 mm long, leaflet pairs 4-6, (50-)-150(-200) x (20-)30(-80) mm, opposite to subopposite, bright green, yellow-green to dark green, ovate to obovate-oblong, leathery, margins somewhat undulate. Plants gynodioecious, with fixed female and inconstant males on different trees. Inflorescence a cymose, drooping, panicle arising from trunk and branches (cauliflorous). Flowers c. 30 mm diam., fleshy. Pedicels short. Calyx divided to base, lobes broad-oblong, abruptly pointed, ciliate, petals linear, 10 mm, spreading, waxy white or greenish. Capsules, woody, broad-obovoid to subglobose, 3-4-celled, c. 25 mm long, green. Seeds 2 per cell, orange or scarlet.

Flowering:

March - June

Fruiting:

April - August

Threats:

Not Threatened. However, where possum and rat numbers are high this species is not actively regenerating. Possums defoliate trees, and will heavily browse inflorescences such that few succeed in flowering and setting fruit. Rats are major seed predators. Only where control of these animals is undertaken, or on possum and rodent-free offshore islands can one see kohekohe flowering, fruiting and regenerating freely. If numbers of these introduced animals remain unchecked, it is clear that kohekohe will decline and vanish from large parts of its natural range.

References and further reading:

Duguid, F. 1985. Kohekohe *Dysoxylum spectabile* as an accidental epiphyte. Wellington Botanical Society Bulletin, 42: 11

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1825



Caption: Colonial Knob Scenic Reserve, Porirua.

Photographer: Jeremy Rolfe



Caption: Colonial Knob Scenic Reserve, Porirua.

Photographer: Jeremy Rolfe

Entelea arborescens

Common Name(s):

Whau

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. Three Kings, North (including Little and Great Barrier Islands) and South Islands. In the North Island, whau is locally common from Te Pahi to about Kawhia and Mahia Peninsula south of there it is known from a few sites in the eastern Wairarapa, at Paekakariki and Wellington. In the South Island it is confined to the Golden Bay area of North-West Nelson. Whau naturalises easily and has established south of these stations from bach and urban plantings.

Habitat:

Coastal to lowland forest or shrubland. Usually in open sites such as around recent slips, tree falls, cliff faces, boulder slopes, sand dunes or on the margins of streams, rivers, lagoons and lakes. Mostly near the coast however it may occur well inland in some places e.g., the Waikato River near Hamilton, Rotorua. Some inland and southern North island occurrences are thought to be derived from deliberate plantings by Maori.

Features*:

Shrub or small spreading tree up to c. 8 m tall; trunk up to 0.25 m dbh; wood-weight very light; bark firm, grey, tearing in long fibrous strips when cut. branches numerous, upright than spreading; branchlets, leaves, petioles, inflorescences densely clad in soft whitish branched hairs; leaf-scars oval or lunate. Leaves alternate, softly membranous (wilting readily if picked), green, bright green to yellow green, ± glossy, venation distinct when fresh or dry; petioles 80-300 mm long, stout; stipules linear-acuminate, ± persistent. Lamina 50-100-150(-300) × 50-100-150-(260) mm, obliquely very broad-ovate, abruptly acuminate, cordate at base, margins doubly crenate-serrate, sometimes obscurely lobed, 3-5-7-subpalmately lobed. Inflorescence a subumbellate many-flowered cyme. Flowers (3-)4-5-merous. Peduncle 100-300 mm long, stout, pedicels 10-40 mm long. Sepals free, 8-10-12 mm long, narrowly lanceolate to triangular, acuminate; petals (3-)4-5, 10-30 × 10-30 mm, orbicular to suborbicular, white, crumpled. Stamens numerous, mostly free sometimes connate at base, filaments 10-18(-20) mm long, white, anthers versatile, yellow. Ovary 5-10 mm long, broadly to narrowly globose or ovoid, hispid, 5-7-locular, ovules numerous, style simple, stigma ± globular to broadly capitate, fringed or toothed. Fruit a bristly capsule 20-35 mm diameter, subglobose to globose, black to charcoal when ripe, invested by numerous, rigid, spinose, black to charcoal coloured hairs 15-25 mm long. Seeds numerous, 1.9-2.9 mm long, obovate, elliptic to broadly elliptic, glabrous, surface granular, orange-yellow, pale brown, or orange-brown. Description of seeds by Webb & Simpson (2001).

Flowering:

August - November

Fruiting:

December - June

Threats:

Not Threatened. However, recent field work gathering samples for a Marsden study into the possible past use of whau by Maori indicates that whau is much less common in the North Island than it once was. browsing pressure from cattle, goats and horses, clearance of coastal scrub of housing and the spread of invasive woody shrubs and trees into many northern coastal areas may be threatening some populations.

*Attribution:

Fact Sheet Prepared for NZPCN by: P.J. de Lange 10 February 2011. Description of seeds by Webb & Simpson (2001).

References and further reading:

Webb, C.J.; Simpson, M.J.A. 2001: Seeds of New Zealand Gymnosperms and Dicotyledons. Christchurch, Manuka Press.

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1377



Caption: Fruits, Auckland
Photographer: John Barkla



Caption: Awhitu Regional Park, Auckland
Photographer: John Sawyer

Geniostoma ligustrifolium var. *ligustrifolium*

Common Name(s):
hangehange

Current Threat Status (2012):
Not Threatened

Threats:
Not Threatened

References and further reading:

Conn, B.J. 1980: A taxonomic revision of *Geniostoma* subg. *Geniostoma* (Loganiaceae). *Blumea* 26: 245-364.

Connor, H.E.; Edgar, E. 1987: Name changes in the indigenous New Zealand flora, 1960–1986 and Nomina Nova IV, 1983–1986. *New Zealand Journal of Botany* 25: 115-170.

Murray, B.G.; de Lange, P. J. 1999: Contributions to a chromosome atlas of the New Zealand flora - 35. Miscellaneous families. *New Zealand Journal of Botany* 37: 511-521

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1923



Caption: Puketi Forest, Northland
Photographer: Dean Baigent-Mercer



Caption: Puketi Forest, Northland
Photographer: Dean Baigent-Mercer

Griselinia littoralis

Common Name(s):

nroadleaf, kapuka, papauma

Current Threat Status (2012):


Not Threatened

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1957



Caption: Mt Frith, Rimutaka

Photographer: John Sawyer



Caption: Coromandel, October

Photographer: John Smith-Dodsworth

Hebe stricta var. *stricta*

Common Name(s):

koromiko

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic to the North Island. Somewhat local in the far North, otherwise common and widespread to about the Manawatu Gorge.

Habitat:

Common in successional habitats from coastal areas to lower montane habitats.

Features*:

Shrub or small tree (1-)2(-4) m tall. Branchlets finely pubescent. Stem internodes longer than stem diameter. Leaf bud without sinus. Leaves, spreading, 50-100(-120) mm, dull green to yellow-green (not glossy), lanceolate, linear-lanceolate, somewhat leathery, apex often acuminate, leaf margin usually entire, occasionally toothed. Inflorescence lateral, racemose, much longer than leaves, drooping, sometimes spiraled, all parts except flower finely pubescent. Flowers sweetly (sometimes over powerfully so) scented, lilac, mauve or white. Corolla tube 6 mm, exceeding calyx, narrow, cylindric, lobes rounded. Capsules < 5 mm long, pendent, all parts pubescent.

Flowering:

(July-) August (-October) but flowering can also occur sporadically throughout the year

Fruiting:

(September-) November (-January) but seed capsules may be found throughout the year

Threats:

Not Threatened

*Attribution:

Fact Sheet Prepared by P.J. de Lange (1 February 2005). Description based on Allan (1961) - see also Bayly & Kellow (2006)

References and further reading:

Allan, H.H. 1961: Flora of New Zealand. Vol. I, Wellington, Government Printer

Bayly M. and Kellow A. (2006). An Illustrated Guide to New Zealand Hebes. Te Papa Press: Wellington

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=2009



Caption: Rotorua, February
Photographer: John Smith-Dodsworth



Caption: Rotorua, February
Photographer: John Smith-Dodsworth

Hoheria populnea

Common Name(s):

Lacebark, houhere, ribbonwood

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. North Island only from North Cape (Pararaki Stream) south to the northern Waikato and Coromandel. However widely planted and often found naturalising throughout the southern North Island, South, Stewart and Chatham Islands.

Habitat:

Coastal to montane usually in Kauri (*Agathis australis*) forest but also in successional forest associated with kauri. Also common in pohutukawa (*Metrosideros excelsa*) dominated coastal forest.

Features*:

Small upright to often spreading tree up to 8 m tall (rarely more); bark of mature trunk and branches dark grey-brown that of younger growth dark red-brown or maroon, branches and branchlets ascending, in some forms pendulous, pliant, slender, often deeply grooved, ± glabrescent, indumentum comprised of short stellate hairs, on mature parts sparse, on young parts and inflorescence sparse to dense. Juvenile foliage identical or nearly so to adult. Leaves subcoriaceous to coriaceous glossy, adaxially dark-green to yellow-green, sometimes with veins more darkly pigmented, abaxially often maroon or purple with darker coloured veins, sometimes green or glaucescent; petioles slender, pliant up to 10 mm long; lamina (5-)10(-30) mm long, broad-ovate to deltoid to suborbicular in outline, margins serrate, usually deeply so, coarsely lobed. Adult leaves similar, on petioles up to 20 mm long; lamina (50-)7(-180) mm × (30-)40(-60) mm, broad-ovate to ovate-lanceolate to elliptic, apex acuminate or acute, obtuse or rounded, base rounded to truncate (rarely subcordate); margins deeply, coarsely, sometimes doubly, serrate-dentate, teeth usually well spaced. Flowers 25-30 mm diameter, both solitary and in (2-)5-10-flowered cymose clusters on same plant; pedicels 8-10(-12) mm long; calyx campanulate, 5-6 mm long, teeth broadly to narrowly triangular; petals 10-12 mm long, white, obliquely oblong, often notched.; stigmas capitate. Mature carpels 5(-6), compressed. Mericarp winged, main body 4.5-6.5 mm long, brown; wing 3.0-8.5 mm long, slightly curved outwards, orange yellow, finely and sparsely covered with stellate hairs. Description adapted from Allan (1961) and Webb & Simpson (2001).

Flowering:

January - March

Fruiting:

April - June

Threats:

Not Threatened

*Attribution:

Fact Sheet Prepared for NZPCN by P.J. de Lange 9 April 2011. Description adapted from Allan (1961) and Webb & Simpson (2001).

References and further reading:

Allan, H.H. 1961: Flora of New Zealand. Vol. I, Wellington, Government Printer.

Webb, C.J.; Simpson, M.J.A. 2001: Seeds of New Zealand Gymnosperms and Dicotyledons. Christchurch, Manuka Press.

Moorfield, J. C. (2005). Te aka : Maori-English, English-Maori dictionary and index. Pearson Longman: Auckland, N.Z.

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=824



Caption: Waitakere Reservoir
Photographer: Gillian Crowcroft



Caption: Hoheria populnea
Photographer: Peter de Lange

Knightia excelsa

Common Name(s):

Rewarewa, NZ honeysuckle

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic monotypic genus. North and South Islands. Common in the North Island, but confined to the Marlborough Sounds in the South Island.

Habitat:

A common tree of coastal, lowland and lower montane shrubland, secondary regrowth, and on occasion mature forest. Frost-tender when young so generally scarce from cooler, frost-prone habitats - nevertheless it can be very common in suitable sites on the Central Volcanic Plateau of the North Island.

Features:

Tall tree with columnar (fastigate) growth-form up to 30 m tall. Trunk up to 1 m diam. Bark dark brown. Branches erect, fastigate, at first angled, clad in red-brown (rust-coloured), velutinous, tomentum. Juvenile leaves yellow-green, 150-300(-400) x 10-15 mm, narrowly linear-lanceolate, sometimes forked 2,3 or 4 times, margins acutely serrated. Adult leaves dark green, 100-150(-200) x 25-40 mm, broad lanceolate to narrow-oblong or oblong, sometimes obovate, occasionally forked, rigid, bluntly and coarsely serrated, covered in deciduous velutinous red-brown pubescence. Inflorescence a stout raceme up to 100(-180) mm x 60 mm, densely flowered. Pedicels and perianth clad in red-brown, velutinous tomentum. Flowers sexually perfect. Perianth 4, exterior covered in red-brown tomentum, interior dark crimson, segments at first cylindric and fused, soon separating and curling spirally. Stamens 4, filaments crimson, short, anthers long, linear, rich golden-yellow. Ovar sessile. Style long, crimson, long persistent. Fruits, follicles 30-40 mm long, 2-valved, woody, pubescent; valves tapering to persistent style. Seeds 10 mm, apex terminated by 15 mm long wing.

Flowering:

(September-)
October-December

Fruiting:

October-January (fruit takes a year to mature, so fruit and flowers may co-occur)

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1383



Caption: Rangitoto Island

Photographer: John Barkla



Caption: Manurewa

Photographer: Gillian Crowcroft

Kunzea ericoides

Common Name(s):

Manuoa, Titira, Atitira, Kanuka

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. New Zealand: Northern South Island only - north of the Buller and Wairau Rivers. Most common in North West Nelson.

Habitat:

Coastal to lowland shrubland, regenerating forest and forest margins, also present in montane forest, ultramafic shrubland and very occasionally present in subalpine shrubland.

Features*:

Trees up to 18 m. Trunk 1–4, 0.10–0.85 m d.b.h. Early bark brown to grey-brown, ± elongate, usually firmly attached, margins elongate sinuous, ± entire with scarcely any flaking; old bark similar. Branches slender, initially ascending soon spreading, apices often pendulous. Branchlets numerous, slender, glabrescent; indumentum sparse, deciduous, hairs divergent 0.02–0.05 mm long; leaves of branchlets densely crowded along stems. Leaves sessile, ± glabrous, except for the margins; lamina 4.0–25.0 × 0.5–1.8 mm, green to yellow-green, linear, linear-lanceolate, to narrowly lanceolate, straight or with upper ¼ weakly recurved, apex acute, sometimes cuspidate, base attenuate; lamina margins initially finely sericeous, glabrate or glabrous; hairs forming a fine, discontinuous band failing just short of lamina apex. Inflorescence a compact corymbiform to shortly elongate 3–15-flowered botryum up 60 mm long. Pherophylls foliose ± persistent, 1 per flower; lamina 3.0–7.8 × 0.9–1.4 mm, elliptic, lanceolate to narrowly lanceolate, apex acute, base attenuate; Pedicels 1.6–3.8 mm long at anthesis, usually glabrous. Flower buds pyriform to narrowly obconic, apex of mature buds weakly domed to flat, calyx lobes distant. Flowers 4.1–8.3 mm diam. Hypanthium 1.4–3.2 × 1.9–4.1 mm; sharply obconic, apex terminating in 5 persistent suberect to spreading calyx lobes; hypanthium glabrous (very rarely with basal ¼ finely, sparsely covered in minute hairs). Calyx lobes 5, suberect to spreading, 0.4–1.0 × 0.4–1.0 mm, orbicular, obtuse to broadly deltoid, red-green, pink or crimson, margins glabrous or finely ciliate. Receptacle green or pink at anthesis, darkening to crimson or dark magenta after fertilisation. Petals 5, 1.4–2.6 × 1.5–2.0 mm, white, orbicular, suborbicular to narrowly ovate, spreading, apex rounded, entire or very finely denticulate, oil glands usually not evident when fresh, ± colourless. Stamens 10–34 in 1–2 weakly defined whorls, filaments white. Anthers dorsifixed, 0.35–0.48 × 0.16–0.24 mm, broadly ellipsoid. Pollen white. Anther connective gland prominent, pink or pinkish-orange when fresh, drying red to orange, ± spheroidal ± coarsely papillate. Ovary 4–5 locular, each with 16–24 ovules in two rows on each placental lobe. Style 1.5–2.2 mm long at anthesis; stigma capitate, about 1¼ × the style diam., flat, cream or white, flushing pink after anthesis, surface very finely granular-papillate. Fruits rarely persistent, 1.9–3.4 × 1.8–3.9 mm, glabrous, dark green to reddish-green, maturing brown to grey-brown to grey-black, cupular, barrel-shaped, shortly cylindrical to hemispherical, calyx valves erect with the apices incurved, split concealed by dried, erect, free portion of hypanthium. Seeds 1.00–1.05 × 0.32–0.50 mm, semi-glossy, orange-brown to dark brown, obovoid, oblong, oblong-ellipsoid, or cylindrical and ± curved, surface coarsely reticulate.

Flowering:

October-February

Fruiting:

November-March

Threats:

Not threatened, though some stands are at risk from clearance for farmland or through felling for firewood.

*Attribution:

Fact Sheet prepared for NZPCN by P.J. de Lange 25 August 2014. Description modified from de Lange (2014).

References and further reading:

de Lange, P.J. 2014: *A revision of the New Zealand Kunzea ericoides (Myrtaceae) complex*. *Phytokeys* 40: 185p doi: 10.3897/phytokeys.40.7973.


For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=885



Caption: *Kunzea ericoides* - tree showing weeping branches characteristic of this species

Photographer: Peter de Lange



Caption: Marahau

Photographer: Peter de Lange

Definitions of botanical terms

A glossary has been provided below with definitions for many of the botanical terms used in the species descriptions.

Glossary

Term	Definition
Abaxial	Facing away from the stem of a plant (especially denoting the lower surface of a leaf).
Acerose	Narrow with a sharp stiff point.
Achene	A simple, dry, one-seeded (one-celled) fruit
Acicular	Needle-shaped.
Acidic	Having a low pH, opposite of basic or alkaline.
Acroscopic	Pointing towards, or on the side of, the apex
Acuminate	Gradually tapered to a point. Sharply pointed.
Acute	Pointed or sharp, tapering to a point with straight sides.
Adnate	Fusion of unlike parts, e.g. stamens fused to petals.
Adventive	A plant that grows in the wild in New Zealand but which was introduced to the country by humans.
Agglutinated	Stuck together.
Allelopath	An organism that releases compounds that are toxic to other species.
Allelopathy	The release by an organism of compounds that are toxic to other species.
Alternate	Attached singly at each node but changing from one side of a stem to the other.
Alveolate	Honeycombed with ridged partitions.
Amplexicaul	clasping or surrounding the stem
Anamorph	Asexual fruiting stage, usually of an ascomycete fungus.
Anastomosing	Rejoining after branching, as in some leaf veins.
Annual	A plant that completes its complete life cycle within the space of a year
Annual evergreen	Plants that lose their over-wintering leaves rapidly in the first half of the growing season. Annual evergreens never present a leafless appearance, but are closer in a functional sense to a deciduous plant than they are to multi-annual evergreens.
Annulus	Line of thickened cells that governs the release of spores from a sporangium
Anterior	Towards the front.
Anther	The pollen-bearing portion of the stamen.
Antheridium	Male reproductive organ formed on the prothallus of a fern
Anthesis	When the flower is fully developed and functioning. The time of pollination or bloom.
Apex	Tip; the point furthest from the point of attachment.
Apices	Plural of apex. Tip, the point furthest from the point of attachment
Apiculate	Bearing a short slender and flexible point.
Apiculus	A small, slender point.
Apomixis	A form of reproduction whereby seed is formed without the usual mode of sexual fusion
Appressed	Pressed against another organ or surface.
Aquatic	Growing, or living in, or frequenting water. Applied to plants and animals and their habitats. Opposite of terrestrial (land living).
Archegonium	Female reproductive organ of a fern formed on the prothallus
Arcuate	Curved into an arch.
Aril	An often fleshy appendage on the outside of a seed.
Artificial thinning	Selectively removing vegetation to create gaps to facilitate natural invasion of native plants, or to plant later successional plants.
Ascending	Growing obliquely upward.
Asexual	Vegetative reproduction, lacking sexual involvement by sperm or egg cells
Attenuate	Narrowing gradually
Auricle	A small, ear-shaped appendage.
Auriculate	Bearing a small, ear-shaped appendage.
Autogamous	Self-fertilising flowers.
Autotrophic	Of or relating to organisms (as green plants) that can make complex organic nutritive compounds from simple inorganic sources by photosynthesis
awn	A stiff or bristle like projection often from the tip or back of an organ
Axil	The upper angle between the leaf and the stem.
Axis	The longitudinal supporting structure around which organs are borne, e.g., a stem bearing leaves.
Barbellate	Barbed, having or covered with protective barbs or quills or spines or thorns or setae
Basal	At the base.
Basisopic	Pointing towards the base
Beak	A prominent extension of an organ
Bifid	Deeply split into two lobes.
Bifurcate	Divided into two.

Term	Definition
Biosecurity	Preventing, eradicating, controlling and managing risks posed by pests and diseases.
Biotic	Pertaining to the living parts of the environment
Bipinnate	With each primary pinna divided to the midrib into a secondary pinna
Biserrate	Doubly serrate.
Blade	The flattened part of a leaf.
Blunt	Not pointed at the ends
Bog	A quagmire covered with specialised plants including sphagnum moss, grasses, sedges, rushes, sundews, umbrella ferns and other plants; has wet, spongy ground, a marsh-plant community on wet, very acid peat. Fed only by rainfall.
Bottleneck	A genetic term; refers to the fact that in smaller populations there could be lower genetic variability
Brachyblasts	Short shoots
Bract	A reduced leaf or leaf-like structure at the base of a flower.
Bracteate	Bearing bracts: leaves or leaf-like structure reduced at the base of a flower.
Bracteolate	With small bracts.
Bracteole	A small bract.
Bracteoles	Bracts directly below the flower
Brevideciduous	Brief (1 month or less) loss of most leaves from the canopy just before flowering or during flushing of a new cohort of leaves.
Bryophyte	Plant group including mosses, liverworts and hornworts
Bryophytes	Plant group including mosses, liverworts and hornworts
Bulbil	A bud produced vegetatively on the stem or frond that is capable of breaking off and growing into a new plant
Bullate	With rounded projections covering the surface as if blistered
Caespitose	Growing in dense tufts
Calli	Circular, warty, stalked thickenings commonly found on the lip (labellum) of the orchid (plural of callus).
Callose	Hardened or thickened.
Callus	Stalked thickening on the lip (labellum) of an orchid.
Calyx	The group of sepals, or outer floral leaves, of a flower
Campanulate	Bell-shaped.
Canaliculate	With longitudinal channels or grooves.
Canopy	The uppermost cover formed by the branches and leaves of trees or the spread of bushes, shrubs and ground covers.
Canopy closure	Stage where canopies of shrub and tree species meet.
Canopy manipulation	Selectively removing vegetation to create gaps to facilitate natural invasion of native plants, or to plant later successional plants.
Capillary	Hair-like
Capitula	Plural of capitulum: A dense head-like inflorescence of many flowers as occurs in most Asteraceae (daisies)
Capitulum	A dense head-like inflorescence of many flowers as occurs in most Asteraceae (daisies)
Capsule	A dry fruit formed from two or more fused carpels that splits open when ripe.
Carbon sinks	Carbon locked away, or sequestered e.g. by trees
Carpel	One unit of the female part of a flower that consists of a basal seed-bearing ovary joined to a receptive stigma by a stalk-like style.
Cauda	Tail-like appendage. (pl. caudae; adj. caudate)
Caudex	The axis of a woody plant, esp. a palm or tree fern, comprising the stem and root.
Cauline	Belonging to the stem, as in cauline leaves emerging from the stem.
Cerise	Bright or deep red.
Chartaceous	Having a papery texture.
Chlorophyll	The green pigment of plants.
Chlorotic	Lacking chlorophyll, therefore yellowish, suffering from chlorosis.
Cilia	Short small hair-like structures on a cell or microorganism
Ciliate	With small hairs (cilia).
Ciliolate	Diminutive of ciliate, i.e., having very small hairs
Cladode	Flattened stem with the function of a leaf
Cladodes	Usually flattened, photosynthetically active branches, these may be leaf-like (e.g., Phyllocladus) or branch-like (e.g., Carmichaelia)
Clavate	Club-shaped, gradually widening towards apex.
Cleft	Having indentations that extend about halfway to the center, as in certain leaves.
Cleistogamous	Flowers that self-fertilise without opening.
Coherent	Sticking together of like parts.
Column	Stamen and stigmas fused to form a single organ.

Term	Definition
Columnar	Shaped like a column
Composite	many small flowers tightly packed together e.g., daisy flowers.
Compound	Composed of several similar parts (cf simple)
Concave	Curved inward.
Concolorous	Of the same colour.
Conical	Cone-shaped.
Connate	Fusion of like parts.
Conspecific	Individuals of the same species.
Cordate	Heart-shaped with the notch at the base.
Coriaceous	Leather-like; thick, tough, and somewhat rigid.
Corolla	The whorl of petals of a flower.
Corymb	Modified raceme where stalks of lower flowers are elongated to same level as the upper flowers.
Cosmopolitan	A species or other taxonomic group that is distributed widely throughout the world.
Costa	The midrib
Crenate	With rounded teeth (bluntly toothed) along the margin.
Crisped	Margin tightly wavy or crinkled, curled or wavy.
Cristate	With a crest.
Crown	The growing point of an upright rhizome or trunk. This usually produces a tuft or ring of fronds.
Crura	The two small projections at the mouth of a utricle in Carex
Cucullate	Hood-shaped.
Culm	The erect stem of a grass.
Cuneate	Wedge-shaped.
Cupular	Cup-shaped.
Cuttings	Stems and/or leaves taken from plants for propagation
Cyathium	A cup-like structure that surrounds the inflorescence in Euphorbia
Cyme	Inflorescence at the terminus of a branch and where new flowering branches emerge laterally below the flower.
Cytorace	Populations (or infraspecific taxa) that differ in chromosome number or chromosome morphology, e.g., <i>Nematoceras trilobum</i> agg. has two cytoraces, a diploid and a tetraploid (in which the chromosomes are doubled).
Cytotype	Populations (or infraspecific taxa) that differ in chromosome number or chromosome morphology, e.g., <i>Nematoceras trilobum</i> agg. has two cytotypes, a diploid and a tetraploid (in which the chromosomes are doubled).
Deciduous	Marked leaflessness in winter, and greater than 90% leaves lost by beginning of spring flush.
Decrescent	Diminishing.
Decumbent	With a prostrate or curved base and an erect or ascending tip.
Decurrent	Attached by a broadened base.
Decurved	Curved downward.
Deflexed	Bent abruptly downward.
Dehiscence	The time of opening at maturity to release the contents, e.g., a capsule releasing the seeds.
Dehiscent	Splitting open at maturity to release contents (of a fruit).
Deltoid	Shaped broadly like an equilateral triangle.
Dentate	Toothed along the margin with the teeth pointing outward, not forward.
Denticles	minute teeth
Denticulate	having a very finely toothed margin
Dichotomous	Divided into two equal branches.
Digitiform	Finger-like.
Dioecious	Having male and female flowers on separate plants of the same species.
Diploid	With two complete sets of chromosomes in each cell.
Disarticulating	Separating at a joint.
Discoid	Disc-shaped.
Disjunct	A species or other taxonomic group that occupies areas that are widely separated and scattered and therefore have a discontinuous distribution.
Distal	Toward the apex, away from the point of attachment (cf. proximal).
Distichous	In two rows on opposite sides of the axis.
Divaricating	Branching at a very wide angle with stiff intertwined stems.
Domatia	small structures on the lower surface of a leaf in some woody dicotyledons, located in the axils of the primary veins and usually consisting of depressions partly enclosed by leaf tissue or hairs.

Term	Definition
Dorsal	Of the back or outer surface relative to the axis. (cf. ventral)
Drupe	A stone fruit, the seed enclosed in a bony covering (endocarp) which is surrounded by a + fleshy layer (mesocarp)
Early successional species	Plants which are able to colonise an open area after disturbance but which are often temporary and are replaced by taller plants in time and shaded out.
Echinate	having sharply pointed spines or bristles.
Ecological district	A characteristic landscape and biological community defined in the PNA (Protected Natural Area) programme.
Ecological restoration	Attempt to reinstate original (pre-disturbance) state of a habitat, plant community or ecosystem.
Ecosourced	Plants sourced from seed collected from similar naturally growing plants in the area of the planting site.
Ecosourcing	Using native plants grown from locally grown seeds. Eco-sourced plants help to preserve the ecological distinctiveness of an area, and ecosourced plants fare better and are adapted to survive in the local conditions.
Eglandular	Without glands.
Elaiosome	Fleshy, oil-rich structure attached to seed that attracts ants which act as dispersers.
Ellipsoid	Elliptic in long section and circular in cross-section.
Elliptic	Broadest at the middle
Emarginate	With a notch at the apex.
Emarginated	Having a shallow notch at the tip, as in some petals and leaves.
Emergent	In an aquatic sense - wetland herbs that are rooted in the substrate below water level, but carry leaves and stems above the water level e.g. rushes and raupo. Found on the shallow margins of lakes, ponds and waterways. In a forest sense - tree that is appearing above the surrounding canopy.
Emergent marginals	An aquatic plant having most of its structure above water. Other aquatic plants are submerged or floating.
Endemic	Unique or confined to a place or region, found naturally nowhere else.
Endophyte	An endosymbiont (usually a bacterium or fungus) that lives within a plant for at least part of its life without causing any apparent disease.
Endophytes	Endosymbionts (usually bacteria or fungi) that live within plants for at least part of their lives without causing any apparent disease.
Endosperm	The nutritive tissue of a seed, consisting of carbohydrates, proteins, and lipids.
Enrichment planting	Returning to a revegetation site and creating gaps, or filling existing gaps, with different plants of plants, usually later successional plants which may not have survived being planted in the first phases of the project.
Ensiform	Sword shaped
Entire	Smooth. Without teeth, notches or divisions.
Entomophilous	Pollinated by insects.
Epicalyx	Calyx-like structure outside, but close to, the true calyx.
Epigeal	Growing on or close to the ground or emerging from the ground after germination (often used for cotyledons).
Epiphyte	A plant that grows upon another plant but is not parasitic and does not draw nourishment from it.
Epiphytic	Growing upon another plant but not parasitic and not drawing nourishment it
Erose	Irregularly toothed, as if gnawed.
Estuarine	Pertaining to the meeting of freshwater and seawater wetlands.
Ethnobotany	The study of people's classification, management and use of plants.
Eusporangia	Sporangia that arise from groups of epidermal cells
Evanescent	Lasting a very short time or running a short distance.
Ex situ	Away from the place of natural occurrence.
Ex-situ	Maintenance of plants as live specimens or propagules in cultivation as insurance against the loss of wild populations and as source for material for translocation.
Excurrent	Having the axis prolonged to form an undivided main stem or trunk (as in conifers).
Extravaginal	Outside an enclosing sheath
Falcate	Hooked or curved like a sickle.
Fastigate	Branches erect and close to central axis.
Fen	A type of wet land that accumulates peat deposits. Fens are less acidic than bogs, deriving most of their water from groundwater rich in calcium and magnesium.
Ferruginous	Rust-like (a colour term)
Fertile frond	Fronds that bear sporangia.
Filamentous	Resembling a filament.
Filiform	Thread like, resembling a filament.
Filiramulate	Branching at a very wide angle with stiff intertwined stems.
Fimbriae	Plural of fimbria: Fringe. A fimbria is composed of many fimbriae (individual hair-like structures).
fimbriate	With fringes.
Flabellate	Fan shaped.
Flaccid	Limp, not rigid, flabby.
Flange	A projecting rim.

Term	Definition
Flexuose	With curves or bends.
Floccose	Having tufts of soft woolly hairs
Floret	A small flower, usually one of a cluster - the head of a daisy for example.
Foliaceous	Leaf-like.
Foliate	Having leaflets.
Founder effect	When a small number of plants (and therefore their genes) from a larger population are selected some genetic information is lost.
Fronid	A leaf, the complete leaf of a fern including the stipe and lamina
Fulvous	Orange–yellow.
Funneliform	Funnel-shaped.
Fusiform	Broadest near the middle and tapering toward both ends.
Galea	Helmet- or hood-shaped.
Galeate	Shaped like a helmet or hood.
Gametophyte	A plant that produces sperm and egg cells and in which sexual reproduction takes place - in ferns this is known as the prothallus
Gene pool	The mixture of all genes and gene variations of a group or population.
Genetic diversity	The variety of genes in a plants or populations.
Genetic variation	Differences displayed by individuals within a plant which may be favoured or eliminated by selection.
geniculate	abruptly bent
Genus	A taxonomic rank of closely related forms that is further subdivided in to species (plural = genera). In a scientific name (e.g., <i>Sicyos australis</i>), the first word is the genus, the second the species.
Gibbous	Swollen or enlarged on one side, as in a gibbous moon.
Glabrescent	Lacking hair or a similar growth or tending to become hairless
Glabrous	Without or devoid of hairs, smooth.
Gland	A structure that secretes a sticky or oily substance.
Glandular	A structure that secretes a sticky or oily substance.
Glaucous	Covered with a fine, waxy, removable powder that imparts a white or bluish cast to the surface.
Gley	A soil prone to seasonal inundation.
Globose	Globe-shaped.
Glume	One of two bracts at the base of a grass spikelet.
Groundwater	Groundwater is the water beneath the surface that can be collected with wells, tunnels, or drainage galleries, or that flows naturally to the earth's surface via seeps or springs. Groundwater is the water that is pumped by wells and flows out through springs.
Gymnosperm	Plants in the class Gymnospermae that have seeds which are not enclosed in an ovary.
Gynodioecious	A species population containing plants that produce bisexual (perfect) flowers, and plants that produce only female (pistillate) flowers.
Gynoeceium	The female reproductive organs of a flower; the pistil or pistils considered as a group. Means literally "womans house" i.e., the overall structure that contains the female sex organs
Hastate	Spear like. Shaped like an arrowhead, but with basal lobes pointing outward rather than downward.
Haustorium	The absorbing organ of a parasite or hemiparasite
Hemi–parasite	Obtains water and nutrients from the roots of other plants but also manufactures food through photosynthesis.
Hemi–parasitic	Obtaining water and nutrients from the roots of other plants then manufacturing food through photosynthesis.
Herbarium	The place where collections of dried/pressed plants are kept.
Hermaphrodite	Having both male and female sexual characteristics and organs.
Heteroblastic	Exhibiting differences in leaf shapes or forms in juvenile and adult phases of the plant.
Heteroblasty	The state of being heteroblastic (i.e., exhibiting differences in leaf shapes or forms in juvenile and adult phases of the plant).
Hirsute	Hairy.
Hyaline	Membranous, thin and translucent.
Hybrid	An individual that is the offspring of a cross between two different varieties or species.
Hybridise	Breeding with a member of a different plant or type.
Hydrophyte	A plant species adapted to growing in or on water or in wet situations. Aquatic or semi-aquatic.
Hymenium	The fertile, spore–bearing layer of a fruitbody.
Hypanthium	A ring–like, cup–shaped, or tubular structure of a flower on which the sepals, petals, and stamens are borne.
Imbricate	Overlapping.
imbricating	Overlapping.
Imparipinnate	Odd–pinnate, a leaf shape; pinnate with a single leaflet at the apex.
In-situ	On site conservation relating to the maintenance of plants in the wild.
Inbreeding	Genetic similarity in offspring of closely related individuals.

Term	Definition
Incoherent	Not sticking together.
Incursion	Entrance of a pest into an area where it is not present
Indumentum	A covering of fine hairs (or sometimes scales)
Indusia	Plural of indusium, a membrane covering a sorus of a fern
Indusium	A thin tissue that covers the sorus in many ferns. Plural: indusia.
Inflorescence	The arrangement of flowers on the stem. A flower head.
Infundibuliform	Funnel-like.
Interkeel	The space between the keel and the leaf blade
Internode	The part of an axis between two nodes; the section of the stem between leaves.
Internodes	Part of a stem between two nodes.
Intramarginal	Within or near the margin.
Involucral bracts	The scales surrounding the flower head or capitula.
Involucre	A group of bracts surrounding a flower head.
Involute	With margins rolled inward toward the upper side.
Irritable	Responding to touch.
Jugate	Paired.
Juvenile	A plant of non-reproducing size.
Keel	A prominent or obvious longitudinal ridge (as in a boat).
Labellar	Pertaining to the labellum: a lip; in orchid flowers referring to the middle petal which usually differs in size, shape or ornamentation from the two lateral petals.
Labellum	A lip; in orchid flowers referring to the highly modified middle petal which usually differs in size, shape or ornamentation from the two lateral petals.
Lacinia	A jagged lobe.
Laciniae	Jagged lobes.
Laciniate	Cut into narrow, irregular lobes or segments.
Lacustrine	Of or having to do with a lake, of, relating to, or formed in lakes, growing or living in lakes.
Lamina	The expanded flattened portion or blade of a leaf, fern frond or petal.
Lanceolate	Lance-shaped; of a leaf several times longer than wide with greatest width about one third from the base, tapering gradually to apex and more rapidly to base
Lateral	On or at the side.
Lax	With parts open and spreading, not compact.
Laxly	With parts open and spreading, not compact
Leaflet	One section of a compound leaf.
Lemma	The lower of two bracts enclosing the flower in grasses.
Lenticillate	Bark that is covered in fine lenticles (breathing pores)
Ligulate	Strap-like, tongue-shaped
Ligule	The membrane between the leaf and the stem of a grass; the "petal" of a ray floret in a composite inflorescence
Linear	Long and narrow with more or less parallel sides.
Littoral	Occurring at the border of land and sea (or lake). On or pertaining to the shore. The shallow sunlit waters near the shore to the depth at which rooted plants stop growing.
Lobe	A recognisable, but not separated, rounded division or segment of a leaf or pinna. Used to describe ferns and leaves in <i>Cotula</i> and <i>Leptinella</i> .
Lobed	Part of a leaf (or other organ), often rounded, formed by incisions to about halfway to the midrib.
Lobule	A small lobe or sub-division of a lobe
Lustrous	Glossy, shiny.
Lycophytes	Seedless vascular plants that belong to the phylum Lycophyta (characterised by microphylls -primitive leaves found in ancient plants).
Lyrate	Pinnatifid or pinnatisect terminal lobe much larger than lower lobes.
Maculate	Blotched or spotted.
Mangrove	Coastal wetland dominated by Manawa or mangrove <i>Avicennia marina</i> var. <i>resifera</i> . Northern New Zealand only, salt marsh replaces it further south.
Margin	The edge or border of a leaf
Marine	Pertaining to the sea and saltwater systems.
Marsh	A tract of wet land principally inhabited by partially-submerged herbaceous vegetation. Has fewer woody plants than swamper habitats.
Mealy	Dry, powdery, crumbly.
Median	In the middle.
Membranous	Very thin, like a membrane.
Mid-lobe	The middle part into which a leaf is divided.
Midrib	The central or principal vein of a leaf or pinna of a fern.
Mire	Synonymous with any peat-accumulating wetland. Term covers bogs and peaty swamps, fens, carr, moor, muskeg and peatland. Term excludes marsh which is non-peat forming.

Term	Definition
Molecular techniques	Where proteins and genes are used to investigate plant relationships
Monitoring	Recording of quantitative data over time to document changes in condition or state of species or ecosystems.
Monoecious	Having male and female flowers on the same plant of the same species.
Montane	Land between 300 and 800 metres above sea level.
Mucronate	Tipped with a short, sharp, point.
Mucronulate	Having a very small mucro; diminutive of mucronate.
Multi-annual evergreen	Overlapping annual cohorts of leaves always present.
Multifid	Cleft into many lobes or segments
Multiseptate	With many septa.
Mycorrhiza	A symbiotic relationship between a fungus and a plant.
Mycorrhizal associations	Symbiotic association between fungi and plant roots which assists plant health by allowing increased ability for uptake of nutrients and promote plant growth.
Napiform	A long swollen but tapering root – like a parsnip, or carrot.
Native	Naturally occurring in New Zealand (i.e., not introduced accidentally or deliberately by humans).
naturalised	Referring to plants that have escaped from cultivation (including gardens or forest plantations) and can now reproduce in the wild (without human assistance)
Nectary	Organ that produces nectar.
Nerve	Prominent vein or rib.
Nerves	Strands of conducting and usually strengthening tissue in a leaves or similar structures
Net veins	Veins that repeatedly divide and re-unite.
Net venation	Feather-like or hand-like venation on a leaf.
Nival	Growing at high altitudes. From Latin: nivalis, snowy etc. from nix, nivis, snow.
Node	The point at which leaves, branches or roots arise on a stem.
Ob-	Prefix meaning inverted, in reverse direction.
Obcordate	Heart shaped with the notch at the apex.
Oblanceolate	Tapering and widest towards the apex or inversely lanceolate.
Oblique	Slanting; of a leaf, larger on one side of the midrib than the other, in other words asymmetrical.
Oblong	Rectangular.
Obovate	Roughly elliptical or reverse egg shaped and widest near the apex (i.e., the terminal half broader than the basal half).
Obtuse	Blunt or rounded at the apex, with the sides meeting at an angle greater than 90°.
Operculate	With a small lid.
Opposite	A pair of organs attached at nodes in pairs on either side of a stem or axis.
Orbicular	Almost or approximately circular.
Outbreeding depression	A reduction in vigor of offspring from distant parents. It can occur when a locally adapted population is moved and mixed with plants adapted to different conditions.
Outer canopy deciduous	Marked reduction in leaf number in the outer canopy in exposed high light environments over winter.
Oval	Planar, shaped like a flattened circle, symmetrical about both the long and the short axis; about twice as long as broad, tapering equally both to the tip and the base. Synonymous with elliptical.
Ovary	Part of a flower containing the ovules and later the seeds.
Ovate	Egg-shaped and widest at base.
Ovoid	Oval; egg-shaped, with rounded base and apex.
Pakihi	A term which in its strict sense refers to open clears within forest dominated by low scrub and rushes. However, more usually used to refer natural and induced wetlands and their associated shrublands. A vernacular most frequently used in the West Coast for impoverished soils and their associated peats, left after forest has been cleared
Palea	The small upper bract enclosing the flower of a grass
Palmately	Radiating from a point, as fingers radiating from the palm of a hand.
Palmatifid	Deeply divided into several lobes arising from more or less the same level.
Palmatisect	Intermediate between palmate and palmatifid, i.e. the segments are not fully separated at the base; often more or less digitate.
Palustrine	Pertaining to wet or marshy habitats. Term covers mires and marshes
Pandurate	Fiddle-shaped.
Panicle	Highly branched (multiple raceme).
Papilla	A short rounded projection.
Papillae	A soft, fleshy projection, usually small and nipple-like.
Papillate	With short rounded projections.
Papillose	Warty, with short rounded projections or gland-dotted
Parallel venation	Veins are parallel along leaf.

Term	Definition
Parasite	An organism that derives all its nourishment from its host.
Patent	Spreading or expanded, e.g., spreading petals.
Peat	A mass of partially carbonised plant tissue formed by partial decomposition in water of various plants and especially of mosses of the genus Sphagnum, widely found in many parts of the world, varying in consistency from a turf to a slime used as a fertiliser, as stable litter, as a fuel, and for making charcoal. Partially carbonized vegetable matter saturated with water; can be used as a fuel when dried. A type of soil deriving from dead organic material situated in a wet area, where the reduced amount of [[oxygen available in the wet conditions results in the organic material not decomposing as much as it usually would do so in the presence of more oxygen. Used in growing media. Represents an important carbon sink –drainage of peat releases large amounts of carbon (CO ₂) to the atmosphere.
Pedicel	The stalk of a single flower in an inflorescence or fruit (either in a cluster or existing singularly).
Peduncle	The stalk of a solitary flower or the main stalk of an inflorescence or flower cluster.
Pedunculate	Describing fruits, which are borne on a stalk (a peduncle).
Pellucid	Transparent.
Peltate	Shield-like, with the stalk attached well inside the margin
Pendent	Hanging down from its support
Pendulous	Hanging or drooping.
Penicillate	With a tuft of hairs at the end, like a brush.
Perennial	A plant lasting for three seasons or more
Perianth	A collective term for the calyx (sepals or tepals) and corolla (petals) of the flower, especially when these are indistinguishable
Petal	Part of flower inside the sepals; usually coloured.
Petiolate	Having a petiole.
Petiole	Leaf stalk.
phloem	The vascular tissue in land plants that is primarily responsible for the distribution of sugars and nutrients manufactured in a shoot.
Photopoint	A monitoring technique where repeat photos are taken of the same scene from the same point over a period of time in order to quantify changes.
Pilose	Bearing long, soft hairs.
Pinna	A segment of a divided lamina that is classified as primary, secondary or tertiary according to the degree of dissection of the lamina.
Pinnae	Divisions of a pinnate leaf
Pinnate	With leaflets arranged regularly in two rows on either side of a stalk as in a feather; the lamina on a fern is divided into separate pinnae
Pinnatifid	Pinnately lobed, cleft more than halfway to the midrib. Not cleft all the way to the rachis.
Pinnatisect	Pinnately divided almost to midrib but segments still confluent.
Pioneer	Plant species are hardy species that should be planted first to establish a good canopy cover that restricts weed growth and promotes natural regeneration. In natural ecosystems these are the first plants to arrive and grow on a site.
Pistil	The female reproductive organ of a flower, consisting of an ovary, style, and stigma.
Pistillate	A flower with one or more pistils, but no stamens.
Plano-convex	Flat on one side, convex on the other.
Plumose	Feathery.
Podzol	Infertile, acidic soil, strongly leached to form a whitish-grey subsoil underlain by a layer enriched in iron, aluminium and organic matter; usually under forest in a wet temperate climate.
Pole	A subcanopy size individual with a long thin trunk and foliage tuft of a potential canopy tree.
Pollinia	Compact masses of orchid pollen.
Population enhancement	Increasing a population for a specific biological purpose, e.g., when a species is already present in an area but extra individuals are added to address a sex imbalance.
Porrect	Extending forward.
Procumbent	Lying and flat along the ground but not rooting
Propagate	To reproduce a plant by sexual (i.e., from seed) or asexual (e.g., from cuttings) means.
Prostrate	A general term for lying flat along the ground. This includes procumbent (that is lying and flat along the ground but not rooting) and decumbent (with a prostrate or curved base and an erect or ascending tip).
Provenance	The place of origin (of a plant that is in cultivation).
Proximal	Toward the base or point of attachment (cf. distal).
Pseudobulb	Thickened surface stem; usually looking like a bulb.
Pseudoterminal	Falsely terminal – as in a bud which appears to occupy a terminal position but does not
Puberulent	Minutely clad in short, soft hairs
Pubescence	Covering of soft, fine hairs
Pubescent	Covered in short, soft hairs.
Pungent	Ending in a stiff sharp point
Pustule	Small blister-like elevation.

Term	Definition
Quadrat	Square, rectangular.
Raceme	An unbranched, elongated inflorescence with pedicellate flowers maturing from the bottom upward i.e., flowers attached to the main stem by short stalks.
Rachis	the axis of an inflorescence or of a compound leaf
Ray	An outer ring of strap-like florets in the head of Asteraceae (daisy) flowers.
Re-introduction	Translocating wild or cultivated individuals to sites where the taxon has been known to occur in the past, but from which it has disappeared.
Recurved	Curved backward.
Reflexed	Bent back on itself
Reniform	Kidney shaped.
Repand	With a slightly wavy margin.
Replum	The outer structure of a pod in which the valves have dehisced (persists after the opening of the fruit)
Restiad	Area dominated by rush-like plants (collectively known as restiads) of the family Restionaceae. Includes Chatham Island and North Island Sporodanthus and oioi (<i>Apodasmia similis</i>)
Retorse	Pointing backward.
Retuse	A shallow notch at the rounded or blunt apex of a leaf.
Rhizoid	Any of various slender filaments that function as roots in mosses and ferns and fungi.
Rhizomatous	With underground creeping stems.
Rhizome	An underground stem (usually spreading horizontally or creeping) or short and erect.
Rhombic	Diamond-shaped.
Rhomboid	Diamond shaped, nearly rhombic.
Riparian	Relating to or living or located on the bank of a natural watercourse (as a river) or sometimes of a lake or a tidewater.
Riparian margin	Refers to the edges of streams, rivers, lakes or other waterways.
Riparian plants	Refers to plants found growing near the edges of streams, rivers or other waterways.
Riparian zone	A strip of land next to streams, rivers, and lakes where there is a transition from terrestrial (land vegetation) to aquatic (water) vegetation. Also known as "berm".
Riverine	Pertaining to rivers, streams and such like flowing water systems.
Rootstock	A short, erect, underground stem.
Rosette	A radiating cluster of leaves.
Rostellum	In orchids, a modified stigma that prevents self-fertilisation.
Rosulate	A dense radiating cluster of leaves.
Rugose	Wrinkled.
Rugulose	Having small wrinkles.
Runcinate	Sharply pinnatifid or cleft, the segments directed downward.
Runner	A trailing stem that roots at the nodes.
Rupestral	Growing on rocks.
Rushes	A group of distinctive wetland plants. They have solid stems (grasses have hollow stems), true rushes <i>Juncus</i> sp. have rounded leaves.
Sagittate	Shaped like the head of an arrow; narrow and pointed but gradually enlarged at base into two straight lobes directed downwards; may refer only to the base of a leaf with such lobes; cf. hastate.
Salt marsh	A coastal wetland, with specialized salt tolerant plants (halophytes).
Sapling	A juvenile tree that has reached the stage of 1 or 2 main stems but is still in the shrub layer.
Saprophyte	A plant lacking chlorophyll and living on dead organic matter.
Saprophytic	Lacking chlorophyll and living on dead organic matter.
Sarcotesta	The fleshy, often highly coloured outer layer of the seed coat in some species, e.g., titoki (<i>Alectryon excelsus</i>).
Scabrid	Roughened or rough with delicate and irregular projections.
Scale	Any thin, flat, membranous structure.
Scape	A leafless flower stem.
Scutiform	Shield-shaped.
Sedges	A group of grass-like or rush-like herbaceous plants belonging to the family Cyperaceae. Many species are found in wetlands some are forest floor plants. Leaves are usually angular. Hence the saying "rushes are round and sedges have edges".
Seedling	A newly germinated plant.
Self sustaining	Able to sustain itself, or replace itself, independently of management i.e. regenerate naturally
Self thinning	Natural tree death in a crowded, even-aged forest or shrubland.
Semi-deciduous	Partial leaflessness in winter, and greater than 50% leaves lost by the beginning of spring flush.
Sepal	Outer part of flower; usually green.
Serrate	Sharply toothed with teeth pointing forwards towards apex.
Serrulate	Finely serrate, i.e., finely toothed with asymmetrical teeth pointing forward; like the cutting edge of a saw.

Term	Definition
Sessile	Attached by the base without a stalk or stem.
Seta	The stalk of a fruiting moss capsule
Sheath	A portion of an organ that surrounds (at least partly) another organ (e.g., the tubular envelope enclosing the stem in grasses and sedges).
Silicles	The flattened usually circular capsule – compared with the narrow, elongated fruit (silique) – containing the seed/seeds. A term used almost exclusively for plants within the cabbage family (Brassicaceae)
Silique	A capsule, usually 2-celled, with 2 valves falling away from a frame (replum) bearing
Simple	Of one part; undivided (cf compound).
Sinuate	With a wavy margin.
Sinus	The space or recess between lobes; in hebes a gap between the margins of two leaves of an opposite pair that may be present in the bud before the pair of leaves separate.
Sorus	A cluster of two or more sporangia on the margin or underside of the lamina of a fern, sometimes protected by an indusium.
Spathulate	Spatula or spoon-shaped, a rounded blade tapering gradually to the base.
Spheroidal	Almost spherical but elliptic in cross section.
Spicate	Arranged in a spike.
Spike	Flowers attached to main stem without stalks.
Spikelet	Collection of individual grass florets borne at the end of the smallest branch of the inflorescence.
Sporangia	Plural of sporangium. Structures in which spores are produced.
Sporangium	Structure in which spores are produced.
Spore	A single-celled reproductive unit similar in function to that of the seed in a flowering plant.
sporophyte	The spore producing plant in ferns that is usually the visible part.
Stamen	The male reproductive organ of a flower where pollen is produced. Consists of an anther and its stalk.
Stamens	The male, pollen bearing organ of a flower.
Standing water	Where water lies above the soil surface for much of the year.
Stellate	Irregularly branched or star shaped.
Stigma	Female part of the flower that is receptive to pollen, usually found at or near the tip (apical end) of the style where deposited pollen enters the pistil.
Stipe	The stalk of a frond.
Stipitate	Borne on a stipe or stalk.
Stipulate	A leaf with stipules.
Stipule	A scale-like of leaf-like appendage at the base of a petiole, usually paired.
Stolon	A stem which creeps along the ground, or even underground.
Stoloniferous	Producing stolons
Stramineous	Chaffy, like straw or straw-colored.
Stria	A fine line or groove.
Striae	Fine lines or grooves.
Striate	Fine longitudinal lines or minute ridges
Style	The elongated part of the flower between the ovary and the stigma.
Sub-	A prefix meaning under, somewhat or almost.
Subglabrous	Very slightly, but persistently, hairy.
Suborbicular	Slightly rounded in outline
Substrate	The surface upon which an orchid grows.
Subtended	Immediately beneath, occupying a position immediately beneath a structure, i.e., flower subtended by bract
Subulate	Slender and tapering to a point.
Succession	Progressive replacement of one species or plant community type by another in an ecosystem.
Successional	Referring to species, plant communities or habitats that tend to be progressively replaced by another.
Succulent	Fleshy and juicy.
Summer-green	Used in New Zealand to indicate herbs or sub-shrubs that die down to a root stock or rhizomatous network.
Supplementary planting	Returning to a revegetation site and creating gaps, or filling existing gaps, with different plants of plants, usually later successional plants which may not have survived being planted in the first phases of the project.
Surface water	Water present above the substrate or soil surface.
Surveillance	Regular survey for pests inside operational and managed areas e.g. nurseries, standout areas on parks.
Survey	Collection of observations on the spatial distribution or presence or absence of species using standardised procedures.
Sustainable Land Management	The use of farming practices which are sustainable both financially and environmentally including management of nutrient runoff, waste disposal or stock effluent, reducing impacts of nutrients on waterways, preventing erosion and soil loss, and protecting native forest and wetland habitats from stock damage.
Swamp	Low land that is seasonally flooded; has more woody plants than a marsh and better drainage than a bog. They are more fertile and less acidic than bogs because inflowing water brings silt, clay and organic matter. Typical swamp plants include raupo, purei and harakeke (flax). Zonation and succession often leads through manuka to kahikatea swamp forest as soil builds up and drainage improves.

Term	Definition
Symbiote	An organism that has an association with organisms of another species whereby the metabolic dependence of the two associates is mutual.
Symbiotic	The relation between two different species of organisms that are interdependent; each gains benefits from the other (see also symbiosis).
Sympatric	Occupying the same geographical region.
Synangia	Structures made up of fused sporangia
Synonym	A botanical name that also applies to the same taxon.
Systematics	The study of taxonomy, phylogenetics, and taxogenetics.
Tabular	Shaped like a rectangular tablet.
Taxa	Taxonomic groups. Used to refer to a group at any level e.g., genus, species or subspecies.
Taxon	A taxonomic group. Used to refer to a group at any level e.g., genus, species or subspecies.
Taxonomy	The process or science of classifying, naming, and describing organisms
Tepal	An individual member of the perianth.
Terete	Cylindrical and tapering.
Terminal	At the tip or apex.
Ternatifid	Leaflets In threes,
Tetrad	A group of four.
Tomentum	A hairy covering of short closely matted hairs.
Translocation	The movement of living organisms from one area to another.
Trifid	Divided into three.
Trifoliate	Having three leaflets.
Trigonous	Three-angled
Tripinnate	With each secondary pinna divided to the midrib into tertiary pinnae
Triquetrous	Triangular in cross section and acutely angled.
Truncate	With the apex or base squared at the end as if cut off.
Tuberculate	Bearing small swellings.
Tubular	Tube-shaped.
turbinate	Top-shaped.
Turgid	Distended through internal pressure
Type locality	The place or source where a holotype or type specimen was found for a species.
Ultramafic	A type of dark, usually igneous, rock that is chemically dominated by magnesium and iron-rich minerals, the partially metamorphosed form of which is serpentinite.
Umbel	Umbrella like; the flower stalks arise from one point at the stem.
Undulate	Wavy edged.
Undulose	Wavy edged.
Unitubular	A tube partitioned once – literally one tube (compare – multitubular – many tubes)
Utricle	A thin loose cover enveloping some fruits (eg., Carex, Uncinia)
Valvate	Opening by valves.
Vascular plant	A plant that possesses specialised conducting tissue (xylem and phloem). This includes flowering plants, conifers and ferns but excludes mosses, algae, lichens and liverworts.
Velutinous	Thickly covered with delicate hairs; velvety.
Ventral	Of the front or inner (adaxial) surface relative to the axis. (cf. dorsal)
Vermiform	Worm-shaped.
Vernicose	Glossy, literally as if varnished, e.g., Hebe vernicosa has leaves that appear as if varnished
Verrucose	Having small rounded warts.
Verticillium	A fungus disease that will cause wilting and death.
Villous	Covered with long, soft, fine hairs.
Water table	The level at which water stays in a soil profile. The zone of saturation at the highest average depth during the wettest season.
Wetland	A site that regularly has areas of open water for part or all of the year, or has a water table within 10 cm of the surface for at least 3 months of the year. Wetland ecosystems support a range of plant and animal species adapted to an aquatic or semi-aquatic environment.
Whipcord	A shrub in which the leaves are reduced to scales that are close-set and pressed against the stem.
Whorl	A ring of branches or leaves arising at the same level around the stem of a plant.