

INDIGENOUS HIGHER PLANTS (PSILOPSIDS, LYCOPODS, FERNS, GYMNO-  
SPERMS, FLOWERING PLANTS) OF AORANGI RANGE, SOUTHERN  
WAIRARAPA, SEA-LEVEL TO 3226 FT/981 M


Highest peak: Mt Ross, grid ref. S28 062693  
(Biosis c.h.)

A.P. Druce, Botany Division, D.S.I.R., June 1974

(Revised July 1975, July 1976, May 1977, Oct. 1977, June 1980,  
Oct. 1982, May 1983, Oct. 1986, Nov. 1986, June 1988, Aug. 1990,  
Nov. 1990, May 1991, Sept. 1991, July 1992)

(Originally published in Bull. Wellington Bot. Soc. 37, 4-29, 1971)

(Rewritten and extended to the coast Aug. 1990)  
(Based on observations 1947-1992)



- ① - Putangitua Stm
- ② - Makotukutuku Stm
- ③ - Mangatoetoe Stm
- ④ - Little Mangatoetoe Stm
- ⑤ - Waitetuna Stm
- ▨ Tertiary siltstone and sandstone

(N) - found only in northern part of Range

(S) - found only in southern part of Range

(L) - found only in limestone area, Ruakohoputuna V.

(C) - found only near coast

(unc) - uncommon ; (/) - not found above 300m, or rarely so

### GYMNOSPERM TREES

*Dacrycarpus dactyloides* (unc)

*Dactyidium cupressinum*

*Podocarpus hallii*

*P. totara* s.s. (unc)

*Prumnopitys ferruginea*

*P. taxifolia*

*Metrosideros robusta* (two live trees seen 1982/3, one dead one in 1972, and two dead ones in 1974)

*Myoporum laetum* var. *laetum*

*Myrsine australis*

*M. salicina*

*Nestegis cunninghamii*

*N. lanceolata* (unc)

*N. montana*

*Nothofagus fusca* (not present south of Mt Barton)

*N. menziesii*

*N. solandri* var. *solandri*

*N. truncata* (from Putangirua, Tauanui and Dry river valleys north-westwards)

*N. fusca* X *N. solandri*

*N. fusca* X *N. truncata* (unc N)

*N. solandri* X *N. truncata*

*Olearia paniculata*

*O. rani* var. *colorata*

*Pennantia corymbosa*

*Pittosporum eugenioides*

*P. tenuifolium* ssp. *tenuifolium*

*Plagianthus regius* var. *regius* (/)

*Pseudopanax arboreus* var. *arboreus*

*P. colensoi* s.s. (found mainly on cliffs or as an epiphyte, but a number of plants present in scrub on Mangataetoe in 1986)

*P. crassifolius*

*P. edgerleyi* (unc)

*P. simplex* (incl. *P. s.* var. *sinclairii*)

*Schefflera digitata*

*Sophora microphylla*

### MONOCOT TREES AND SHRUBS

*Cordyline australis*

*C. banksii*

*C. indivisa* (unc)

*C. australis* X *C. banksii* (unc)

*Rhopalostylis sapida* (/)

### DICOT TREES

*Alectryon excelsus* var. *excelsus* (/)

*Aristotelia serrata*

*Beilschmiedia tawa* (unc/)

*Carpodetus serratus*

*Corynocarpus laevigatus* (/)

*Dodonaea viscosa* var. (/)

*Elaeocarpus dentatus* (incl. *E. d.* var. *obovatus*)

*E. hookerianus* (unc)

*Entelea arborescens* (unc C)

*Fuchsia excorticata*

*Griselinia littoralis*

*G. lucida*

*Hedycarya arborea*

*Hoheria populnea* var. *lanceolata*

*Knightia excelsa*

*Laurelia novae-zelandiae* (unc/)

*Lophomyrtus obscurata* (/)

*Macropiper excelsum* var. *excelsum*

*Melicope ternata* (unc C)

*Melicytus ramiflorus* ssp. *ramiflorus*

*Streblus heterophyllus* (1)

*Weinmannia racemosa* var. *racemosa*

### DICOT SHRUBS

- Alseuosmia pusilla* (unc)  
*Brachyglottis grayi* var. *grayi* (S - south of Mt Barton)  
*B. repanda* s.s.  
*B. rotundifolia* var. (*B. elaeagnifolia*) (mainly south of Mt Barton)  
*Carmichaelia* sp. (subgenus *Carmichaelia*)  
*Cassinia leptophylla* var. *leptophylla*  
*Coprosma acerosa* ssp. *acerosa* (unc C)  
*C. atrolata*  
*C. grandifolia*  
*C. colensoi* s.s.  
*C. crassifolia*  
*C. foetidissima* (unc)  
*C. linariifolia* (unc N)  
*C. lucida* s.s.  
*C. microcarpa*  
*C. propinqua* ssp. *propinqua*  
  
*C. repens* (C)  
*C. rhamnoides* (incl. *C. polymorpha*)  
  
*C. rigida*  
*C. robusta*  
*C. rotundifolia*  
*C. rubra*  
*C. virescens* (unc / S - Waitatuna and Mangatoetoe Stms)  
*C. sp.* (unnamed; included in *C. parviflora* by Oliver, 1935, and others) (If surface not minutely hairy as in usual in *C. parviflora*; drupes usually unpigmented, occasionally purplish red)  
*C. propinqua* X *C. robusta* (unc)  
*C. repens* X *C. rhamnoides* (unc C)

*Coriaria arborea*

- C. sarmentosa* (unc S - south of Mt Barton)  
*C. sp.* (unnamed; aff. *C. plumosa* and *C. pteridoides*) (S - Mangatoetoe Stm)  
*C. arborea* X *C. sarmentosa* (unc S)  
*C. arborea* X *C. sp.* (unnamed) (unc S)  
*Cyathodes sp.* (*C. juniperina* agg.) (lvs small, vs.  $\angle$  1cm X  $\angle$  1mm)  
*Dracophyllum longifolium* var. (incl. *D. filifolium* and *D. oliveri*) (mainly south of Mt Barton)  
*Fuchsia perscandens* X *F. excorticata* (N. - Ruakokoputuna V.)  
*Gaultheria antipoda*  
*G. rupestris* (incl. *G. rupestris* var. *subcorymbosa*) (mainly south of Whananui)  
*G. antipoda* X *G. rupestris*  
*Geniostoma rupestre* var. *ligustifolium* (unc)  
*Hebe stricta* var. *atkinsonii*  
*H. stricta* var. *macrooura* (C)  
*H. stricta* var. *atkinsonii*  $\rightleftharpoons$  var. *macrooura* (C)  
*H. venustula* (incl. *H. brachysiphon*) (S)  
*H. sp.* (*Veronica arborea*)  
*Helichrysum aggregatum* var. *aggregatum*  
 [Helichrysum hybrids - see under DICOT HERBS]  
*Korthalsella satiocornoides* (1)  
*Kunzea ericoides* agg.  
*Leptospermum scoparium*  
*Leucopogon fasciculatus*  
*L. fraseri* s.s. (mainly south of Mt Barton)  
*Melicope simplex*  
*Melicytus crassifolius* var. *crassifolius* (1)  
*Oleaia arborescens*  
*O. solandri* (1)

- Pimelea gnidia* (S - south of Mt Barton)
- P. longifolia* (S - south of Mt Barton)
- P. prostrata*
- P. sp.* (cf. *P. urvilleana*) (incl. *P. prostrata* var. *quadrifaria*)
- Pittosporum cornifolium* (unc)
- Plagianthus divaricatus* (C)
- Pomaderris ericifolia* (N)
- Pseudopanax anomalus* (unc)
- Pseudowintera colorata*
- Solanum aviculare* var. *aviculare*
- S. laciniatum* (N)
- Urtica ferox*

**MONOCOT LIANES**

- Freycinetia bavariana* ssp. *banksii* (I)
- Ripogonum scandens*

**DICOT LIANES AND RELATED PLANTS**

- Calystegia soldanella* (C)
- C. tuguriorum*
- Clematis afoliata* (common in Makotukotuku V, unc. in Little Mangataotoe V.)
- C. foetida*
- C. forsteri*
- C. paniculata*
- C. afoliata* x *C. foetida* (unc)
- Convolvulus sp.* (unnamed; included in *C. vetecundis*, as ssp. *waitaha*, by Sykes 1987)
- Metrosideros colensoi* (incl. *M. c.* var. *pendula*) (S)
- M. diffusa*
- M. perforata*
- Muehlenbeckia australis*
- M. complexa*
- M. ephedroides* (unc C)
- M. australis* x *M. complexa* (unc)
- Parsonsia capsularis* var.
- P. heterophylla*
- Passiflora tetrandra*
- Robus cissoides* var. *cissoides*
- R. schmidelioides* var. *schmidelioides*
- R. squarrosus*
- Tetragonia trigyna* (C)
- P. capsularis* x *P. heterophylla* (unc)

**PSILOPSIDS AND LYCOPODS**

- Tmesipteris elongata* (incl. *T. elongata* ssp. *robusta*) (unc)
- Lycopodium fastigiatum* (S - south of Mt Barton)
- L. scariosum*
- L. varium* (incl. *L. billardierei* and *L. novae-zelandicum*)
- L. volubile*

**FERNS**

- Adiantum cunninghamii*
- A. diaphanum* (unc I)
- Anarthropteris lanceolata* (I)
- Anogramma leptophylla* (unc I)
- Asplenium bulbiferum*
- A. flabellifolium* agg.
- A. flaccidum*
- A. gracillimum*
- A. hookerianum*
- A. lyallii* (L)
- A. oblongifolium*
- A. polyodon*
- A. terrestre* ssp. *maritimum* (C)
- A. terrestre* ssp. *terrestre*
- A. bulbiferum* x *A. hookerianum*
- A. flaccidum* x *A. gracillimum*
- A. flaccidum* x *A. terrestre* (unc)
- A. gracillimum* x *A. hookerianum*
- A. hookerianum* x *A. lyallii* (unc L)
- A. lyallii* x *A. oblongifolium* (unc L)
- Azolla filiculoides*
- Blechnum banksii* (incl. *B. durum*)
- B. chambersii*
- B. colensoi* (unc)
- B. discolor*
- B. filiforme*
- B. fluviatile* agg.
- B. membranaceum* (I)
- B. minus*
- B. penna-marina*
- A. sp.* (unnamed; aff. *A. trichomanes*) (unc L)

- B. procerum
- B. vulcanicum (unc)
- B. sp. (i) (unnamed; common lowland sp.) (lower pinnae reduced in length)
- B. sp. (ii) (unnamed; cliff sp.) (lower pinnae not reduced in length; "The Green Bay Species" of Crookes 1963)
- B. sp. (iii) (unnamed; mountain sp.) (lower pinnae reduced in length) ("The Dunedin-Cass Variety" of Crookes 1963)
- B. chambersii X B. membranaceum (/)
- B. procerum X B. sp. (i)
- Bettrichium biforme (unc)
- Cheilanthes distans (unc /)
- C. homilis (/)
- Ctenopteris heterophylla
- Cyathea colensoi (unc)
- C. dealbata
- C. medullaris
- C. smithii
- Dicksonia fibrosa (N - Ruakohoputona V.)
- D. squarrosa
- Grammitis billardierei
- G. ciliata (unc)
- G. poeppigiana (S - south of Mt Barton)
- Histiopteris incisa
- Hymenophyllum bivalve
- H. cupressiforme
- H. demissum
- H. dilatatum (unc)
- H. flabellatum
- H. minimum (unc S - south of Mt Barton)
- H. multifidum
- H. peltatum (unc)
- H. ratum
- H. revolutum (unc)
- H. sanguinolentum agg.
- H. scabrum (/)
- H. pulcherrimum (unc)
- G. magellanica ssp. nothofageti

- Hypolepis ambigua
- H. millefolium (unc S - south of Mt Barton)
- H. rufobarbata
- H. ambigua X H. rufobarbata
- Lastreopsis glabella (/)
- L. hispida (/)
- L. microsora ssp. pentangularis (unc /)
- L. velutina (/)
- Leptolepia novae-zelandiae (unc)
- Ophioglossum ceriaceum agg. (unc S - south of Mt Barton)
- Paesia scaberula
- Pellaea calidrupium (unc c)
- P. rotundifolia
- Phymatosorus pustulatus
- Pleurosorus rotifolius (unc)
- Pneumatosorus pennigera
- Polystichum richardii
- P. silvaticum
- P. vestitum
- P. richardii X P. vestitum (unc)
- P. silvaticum X P. vestitum (unc)
- Pteridium esculentum
- Pteris pendula
- P. tremula (/)
- Pyrosia alcagnifolia
- Rumohra adiantiformis
- Trichomanes reniforme
- T. venosum (unc)
- { Leptopteris hymenophylloides
- { L. superba
- { L. hymenophylloides X L. superba

ORCHIDS

- Acianthus sinclairii (/)
- Caladenia carnea agg.
- Chiloglottis cornuta
- Corybas macranthus
- C. rivalaris ss. (unc /)
- C. trilobus ss.
- Cyrtostylis reniformis ss. (unc /)

*P. cardiostigma* (unc)*Dendrobium cunninghamii**Drymeanthus adversus* (unc)*D. sp.* (unnamed) (unc - Mt Surf)*Earina autumnalis**E. mucronata* s.s.*Gastrodia cunninghamii**Genoplesium nudum* (unc/)*Microtis unifolia**Orthoceras novae-zeelandiae* (unc/)*Prasophyllum colensoi* (unc S)*Pterostylis alobula* (/)*P. foliata* (unc/)*P. banksii**P. graminea**P. irsoniana* (unc S - south of Mt

Barton)

*P. sp.* (unnamed; aff. *montana*) (unc N)*P. trullifolia* (unc/)*Thelymitra hatchii* (unc)*T. longifolia**T. pauciflora**T. pulchella* (unc/)*T. sp.* (aff. *T. ixioides*) (unc/)*Festuca multinodis**Hierochloa pedolens**Lachnagrostis filiformis* s.s. (unc/)*L. richardii**L. sp.* (unnamed; included in *L. filiformis*, as var. *semiglabra*, by Zotou 1965) (unc L)*Microlaena avenacea**M. stipoides**Poa anceps* ssp. *anceps**P. breviglumis**P. cita* ssp. (lvs narrow, folded)*P. colensoi* (unc S - south of Mt

Barton)

*P. imbecilla**P. anceps* x *P. cita* (?)*Rytidosperma buchananii* (unc)*R. clavatum**R. gracile**R. merum* (unc L)*R. petrosum* (/)*R. unarede**Spinifex sericeus* (C)*Trisetum sp.* (i) (unnamed; aff. *T. antarcticum*)*T. sp.* (ii) (unnamed) (plants slender; lf sheath glabrous or nearly so)*Zoysia minima* (C)**GRASSES***Anemantliele lessoniana**Austrofestuca littoralis* (C)*Chionochoa beddiei* (S - south of

Mt Barton)

*C. cheesemani**C. conspicua* ssp. *cunninghamii*

(unc - Mt Surf and Mt Barton)

*C. beddiei* x *C. cheesemani* (unc)*Cortaderia fulvida**C. toetoe**Deyeuxia avenoides* (incl. *D. avenoides* var. *brachyantha*)*D. billardièrei* (C)*Dichelachne ctinita**Echinochloa ovatus**Elymus multiflorus* (Hooker 1864, as *Triticum multiflorum*)*E. sp.* (aff. *E. rectisetus*)**SEDGES***Baumea rubiginosa* (C)*Carex breviculmis**C. buchananii* (unc C)*C. dissita* s.s.*C. flagellifera**C. forsteri**C. geminata* (unc)*C. lambertiana**C. lessoniana* (unc)*C. pumila* (C)*C. raouli**C. resectans* (unc C)

*S. maschalinus* (1)*C. secta* ss. (1)*C. solandri**C. spinulosus* (unc 1)*C. testacea* ss.*C. virgata* (unc 1)*Cyperus ustulatus* (C)*Dasymaschonus spiralis**Eleocharis acuta* (unc)*E. gracilis* (unc)*Gahnia pauciflora* (unc)*Isolopis cernua* (C)*I. habra* (unc N - Te Maunga ridge)*I. nodosa* (C)*I. pottsii**I. prolifer* (C)*I. reticularis* (unc)*Schoenoplectus pungens* (C)*Schoenus apogon* (unc N)*S. nitens* (Hooker 1864)*Uncinia banksii* (1)*U. clavata**U. ferruginea**U. filiformis* (N)*U. gracilentata* (unc N)*U. laxiflora* (unc)*U. leptostachya**U. rupestris* (incl. *U. angustifolia*)*U. scabra**U. silvestris* (incl. *U. affinis*)*U. uncinata**U. zotovii**Leptocarpus similis* (C)*Luzula banksiana* ss. (C)*L. migrata* (S - south of Mt Barton)*L. picta* ss.*L. subclavata* (unc N)*L. sp.* (unnamed; included in *L. picta*, as var. *limosa*, by Edgar 1966) (unc)*L. banksiana* X *L. picta**L. migrata* X *L. picta* (unc)MONOCOT HERBS OTHER THAN  
ORCHIDS GRASSES, SEDGES, RUSHES*Arthropodium candidum**Astelia fragrans**A. solandri**A. sp.* (unnamed; aff. *A. nervosa*)(lvs not conspicuously white ad-axially as in *A. nervosa*)*Collospermum hastatum* (unc)*C. microspermum**Dianella nigra**Libertia grandiflora* (plants strongly rhizomatous and forming large patches)*L. ixoides**Lemna sp.* (*L. minor* auct. N.Z.)*Luzuriaga parviflora* (unc N - Te Maunga and Mt Ross ridges)*Phormium cookianum**P. tenax* (C)*P. cookianum* X *P. tenax* (C)*Triglochin striata* (C)*Typha orientalis* (C)RUSHES*Juncus antarcticus* (unc S - Mt Barton)*J. australis**J. caespitosus* (C)*J. distegus**J. gregiflorus**J. novae-zelandiae* (unc)*J. pallidus**J. planifolius**J. sarophorus**J. sp. or hybrid* (undetermined) (unc)COMPOSITE HERBS*Anaphalis keriensis* ss.*A. subrigida* (L)*A. sp.* (*Helichrysum bellidoides* s.s.)[*Anaphalis hybrids* - see below under *Helichrysum*]*Brachyglottis lagopus* var. *lagopus**Brachyscome radicata* (S)*Celmisia spectabilis* ssp. *laucedata* (S)

G. involucratum X G. limosum(?) (unc)

- Cotula australis
- C. coronopifolia (C)
- Craspedia minor var. minor (incl. C. major)
- C. uniflora var. grandis
- C. viscosa (unc L)
- Gnaphalium aurdax
- G. gymnocephalum
- G. involucratum (unc)
- G. limosum s.s. (unc)

- Senecio glomeratus (I)
- S. hispidulus (I)
- S. laetus ssp. laetus
- S. minimus
- S. quadridentatus
- S. rufiglandulosus var. rufiglandulosus (S - Mt Barton)
- Sonchus kirki (C)
- Vittadinia australis agg.

G. sphaericum

- Helichrysum filicaule
- Helichrysum - Anaphalis - Raoulia hybrids:
- (A) - between early-flowering spp.
- Anaphalis keriensis X A. subtrigida (unc L)
- A. keriensis X A. sp. (Helichrysum bellidioides s.s.) (unc)
- A. subtrigida X Raoulia tenuicaulis (unc)
- A. sp. (Helichrysum bellidioides s.s.) X Helichrysum aggregatum (unc)
- (B) - between late-flowering spp.
- Helichrysum filicaule X Raoulia glabra (unc)

- Lagenifera pumila
- L. strangulata
- Leptinella pusilla (unc C)
- Microseris scapigera var. (C)
- Pseudognaphalium sp. (I) (unnamed; aff. P. luteoalbum) (lvs mostly linear-spathulate to linear, acute)

- P. sp. (ii) (unnamed; aff. P. luteoalbum) (lvs mostly oblong-spathulate to narrowly oblong, rounded) (C)

- Raoulia australis agg.
- R. glabra
- R. tenuicaulis (incl. R. t. var. dimorpha and R. t. var. pusilla)
- [Raoulia hybrids - see above under Helichrysum]

DICOT HERBS OTHER THAN COMPOSITES

- Acaena ansetinifolia
- A. juvenca
- Aciphylla squarrosa s.s.
- Apium prostratum ssp. (C)
- A. sp. (slender) (C)
- Australina pusilla (unc I)
- Cardamine sp. (i) (C. debilis agg.) ("Long Style" of Pritchard 1957)
- C. sp. (ii) (C. debilis agg.) ("Narrow Petal" of Pritchard 1957)
- C. sp. (iii) (C. debilis agg.) ("Glossy Leaf" of Pritchard 1957)
- C. sp. (iv) (cf. C. corymbosa) ("Mainland Coastal Race" of Pritchard 1957) (unc N)
- Centella uniflora (I)
- Chenopodium glaucum ssp. ambiguum (C)
- Colobanthus muelleri (C)
- C. strictus
- Crassula mataikona (unc)
- C. moschata (unc C)
- C. pedunculatis (C)
- C. sieberiana
- Daucus glochidiatus (unc I)
- Dichondra repens var.
- D. sp. (i) (lvs 5-10 mm wide, emarginate, slightly fleshy, both surfaces initially covered with scattered, appressed hairs, becoming glabrous; fls 6-13 mm diam.; sepals < corolla lobes, between which they are often only just visible; corolla lobes 1.5-4 mm wide, rounded, creamy white, margins smooth, or irregularly



lobed or-toothed; peduncles up to 2-3cm long, reflexed at fruiting stage) (C)  
 D. sp. (1) (unnamed) (stems slender 0.5-1mm diam.; lvs 5-13-(20)mm wide, very slightly emarginate, both surfaces covered with scattered hairs, erect on upper surface, appressed on lower surface; fls 4-5mm diam.; sepals = or > corolla lobes, between which they are conspicuous; corolla lobes c. 1mm wide, subacute, very pale creamy white; peduncles up to 2.7cm long, reflexed at fruiting stage) (unc)

- Disphyma australe ssp. australe (C)
- Drosera peltata ssp. auriculata (N)
- Einadia allanii (C)
- E. triandra (C)
- Epilobium alsinoides s.s. (unc)
- E. atriplicifolium s.s. (mainly S of Mt Barton)
- E. brunnescens s.s.
- E. cinereum (1)
- E. cockayneanum (unc S - Mangatootoe)
- E. glabellum s.s. (unc S - south of Mt Barton)
- E. insulare (unc)
- E. komarovianum
- E. linnaeoides
- E. microphyllum
- E. nerteroides
- E. nummularifolium
- E. pedunculare s.s. (unc)
- E. pubens
- E. rotundifolium
- Eryngium vesiculosum
- Euphasia cuneata
- Galium propinquum
- G. trilobum (C)
- Geranium <sup>sp. (unnamed; aff. G. microphyllum)</sup>
- G. sessiliflorum var. novae-zelandiae
- Gonocarpus aggregatus (unc S - south of Mt Barton)

- G. micranthus ssp. micranthus (unc)
- G. montanus (unc)
- Gunnera monoica
- Haloragis erecta ssp. erecta
- Hydrocotyle elongata
- H. heterometra
- H. hydrophila (unc L)
- H. microphylla
- H. moschata
- H. novae-zeelandiae s.s. (1)
- Jovellana repens (unc)
- Lilaeopsis sp. (C)
- Limosella lineata
- Linum monogynum
- Lobelia anceps
- Mazus novaezeelandiae (C)
- Montia fontana s.s. (C)
- Miyosotis forsteri (unc)
- M. sp. (unnamed; included in M. pygmaea, as var. minutiflora, by Simpson and Thomson, 1943) (unc C)
- Myosurus minimus ssp. novae-zeelandiae (Hooker 1864, as M. australis; Garnock-Jones 1986)
- Myriophyllum sp. (unc C)
- Nerteta depressa (incl. N. cunninghamii)
- N. sp. (unnamed; aff. N. dichondrifolia) (hairs present on both surfaces of lf, straight, not hooked as in N. dichondrifolia)
- Oreomyza sp. (unnamed; aff. O. ramosa) (unc S - near Mt Barton)
- Oxalis exilis
- O. magellanica
- Parakebe diffusa ssp. diffusa (unc S - south of Mt Barton)
- Parietaria debilis (1)
- Plantago masoniae (unc C)
- P. raoulii
- P. spathulata ssp. spathulata (S - south of Mt Barton)
- P. raoulii x P. spathulata (?) (unc C)

*Potentilla anserinoides* (unc L)  
*Pratia angulata* agg.  
*Ranunculus acaulis* (c)  
*R. multiscapus* s.s. (unc S - south of  
 Mt Barton)  
*R. reflexus*  
*R. sp.* (*R. amphitrichus* or *R. glabrifolius*) (unc C)  
*Samolus repens* var. *repens* (c)  
*Sarcocornia quinqueflora* (c)  
*Scandia geniculata* (unc S - south of  
 Mt Barton)  
*Schizaelema trifoliolatum*  
*Scleranthus biflorus* (mainly south of  
 Mt Barton)  
*Selliera radicans* agg. (c)  
*Spergularia media* (c)  
*Stellaria decipiens* (incl. *S. parviflora*)  
*Urtica incisa*  
*U. linearifolia* (unc L)  
*Viola filicaulis*  
*Wahlenbergia coleusii*  
*W. gracilis* s.s. (c)  
*W. rupestris*

No. of species: 497.

Records not accepted

*Anaphalis trinervis* (Druce 1971, as *Gnaphalium trinerve*) [= *A. ketiensis*]  
*Astelia nervosa* (Druce 1971) [= *A. sp.* (unnamed)]  
*Chionochloa bromoides* (Cheeseman 1925 and Druce 1953, as *Danthonia*  
*bromoides*) [= *C. beddiei*]  
*Coprosma parviflora* (Wardle 1967) [= *C. sp.* (unnamed)]  
*Pterostylis venosa* (Wardle 1967) [= ?]  
*Lagenifera petiolata* (Druce 1971) [= *L. strangulata*]  
*Pteris macilentata* (Druce 1971) [= *P. penidula*]  
*Rytidosperma setifolium* (Druce 1953, as *Danthonia setifolia*) [= *R.*  
*patrosum*]  
*Plantago triandra* (Druce 1953) [= *P. masoniae*]  
*Parahebe catarractae* (Druce 1953) [= *P. diffusa*]  
*Dichondra brevifolia* (Druce 1953) [= *D. sp.* (unnamed)]

REFERENCES

- Cheeseman, T.F. 1925: "Manual of the New Zealand Flora", 2nd ed. (Ed. W.R.B. Oliver). Government Printer, Wellington. 1163 pp.
- Crookes, M. 1963: "New Zealand Ferns". Ed. 6. Whitcombe & Tombs. 408 pp.
- Druce, A.P. 1953: Plant Distribution records (1). Bull. Wellington bot. Soc. 26: 18-22.
- 1971: The flora of the Aorangi Range, southern Wairarapa, with notes on the vegetation. Bull. Wellington bot. Soc. 37: 4-30.
- Edgar, F. 1966: Luzula in New Zealand. N.Z. J. Bot. 4: 159-184.
- Garnock-Jones, P.J. 1986: A new status for the New Zealand mouse-tail (Myosurus, Ranunculaceae). N.Z. J. Bot. 24: 351-354.
- Hooker, J.D. 1864: "Handbook of the New Zealand Flora". London.
- Oliver, W.R.B. 1935: The genus Coprosma. Bishop Mus. Bull. 132. 207 pp.
- Pritchard, G.G. 1957: Experimental taxonomic studies on species of Cardamine Linn. in New Zealand. Trans. R. Soc. N.Z. 85: 75-89.
- Simpson, G. ; Thomson, J.S. 1943: Notes on some New Zealand plants and descriptions of new species. Trans. R. Soc. N.Z. 73: 155-171.
- Sykes, W.R. 1987: Pp. 153-4. In: Name changes in the indigenous New Zealand flora, 1960-1986. N.Z. J. Bot. 25: 115-170.
- Wardle, J. 1967: Vegetation of the Aorangi Range, southern Wairarapa. N.Z. J. Bot. 5: 22-48.
- Zotov, V.D. 1965: Grasses of the subantarctic islands of the New Zealand region. Rec. Dom. Mus. Wellington 5: 101-46.

