

HIGHER PLANTS (LYCOPODS, FERNS, GYMNOSPERMS, FLOWERING PLANTS) OF HAUPIRI RANGE AND BOULDER LAKE, N.W. NELSON


3200-5055 FT (970-1535 M)

A.P.Druce, Wellington, May 1985 (revised July 1985, June 1986,  
June 1988, July 1990,

Highest point: Slate River Peak, 5055 ft, grid ref. S8 031749  
July 1991,  
July 1992,  
May 1993)

(Based on visits Jan 1977 and Feb. 1985)

Rocks: shale, argillite, sandstone, conglomerate, quartzite; phyllite, schist; volcanics; alluvium (vicinity of Boulder L.)



## INDIGENOUS SPECIES

(unc) - uncommon or local

### GYMNOSPERM TREES AND SHRUBS

- Halocarpus biformis*
- Lepidothamnus intermedius*
- L. laxifolius*
- L. intermedius* × *L. laxifolius*
- Libocedrus bidwillii*
- Phyllocladus alpinus* agg.
  
- Podocarpus hallii* (unc)
- P. nivalis*
- P. hallii* × *P. nivalis* (unc)

### MONOCOT TREES

- Cordyline indivisa*

### DICOT TREES

- Dracophyllum traversii* (incl. *D. pyramidale*)
- Elaeocarpus hookerianus*
- Fuchsia excorticata* (unc)
- Griselinia littoralis*
- Metrosideros umbellata*
- Nothofagus fusca*
- N. menziesii*
- N. solandri* var. *eliffortioides*
- Pittosporum dallii* (unc)
- P. patulum* (unc)
- Pseudopanax crassifolius* (unc)
- P. linearis*
- P. simplex* (incl. *P. s. var. sinclairii*)
- P. sp.* (unnamed; included in *P. colensoi*, as var. *fiordensis* and as var. *ternatus*, by Wardle 1968)
- Wainmannia racemosa* var. *racemosa* (unc)

### DICOT SHRUBS

- Archeria traversii* (incl. *A. t. var. australis*)

*Aristotelia fruticosa* var.

- Brachyglottis adamsii*
- B. bidwillii* var. *vitidis*
- B. rotundifolia* var. (incl. *B. buchananii* and *B. cockaynei*) (unc)
- B. bidwillii* × *B. rotundifolia* (unc)
- Cassinia vauvilliersii*

*Coprosma cheesemanii* s.s.

*C. ciliata* ssp. *ciliata*

*C. colensoi* s.s.

*C. crenulata*

*C. depressa*

*C. foetidissima*

*C. perpusilla* ssp. *perpusilla*

*C. propinqua* ssp. *propinqua*

*C. pseudocuneata* ssp.

*C. hammoides* (incl. *C. polymorpha*)

*C. rugosa* ssp. *rugosa* (unc)

*C. setulata* (unc)

*C. sp. (i)* (unnamed; aff. *C. pseudocuneata*) (low-growing shrubs with small, thick, arcuate lvs)

*C. sp. (ii)* (unnamed; aff. *C. colensoi*) (low-growing shrubs with trailing branches) (unc)

*C. sp. (iii)* (unnamed) (*C. parviflora* var. *dumosa* sensu Allan 1961, non *C. p. var. dumosa* Cheeseman)

*C. propinqua* × *C. robusta* (unc)

*Cotinia sarmentosa*

*Cyathodes dealbata* (incl. *C. pumila*)

*C. impetifolia*  
*C. sp. (C. juniperina agg.)*  
*Dracophyllum longifolium* var.  
 (incl. *D. filifolium* *D. oliveri*)  
  
*D. palustre* (unc)  
*D. pubescens* var. *pubescens*  
*D. uniflorum* var. *uniflorum*  
*D. sp.* (unnamed; aff. *D. kirkii*) (low-growing shrubs; branches often rooting; infl of few-fld racemes)  
 (unc)  
*Epactis alpina*  
*Exocarpus bidwillii*  
*Gaultheria antipoda* (unc)  
*G. crassa*  
*G. depressa* s.s.  
*G. rupestris* (incl. *G. sulcorymbosa*)  
*G. sp.* (unnamed; included in *G. depressa*, as var. *novae-zelandiae*, by Franklin 1962)  
*Hebe albicans* var. *albicans*  
*H. canterburiana*  
*H. ciliolata*  
*H. cockayneana* var.  
*H. gracilima*  
*H. haastii* var. (aff. var. *humilis*;  
 see Moore 1961, p. 940) (unc)  
*H. macrantha* var. *brachyphylla*  
*H. salicifolia* var. *salicifolia*  
*H. tefrigona* var. (*H. coarctata*)  
*H. topiaria*  
*H. sp(i)* (*Leonohebe masoniae*)\*  
*H. sp(ii)* (*L. mooreae*)  
*H. salicifolia* X *H. sp.* (unc)  
*Helichrysum intermedium* var.  
*Leptospermum scoparium*

*Leucopogon fasciculatus* var.  
*L. fraseri* s.s.  
*L. suaveolans* (incl. *Cyathodes colensoi*)  
*Melicytus alpinus* s.s.  
  
*Myrsine divaricata* s.s.  
*M. nummularia*  
*Olearia atborescens*  
*O. avicenniifolia*  
*O. colensoi* var. *colensoi* (unc)  
*O. lacunosa* var. *lacunosa*  
*O. nummularifolia* var. (Ivs  
 distinctly longer than broad,  
 rather than suborbicular as  
 in var. *nummularifolia*) (unc)  
*Pentachondra pumila*  
*Peraxilla tetrapetala* (unc)  
*Pimelea suteti* var. (*P. orzophila*)  
*Pittosporum anomalum*  
*P. rigidum* var. (*P. cassinoides*)  
*Pseudopanax anomalous*  
*Pseudowintera traversii*  
*Traversia baccharoides* (unc)

#### DICOT LIANES AND RELATED

##### TRAILING PLANTS

*Muehlenbeckia axillaris*  
*Rubus cissoides* var. (leaflets  
 broader than in var. *cissoides*)  
*R. parvus* (unc)  
*R. schmidelioides* var. *schmidelioides*

#### LYCOPODS

*Lycopodium australianum*  
*L. fastigiatum*  
*L. latifoliale* (incl. *L. tamulosum*?)  
*L. scariosum*  
*L. vatum* (incl. *L. billardieri*  
 and *L. novae-zelandicum*) (unc)

\* (incl. *L. m. var. rotundata*)

## FERNS

- Asplenium flaccidum*  
*A. hookerianum* (unc)  
*Blechnum discolor*  
*B. fluviale* agg.  
*B. penna-marina*  
*B. procerum*  
*B. sp. (i)* (unnamed) (common lowland sp;  
 lower pinnae reduced in length)  
*B. sp. (ii)* (unnamed) (mountain sp;  
 lower pinnae not reduced in length  
 ("The Dunedin-Cass Variety" of Crozat)  
*Cyathea colensoi*  
*Cystopteris tasmanica*  
*Gleichenia dicarpa*  
*Grammitis billardierei*  
*G. magellanica* ssp. *nothofageti*  
*G. patagonica*  
*G. poeppigiana*  
*Hymenophyllum flabellatum* (unc)  
*H. multifidum*  
*H. latum* (unc)  
*H. flavescens* (unc)  
*H. villosum*  
*Hypolepis millefolium*  
*Ophioglossum coriaceum* agg.  
*Phymatosorus diversifolius* (unc)  
*Polystichum vestitum*  
*Pittosporum eleagnifolia* (unc)  
*Schizaea australis*  
*Sticherus cunninghamii* (unc)

## ORCHIDS

- Aporostylis bifolia*  
*Chiloglottis cornuta* (unc)  
*Corybas trilobus* ss.  
*Gastrodia cunninghamii* (unc)  
*Microtis oligantha*

*Prasophyllum colensoi*  
*Pterostylis venosa* (unc)  
*Thelymitra cyanea*  
*T. hatchii*

## GRASSES

- Agrostis dyeri* s.s.  
*A. muellerana*  
*A. sp. (i)* (unnamed; aff. *A. dyeri*)  
 (panicle spreading, not contracted  
 as in *A. dyeri*)  
*A. personata*  
*Chionochloa australis*  
 (1963) *C. cheesemanii*  
*C. conspicua* ssp. *conspicua*  
*C. pallens* ssp. *pallens*  
*C. rubra* ssp. *occulta*  
*C. sp.* (unnamed) (*C. flavescens*  
 auct. South Island, non *C.*  
*flavescens* Zotov)  
*C. cheesemanii* × *C. conspicua*  
 (unc)  
*C. conspicua* × *C. rubra* (unc)  
*Deschampsia novae-zelandiae*  
 (unc)  
*D. tenella*  
*Dexauxia aucklandica*

- Elymus hardyioides* (unc)  
*E. sp.* (aff. *E. rectisetus*) (unc)  
*Festuca* sp. (i) (cf. *F. mathewsii*)  
 (plant tufted, but not tussock-forming,  
 glaucous)  
*F. sp. (ii)* (unnamed; aff. *F. rubizii*)  
*Hierochloe novae-zelandiae*  
*Koeleria cheesemanii*  
*Lachnagrostis* sp. (unnamed;  
 included in *L. filiformis*, as  
 var. *semiglabra*, by Zotov 1965)  
*Mictolaelia avenacea*

*Poa breviglumis*  
*P. cita* ssp. *cita*  
*P. colensoi*  
*P. sp.(i)*(*P. novae-zelandiae* or *P. sub-  
vestita*)  
*P. sp.(ii)*(unnamed; aff. *P. novae-zelandiae*)

*Schoenus pauciflorus* s.s. ("slend-  
form" of Edgar 1970, p. 194)  
*Uncinia clavata*  
*U. divaricata*  
*U. egmontiana*  
*U. filiformis*  
*U. fuscovaginata* var. (plants glaucous)  
*U. nervosa* ss.  
*U. tubra*  
*U. rupestris* (incl. *U. angustifolia*)  
*U. silvestris* (incl. *U. affinis*)  
*U. zotovii*  
*U. uncinata*  
*U. sp.(unnamed)*(*U. affinis* sensu  
Hamlin 1959, non *U. riparia* var.  
*affinis* Col. ex Clarke; see Nootboom  
1978)

### SEDGES

*Carex acicularis*  
*C. dallii*  
*C. echinata* var. (*C. stellulata* var. *australis*)  
*C. flavaformis*  
*C. forsteri* (unc)  
*C. gaudichaudiana*  
*C. kirkii* (unc)  
*C. libera*  
*C. sinclairii*  
*C. sp.(unnamed; aff. *C. testacea*)*  
(Ivs upto 5mm wide, reddish;  
utricle margins not scabrid)

*Carex alpina* s.s.  
*Eleocharis acuta*  
*Gahnia procera*  
*Isolepis aucklandica*  
*I. habra*  
*I. subtilissima*  
*Oreobolus impar*  
*O. pectinatus*  
*O. strictus*

### RUSHES AND ALLIED PLANTS

*Centrolepis ciliata*  
*Empodium minus*  
*Juncus antarcticus*  
*J. gregiflorus* (unc)  
*J. novae-zelandiae*  
*Luzula colensoi*  
*L. migrata*  
*L. rufa* s.s.  
*L. subclavata*  
*L. sp.(i)*(unnamed; included in *L.*  
*chinata*, as var. *petrieana*, by  
Edgar 1966)  
*L. sp.(ii)*(unnamed; included in *L.*  
*picta*, as var. *limosa*, by Edgar  
1966)  
*Marsippospernum gracile*

### MONOCOT HERBS OTHER THAN ORCHIDS, GRASSES, SEDGES, RUSHES

*Astelia graminea*  
*A. lineatis* var. *linearis*

*A. nervosa* s.s. (unc)

*A. petriei*

*A. skottsbergii*

*A. sp.(i)* (unnamed; aff. *A. nervosa*)  
(lvs broad, not conspicuously  
white adaxially as in *A. nervosa*)

*A. sp.(ii)* (unnamed; included in *A.*  
*nivicola*, as var. *moticeae*, by  
Moore 1966)

*Libertia ixiooides* (unc)

*L. pulchella*

*Phormium cookianum*

*Potamogeton cheesemanii*

*P. suboblongus*

### COMPOSITE HERBS

*Abrotanella cassiniflora*

*Anaphalis* sp. (*Helichrysum bellidioides* ss.)

*Brachyslektis* sp. (unnamed; included  
in *B. bellidioides*, as var. *crassa*, by  
Nordenstam 1978)

*Brachyscome radicata*

*Celmisia bellidioides*

*C. dallii*

*C. hieracifolia* ssp.

*C. incana* (incl. *C. discolor*)

*C. laticifolia*

*C. lateralis* (incl. *C. l.* var. *villosa*)

*C. montei* var. (*C. coriacea* var. *lancifolia*)

*C. petriei*

*C. sessiliflora*

*C. similiis*

*C. sinclairii* (incl. *C. calliantha* & *C. dufratii*)

*C. spectabilis* ssp.

*C. traversii*

*C. sp.(i)* (unnamed; aff. *C. gracilenta*)  
(plants strongly rhizomatous)

*C. sp.(ii)* (unnamed; aff. *C. gracilenta*)

(lvs linear to linear-lanceolate;  
upper surface and margins  
thinly covered in tomentum;  
plants tufted;

lower surface glabrous, exc. for  
midrib)

*C. spectabilis* × *C. traversii* (Patterson  
and Patterson, 1956)

*Craspedia* sp.(i) (unnamed; includ-  
ed in *C. lanata*, as var. *elongata*,  
by Allan 1961) (lvs white or greenish  
white, both surfaces clad in cottony  
hairs; fls white)

*C. sp.(ii)* (unnamed) (summer-green;  
lvs green, both surfaces densely  
clad in short glandular hairs < 0.3  
mm long; margins + faintly white  
with cottony hairs, fls white)

*Gnaphalium avda*

*G. laterale*

*G. mackayi* s.s.

*G. paludosum* s.s.

*G. polylepis* (unc)

*G. ruhinicum*

*Helichrysum filicaule*

*Lagenifera cuneata*

*L. pumila*

*L. strangulata*

*Leptinella* sp.(i) (*L. pyrethrifolia* agg.)

*L. sp.(ii)* (unnamed; included in *L.*  
*squalida*, as ssp. *mediaea*, by  
Lloyd and Webb, 1987)

*Microrseris scapigeta* var.

*Pseudognaphalium* sp. (unnamed; aff.  
*P. luteoalbum*) (lvs mostly linear-  
spatulate to linear, acute)

*Raoulia eximia* var. (*R. rubra*)

*R. glabra*

*R. grandiflora* var. *grandiflora*

*R. tenuicaulis* (incl. *R.t.* var. *dimorpha*  
and *R.t.* var. *pusilla*)  
*Senecio wairauensis*  
*Taraxacum magellanicum*

*E.sp.(ii)*(unnamed; included in  
*E.brunnescens*, as ssp. *minuti-*  
*florum*, by Raven and Engel-  
horn, 1971)

**DICOT HERBS OTHER THAN COMPOSITES**

*Acaena ansetinifolia*

*A.inermis*

*A.profundèincisa*

*A.ansetinifolia* x *A.inermis*

*Aciphylla anomala*

*A.ferox*

*A.polita* var. (plants more robust)

*Anisotome aromatica*

*A.deltoidea*

*A.haastii*

*A.pilifera*

*Cardamine sp.(i)*(*C.debilis* agg.)

("Glossy Leaf" of Pritchard 1957)

*C.sp.(ii)*(unnamed) (lvs mostly simple;  
blades ± reniform)

*C.sp.(iii)*(undetermined)

*Cheesemania latisiliqua* (incl.  
*C.gibbsii*)

*Colobanthus canaliculatus*

*C.wallii* (incl. *C.masoniae*)

*Donatia novae-zelandiae*

*Drosera arcturi*

*D.späthulata*

*Epilobium brunnescens* s.s.

*E.chlorifolium*

*E.cockayneanum*

*E.glabellum* s.s.

*E.komarovianum*

*E.linnacoides*

*E.margaretiae*

*E.petnitens*

*E.pychostachyum*

*E.sp.(i)*(unnamed; aff. *E.glab-*  
*ellum*)

*Euphtasia townsonii* s.s.

*E.zelandica* agg.

*E.sp.(i)*(unnamed; aff. *E.laingii*)

*E.sp.(ii)*(unnamed; aff. *E.petriei*)

*Forstera mackayi*

*F.tanella*

*F.sp.*(unnamed)(lvs closely spaced,  
not recutted, persisting well down  
stem, becoming hair translucent; cost

*Galium perpusillum* s.s. not apparent

*Gentiana bellidifolia* var.

*G.gracilitolia* (incl. *G.vernicosa*)

*G.montana* (incl. *G.m.* var. *stolonifera*,

*G.patula*, *G.bellidifolia* var. *pulchella*)

*G.spaceri* (ung)

*G.sp.(i)*(unnamed; aff. *G.cotymbifera*)

*G.sp.(ii)*(unnamed; aff. *G.bellidifolia*)  
(plants forming small to large patch-  
es; stems decumbent; fls large,  
creamy white)

*Geranium sp.*(aff. *G.nudophyllum*)

*Geum leiospermum*

*G.sp.*(*G.parviflorum* ags.) (stock  
short, terminal leaflet about as  
long as broad; fls up to 13 mm  
diam.)

*Gingidia decipiens*

*Gonocarpus aggregatus*

*Gunnera dentata* var. *dentata*

*G.monoica*

*Hydrocotyle microphylla*

*H.sulcata*

*H.sp.*(unnamed; included in *H.*  
*novae-zelandiae* as var.  
*montana*, by Kirk 1899)

*Hypsela* sp.(unnamed)(minute  
herb; fls 3mm diam.; corolla  
lobes narrowly lanceolate)

*Jovellana repens*  
*Kelleria dieffenbachii(?)*  
*K. laxa*  
*K. multiflora*  
*K. tessellata*  
  
*Lobelia roughii*  
*Mentha cunninghamii*  
*Neopaxia* sp. (incl. *Montiacalycina* ?)  
*Claytonia austrolasica* var. *biflora*)  
*Myosotis macrantha*  
*M. sp.* (unnamed; included in *M. pygmaea*, as var. *drucei*, by Moore 1961) (unc)  
*Myriophyllum pedunculatum* ssp. n.3.  
*Nertera balfouriana*  
*N. ciliata*  
*N. depressa* (incl. *N. cunninghamii*)  
*N. sp.* (unnamed; aff. *N. dichondrifolia* (hts present on both surfaces of lf, straight, not hooked as in *N. dichondrifolia*))  
*Notothlaspi* sp. (unnamed) (*N. austrole* auct., non *Thlaspi australe* Hook.f.)  
*Oreomythris colensoi* s.s. (incl. *O.c.* var. *hispida* and *O.c.* var. *multifida*)  
*Oreostylidium subulatum*  
*Ovisia caespitosa* ssp. *caespitosa*  
*O. lactea* ssp. *lactea*  
*O. macrocarpa* ssp. *calycina*  
*O. Simpsonii*  
*Oxalis magellanica*  
*Parahedbe decora* (unc)  
*P. lyallii*  
*Phyllachne colensoi*  
*Plantago raoulii*  
  
*P. triandra* s.s.  
*P. uniflora*

*Pratia angulata* agg.  
*Ranunculus foliosus*  
  
*R. insignis*  
*R. membranifolius*  
*R. verticillatus*

*Schizolema haastii* ss.  
*S nitens*  
*Scleranthus brockiei* (unc)  
*Stellaria decipiens* (incl. *S. pativ-flora*)  
*Urtica incisa*  
*Viola cunninghamii*  
*V. filicaulis*  
*V. lyallii*  
*Wahlenbergia albomarginata*  
 var. *albomarginata* (incl. *W. pygmaea* var. *laxa*, *W. brockiei*, and *W. Simpsonii*)

#### ADVENTIVE SPECIES

#### GRASSES

*Agrostis capillaris*  
*Aira caryophyllea*  
*A. praecox*  
*Anthoxanthum odoratum*  
*Cynosurus cristatus* (unc - near hut)  
*Dactylis glomerata*

(continued on next page)

*Festuca nigrescens*  
*Holcus lanatus*  
*Poa annua*

**SEDGES**

*Carex ovalis*

**RUSHIES**

*Juncus articulatus*  
*J. effusus*  
*J. tenuis* (unc - neat hut)  
J. sp. (small - undetermined)  
*Luzula congesta*

**COMPOSITE HERBS**

*Cirsium vulgare*

*Crepis capillaris*  
*Hypochaeris radicata*  
*Leontodon taraxacoides*  
*Mycelis muralis*

*Taraxacum officinale* s.l. (unc)

**DICOT HERBS OTHER THAN COMPOSITES**

*Cerastium fontanum* ssp. *vulgare*  
*Linum catharticum*  
*Lotus pedunculatus* (unc)  
*Rumex acetosella*  
*Sagina procumbens*  
*Trifolium dubium*  
*T. repens*

No. of indigenous spp.: 367.  
No. of adventive spp.: 28.

Doubtful and erroneous records:

*Brachyscome cassinioides* (Petterson & Petterson, 1956, as *Senecio cassinioides*)  
"*Celmisia novae-zelandiae*" (Petterson & Petterson, 1956)  
*Forsteria bidwillii* (Petterson & Petterson, 1956)  
*Pseudopanax colensoi* (Petterson & Petterson, 1956, as *Nothopanax colensoi* var. *montanum*)  
*Schizilema roughii* (Petterson & Petterson, 1956)

REFERENCES

Allan, H.H. 1961: "Flora of New Zealand". Government Printer, Wellington. Vol. 1. 1085 pp.

Crookes, M. 1963 :  
and Tombs. 408 pp.

"New Zealand Flora". Ed. 6. Whitcombe

Edgar, E. 1966: Luzula in New Zealand. N.Z. Jl Bot. 4: 159-84.

— 1970: In "Flora of New Zealand". Government Printer, Wellington. Vol. 2. 354 pp.

Franklin, D.A. 1962: The Ericaceae in New Zealand (Gaultheria and Pernettya). Trans R. Soc. N.Z. Bot. 1: 155-73.

Hackel, E. 1903: On new species of grasses in New Zealand. Trans N.Z. Inst. 35: 377-85.

1> Kirk, T. 1899: "The Student's Flora of New Zealand". Government Printer, Wellington. 408 pp.

3> Moore, L.B. 1961: In "Flora of New Zealand". Government Printer, Wellington. Vol. 1. 1085 pp.

— 1966: Australasian asteliads (Liliaceae). N.Z. Jl Bot. 4: 1

2> 201-40.

Patterson, J.; Patterson, G.. 1956: In search of Pithosporum dallii. Bull. Wellington bot. Soc. 28: 14-23.

Pritchard, G.G. 1957: Experimental taxonomic studies on species of Cardamine in New Zealand. Trans R. Soc. N.Z. 85: 75-89.

Raven, P.H.; Engelhorn, T. 1971: New taxa and new combinations in Australasian Epilobium (Onagraceae). N.Z. Jl Bot. 9: 345-50.

Wardle, P. 1968: The taxonomy and distribution of the stipulate species of Pseudopanax in New Zealand. N.Z. Jl Bot. 6: 226-36.

Zotov, V.D. 1965: Grasses of the subantarctic islands of the New Zealand region. Rec. Dom. Mus. Wellington 5: 101-46.

— . —  
1> Hamlin, B.G. 1959: A revision of the genus Uuncinia (Cyperaceae - Caricoideae) in New Zealand. Dom. Mus. Bull. 19. 106 pp.

2> Nooteboom, H.P. 1978: A taxonomic revision of the Malayan and Australian species of Uuncinia (Cyperaceae). Blumea 24: 511-20.

Nordanstam, B. 1978. Opera Bot. 44: 1-84.

3> Lloyd, D.G.; Webb, C.J. 1987: The reinstatement of Leptinella at generic rank, and the status of the 'Cotuleae' (Asteraceae, Anthemideae).

N.Z. Jl Bot. 25: 99-105.

— . —