

(2/2)

HIGHER PLANTS (LYCOPODS, FERNS, GYMNOSPERMS, FLOWERING
PLANTS) OF MT PEEL* AND TABLELAND, NORTH-WEST
NELSON, EXCLUDING AREAS OF TERTIARY ROCKS†
900-1654 M (3000-5400 FT)

A.P.Druce, Wellington; and P.A.Williams, Botany Division,

D.S.I.R., Nelson; May 1987


(revised Sept. 1987, Oct. 1987, June 1988, June 1990, June 1991)

(Based on visits April 1969, Dec. 1986, May 1987, Dec. 1988)

* Grid ref. M27 757075

† For plants of these areas see List No. 285

July 1991



INDIGENOUS PLANTS

(unc) - uncommon or local

GYMNOSPERM TREES AND SHRUBS [DICOT SHRUBS]

- Halocarpus bidwillii*
- H. biformis*
- Lepidothamnus laxifolius*
- L. intermedius intermedius* X
 L. laxifolius (unc)
- Libocedrus bidwillii*
- Phyllocladus alpinus* agg.
- Podocarpus hallii*
- P. nivalis*
- P. hallii* X *P. nivalis* (unc)

MONOCOT TREES

- Cordyline indivisa* (unc)

DICOT TREES

- Carpodetus serratus*
- Dracophyllum traversii* (incl. *D. pyramidale*)
- Fuchsia excorticata* (unc)
- Griselinia littoralis*
- Hohertia lyallii* var. (*H. glabrata*)
- Nothofagus fusca*
- N. menziesii*
- N. solandri* var. *cliffortioides*
- Pseudopanax colensoi* s.s.
- P. linearis* (unc)
- P. simplex* (incl. *P. s.* var. *sinclairii*)
- P. sp.* (unnamed; included in *P. colensoi*, as var. *fiordensis* and as var. *ternatus*, by Wardle 1968)

- Aristotelia fruticosa* vat.
- Brachyglottis adamsii*
- B. bidwillii* var. *viridis*
- B. rotundifolia* vat. (incl. *B. buchananii* & *B. cockaynei*) (unc)
- Cassinia vauvilliersii*

- Coprosma atropurpurea*
- C. cheesemanii* ss.
- C. ciliata* ssp. *ciliata*
- C. colensoi* ss.
- C. depressa*
- C. foetidissima*
- C. microcarpa*

- C. petraea* ssp. *petraea*
- C. patricii* (Oliver 1935)
- C. pseudocuneata* ssp.
- C. rhomboides* (incl. *C. polymorpha*)
- C. setulata* (unc)
- C. sp. (i)* (unnamed; aff. *C. pseudocuneata*) (low-growing shrubs with small, thick, arcuate lvs)
- C. sp. (ii)* (unnamed; included in *C. parviflora* by Oliver 1935, and others) (lf undersurface not minutely hairy as is usual in *C. parviflora*; drupes vs. unpigmented, occ. purplish red)
- C. sp. (iii)* (unnamed) (*C. parviflora* var. *dumosa* sensu Allan 1961, not *C. p.* var. *dumosa* Cheeseman)
- C. cheesemanii* X *C. ciliata* (unc)

- Coriaria sarmentosa*
Cyathodes dealbata (incl. *C. pumila*)
C. ampetifolia
Dracophyllum longifolium var. (incl. *D. filifolium* and *D. olivaceum*)
D. pubescens var. *pubescens* (unc)
D. uniflorum
Exocarpus bidwillii
Gaultheria antipoda (unc)
G. crassa
 { *G. sp.* (unnamed; included in *G. depressa*, as var. *novae-zelandiae*, by Franklin 1962)
G. nubicola
Habea albicans var. *albicans*
H. canterbutiensis
H. ciliolata
H. cockayneana var.
H. opacidea var. *opacidea* (unc)
H. haastii var. (aff. var. *humilis*; see Moore 1961, p. 940)
H. macrantha var. *brachiphylla*
H. ochracea
H. odora s.s.
H. salicifolia var. *salicifolia*
H. tetragona var. (*H. coarctata*)
H. topiaria
H. vernicosa
H. sp. (*Leonotis* *masoniae*) *
Helichrysum intermedium
 var.
Leucopogon fraseri s.s.
L. suaveolens (incl. *Cyathodes colensoi*)
Malicytus alpinus s.s.
- Myrsine divaricata* s.s.
M. nummularia
Olearia capillaris (unc)
O. colensoi var. *colensoi* (unc)
O. lacunosa var. *lacunosa*
O. nummulariifolia var. (lvs distinctly longer than broad, rather than suborbicular as in var. *nummulariifolia*)
O. virgata var. *implicata* (incl. *O. v.* var. *divaricata* and *O. v.* var. *laxiflora*) (unc)
O. arborescens × *O. capillaris* (unc)
O. capillaris × *O. lacunosa* (unc)
Pentachondra pumila
- Pimelea suteri* var. (*P. oreophila*)
Pittosporum anomalum
P. rigidum var. (*P. crassicaule*)
Pseudopanax anomalous
Pseudowintera colorata
Traversia baccharoides

DICOT LIANES AND RELATED

TRAILING PLANTS

- Muehlenbeckia axillaris*
M. australis × *Maxillaris* (unc)
Rubus cissoides var. (leaflets broader than in var. *cissoides*)

LYCOPODS

- Lycopodium australe*
L. fastigiatum
L. scariosum
L. vatum (incl. *L. billardierei* and *L. novae-zelandicum*)

* (incl. *L. m.* var. *rotundata*)

FERNS

- Asplenium flaccidum*
A. richardii
Blechnum fluviatile agg.
B. penna-martini
B. proctatum
B. sp. (I) (unnamed) (common lowland sp.; lower pinnae reduced in length)
B. sp. (ii) (unnamed) (mountain sp.; lower pinnae not reduced in length) ("The Dunedin-Cass Variety of Crookes 1963")
Cyathea colensoi
Cystopteris tasmanica
Grammitis billardierei
G. givenii (unc)
G. magellanica ssp. *nothofageti*
G. patagonica
G. poeppigiana
Histiopteris incisa
Hymenophyllum multifidum
H. ratum
H. villosum
Hypolepis millefolium
Paesia scaberula
Polystichum vestitum
Pteridium esculentum (unc)

- Prasophyllum colensoi*
Pterostylis australis
P. humiliis (unc)
Thelymitra cyanea

GRASSES

- Agrostis dycti* ss.
A. muelleriana
A. personata
A. sp. (unnamed; aff. *A. dycti*)
 (panicle spreading, not contracted as in *A. dycti*)
Chionochloa australis
C. conspicua ssp. *conspicua*
C. pallens ssp. *pallens*
C. rubra ssp. *occulta*
C. sp. (unnamed) (*C. flavesans* avct. South Island, non *C. flavesans* Zotov)
C. rubra X *C. sp.* (unnamed) (unc)
Deschampsia tenella
Deyeuxia aucklandica
D. avenoides (incl. *D. a. var. brachyantha*)
Elymus sp. (aff. *E. rectisetus*)
 (lvs ± glaucous)
Festuca sp. (aff. *F. matthewsii*)
 (tufted, but not tussock-forming, glaucous)
Hierochloe novae-zelandiae
Koeleria cheesemanii
K. sp. (*K. novo-zelandica* agg)
 (plants slender, rhizomatous; lvs up to 1.5mm wide) (unc)

ORCHIDS

- Apotostylis bifolia*
Caladenia lyallii
Chiloglottis cornuta
Corybas macranthus
C. trilobus s.s.

<i>Lachnagrostis</i> sp. (unnamed; included in <i>L. filiformis</i> , as var. <i>semi-glabra</i> , by Zotov 1965)	<i>Oreobolus pectinatus</i>
<i>Microlaena colensoi</i> s.s.	<i>O. strictus</i>
<i>Poa breviglumis</i>	<i>Schoenus pauciflorus</i> ss. ("slender form" of Edgar 1970, p. 194)
<i>P. celsa</i>	<i>Uncinia caespitosa</i>
<i>P. colensoi</i>	<i>U. clavata</i>
<i>P. dipsacea</i>	<i>U. divaricata</i>
<i>P. imbecilla</i>	<i>U. filiformis</i>
<i>P. kirkii</i>	<i>U. fuscovagina tavar. (plants glaucous)</i>
<i>P. maia</i>	<i>U. involuta</i>
<i>P. subvestita</i>	<i>U. netvosa</i> s.s.
<i>P. sp. (unnamed; aff. <i>P. novae-zelandiae</i>)</i>	<i>U. rupestris</i> (incl. <i>U. angustifolia</i>)
<i>Rytidosperma buchananii</i>	<i>U. silvestris</i> (incl. <i>U. affinis</i>)
<i>R. gracile</i>	<i>U. sp. (unnamed) (U. affinis sensu Hamlin 1959, non <i>U. riparia</i> var. <i>affinis</i> Col. ex Clarke; see Nootboom 1978)</i>
<i>R. setifolium</i>	
<i>Simplicia buchananii</i> (unc)	
<i>Trisetum youngii</i>	
<i>T. sp. (unnamed) (included in <i>T. antarcticum</i>, as ssp. <i>tenellum</i>, by Petrie 1912)</i>	

SEDGES

RUSHES AND ALLIED PLANTS	
<i>Carex aciculatis</i>	<i>Centrolepis ciliata</i>
<i>C. echinata</i> var. (<i>C. stellulata</i> var. <i>australis</i>)	<i>Empodisma minus</i>
<i>C. flavaformis</i> (unc)	<i>Juncus antarcticus</i>
<i>C. gaudichaudiana</i>	<i>J. novae-zelandiae</i>
<i>C. libera</i>	<i>J. pusillus</i> (unc)
<i>C. pyrenaica</i> var. <i>cephalotes</i>	<i>Luzula colensoi</i>
<i>C. sinclairii</i>	<i>L. migrata</i>
<i>C. sp. (unnamed; aff. <i>C. coriacea</i>)</i> (lvs slender, bright green, not becoming stiff and yellow-green in exposed sites)	<i>L. picta</i> ss.
<i>Catpha alpina</i> s.s.	<i>L. rufa</i> ss.
<i>Eleocharis acuta</i>	<i>L. sp.(i) (unnamed; included in <i>L. picta</i>, as var. <i>limosa</i>, by Edgar 1966)</i>
<i>Gahnia procera</i>	<i>L. sp.(ii) (unnamed; included in <i>L. crinita</i>, as var. <i>petricana</i>, by Edgar 1966)</i>
<i>Isolepis aucklandica</i>	<i>L. colensoi</i> X <i>L. migrata</i>
<i>I. habra</i>	<i>L. migrata</i> X <i>L. rufa</i>

MONOCOT HERBS OTHER THAN
ORCHIDS, GRASSES, SEDGES, RUSHES
AND ALLIED PLANTS

<i>Arthropodium candidum</i>	<i>Calminia alpina</i> ss.
<i>Astelia graminea</i>	<i>C. bellidioides</i>
<i>A. linearis</i> var. <i>linearis</i>	<i>C. dallii</i>
<i>A. nervosa</i> ss.	<i>C. hieracifolia</i> ssp.
<i>A. nivicola</i> ss.	<i>C. incana</i> (incl. <i>C. discolor</i>)
<i>A. petriei</i>	<i>C. laeticifolia</i>
<i>A. sp.(i)</i> (unnamed; included in <i>A. nivicola</i> , as var. <i>motteae</i> , by Moore 1966)	<i>C. lateralis</i> (incl. <i>C. l.</i> var. <i>villosus</i>)
<i>A. sp.(ii)</i> (unnamed; aff. <i>A. nervosa</i>) (lvs broad, not conspicuously white adaxially as in <i>A. ner- vosa</i>)	<i>C. montoi</i> ssp. (<i>C. cotacea</i> var. <i>lancifolia</i>)
<i>Bulbinella hookeri</i>	<i>C. rupestris</i>
<i>Herpolitha novae-zelandiae</i>	<i>C. sessiliflora</i>
<i>Libertia peregrinans</i>	<i>C. similis</i>
<i>L. pulchella</i>	<i>C. sinclairii</i> (incl. <i>Callanii</i> & <i>C. dumetzi</i>)
<i>Phormium cookianum</i>	<i>C. spectabilis</i> ssp.
<i>Potamogeton cheesemanii</i> (unc)	<i>C. traversii</i>
<i>P. suboblongos</i> (unc)	<i>C. sp.(i)</i> (aff. <i>C. gracilenta</i>) (plants strongly rhizomatous)
	<i>C. sp.(ii)</i> (unnamed; aff. <i>C. graci- lenta</i>) (plants tufted; lvs linear to linear-lanceolate, up to 4.5 x 1.2 cm; upper surface ± margins thinly covered in tomentum; lower surface densely covered)

COMPOSITE HERBS

<i>Abrotanella caespitosa</i>	<i>C. sessiliflora</i> × <i>C. spectabilis</i> (unc)
<i>A. linearis</i> ss.	<i>C. s.</i> × <i>C. traversii</i> (unc)
<i>Anaphalis</i> sp. (<i>Helichrysum bellidi- oides</i> ss.)	<i>C. aspera</i> sp.(i) (unnamed) (plants summet-green; lvs green, both surfaces densely clad in short glandular hairs; margins ± white; fls white)
<i>Brachyglottis</i> sp. (unnamed; included in <i>B. bellidioides</i> , as var. <i>crassa</i> , by Nordenstam 1978)	<i>C. sp.(ii)</i> (unnamed)
<i>Brachyscome radicata</i>	<i>C. sp.(iii)</i> (unnamed)
<i>B. sinclairii</i> ss.	

- Anisotome aromatic
- | | |
|--|---|
| <i>Leptinella</i> sp.(i) (<i>L.pyrathrifolia</i> agg.)
<i>L.</i> sp.(ii) (unnamed; included in <i>L.</i>
<i>squalida</i> as ssp. <i>mediana</i> ,
by Lloyd and Webb, 1987)
<i>Gnaphalium mackayii</i> s.s.
<i>G. paludosum</i>
<i>G. traversii</i>
<i>Haastia sinclairii</i> s.s.
<i>Helichrysum filicaule</i>

<i>Lagenifera pinnatifida</i>
<i>L.pumila</i>
<i>L.strangulata</i>
<i>Leucogenes grandiceps</i>
<i>Microrhiza scapigera</i> var.
<i>Pseudognaphalium</i> sp. (unnamed;
aff. <i>P. luteoalbum</i>) (mountain sp.; lvs
mostly oblong-spathulate to
narrowly oblong, rounded)
<i>Raoulia eximia</i> var. (<i>Rubra</i>)
<i>R.glabra</i>
<i>R.grandiflora</i> var. <i>grandiflora</i>
<i>Senecio glaucophyllum</i> ssp. <i>glaucophyllum</i> (unc)
<i>S.wairauensis</i> | <i>A. deltoidea</i>
<i>A.filifolia</i>
<i>A.pilifera</i>
<i>A.</i> sp. (unnamed; included in <i>A. imbricata</i> , as var. <i>prostrata</i> , by Dawson
1961)

<i>Cardamine</i> sp.(i) (<i>C.debilis</i> agg.)
("Narrow Petal" of Pritchard 1957)
<i>C.</i> sp.(ii) (<i>C.debilis</i> agg.)
(cf. "Long Style" of Pritchard 1957)
<i>C.</i> sp.(iii) (unnamed) (lvs mostly simple
blade + reniform)
<i>C.</i> sp.(iv) (aff. <i>C. corymbosa</i>)
<i>Cheesemania latisiliqua</i> (incl.
<i>C.gibbsii</i>)
<i>Colobanthus aciculatus</i>
<i>C.apetalus</i> (incl. <i>C.a.</i> var. <i>alpinus</i>)
<i>C.canaliculatus</i>
<i>C.strictus</i>
<i>Crassula sinclairii</i>
<i>Donatia novae-zelandiae</i>
<i>Drosera arcturi</i>
<i>D.spathulata</i>
<i>D.stenopetala</i>
<i>Epilobium atriplicifolium</i> s.s.
<i>E.brunnescens</i> s.s.
<i>E.chlorifolium</i> |
|--|---|

[DICOT HERBS OTHER THAN COMPOSITES]

- Acaena fissistipula*
A.profundeincisa
A.saccaticupula
Aciphylla anomala
A.ferox
A.polita var.
A. sp. (unnamed; aff. *A.glaucocaeus*) (pinnae rigid)
A.anomala X *A. ferox* (Dawson
 and Le Comte, 1978)

- E.cochayneanum*
E.glabellum s.s.
E.gracilipes (unc)
E.macropus
E.margaretae
E.netteroides
E.pernitens
E.pubens

- E. pycnostachyum*
E. rubromarginatum
E. vernicosum
E. sp.(i)(unnamed; included in *E. brunneescens*, as ssp. *minutiflorum*, by Raven and Engelhorn, 1971)
E. sp.(ii)(unnamed; aff. *E. glabellum*)
Euphrasia cheesemanii
E. townsonii s.s.
E. sp.(i)(unnamed; aff. *E. laingii*)
E. sp.(ii)(unnamed; aff. *E. townsonii*)
Forsteria mackayi
Galium perpusillum s.s.
Gentiana bellidifolia var.
G. sp.(i)(unnamed; aff. *G. corymbifera*)
G. gracilifolia (incl. *G. vernicosa*)
G. montana (incl. *G. m.* var. *stolonifera*, *G. patula*, and *G. bellidifolia* var. *pulchella*)
G. sp.(ii)(unnamed; aff. *G. bellidifolia*) (plants forming small to large patches; lvs very glossy; fls large, creamy white)
Geranium sp. (aff. *G. microphyllum*)
Geum leiospermum
G. sp.(i) (*G. parviflorum* agg.) (stock short; terminal leaflet about as long as broad; fls up to 13 mm diam.)
G. sp.(ii) (*G. parviflorum* agg.) (stock creeping and branching; terminal leaflet slightly longer than broad; fls up to 8 mm diam.)
Gingidia decipiens
Gonocarpus aggregatus
G. micranthus ssp. *micranthus*
G. montanus
Hydrocotyle heteromeria (unc)
H. moschata (unc)
H. sulcata
Kelleria dieffenbachii
K. laxa
K. multiflora
K. tessellata
Lobelia roughii
Mentha cunninghamii
Montia fontana s.s.

Myosotis macrantha
M. sp.(i)(unnamed; included in *M. pygmaea*, as var. *drucei*, by Moore 1961)
M. sp.(ii) (*M. australis* agg.) (fls yellow)
M. sp.(iii) (*M. australis* agg.) (fls small, white)
Myriophyllum triphyllum (unc)
Neopaxia sp. (incl. *Montia calycina*)
Claytonia australasica var. *biflora*
Nertera balfouriana
N. ciliata
N. sp.(unnamed; aff. *N. dichondrifolia*) (hairs present on both surfaces of lf, straight, not hooked as in *N. dichondrifolia*)
Nototlaspi sp. (*N. austriale* auct., non *Thlaspi austriale* Hook.f.)
Oreomyrtis colensoi s.s. (incl. *O. c.* var. *hispida* and *O. c.* var. *multifida*)
Oreostylidium subulatum

Ouisia caespitosa ssp. *caespitosa* *Viola cunninghamii*
V. filicaulis
O. lactea ssp. *lactea* *Wahlenbergia albomarginata*
O. simpsonii var. *albomarginata* (incl. *W. brockiei*, *W. pygmaea* var. *laxa*,
Oxalis magellanica and *W. simpsonii*)
Parahelle cheesemanii
P. decora
P. lyallii No. of species: 363.
Phyllachne colensoi var.
Plantago lanigera s.l. (incl. *P. n.z.*)
P. raoulii
P. uniflora
Pratia angulata agg.
P. macrodon s.s.
Psychrophila novae-zealandiae
Ranunculus foliosus

R. insignis
R. membranifolius
R. mitus
R. reflexus
R. verticillatus

Schizolema colensoi
S. haastii s.s.
S. nitens
S. roughii
Scleranthus uniflorus
Stellaria decipiens (incl. *S. parviflora*)
S. gracilenta

ADVENTIVE PLANTS

GRASSES

Agrostis capillaris
Anthoxanthum odoratus
Dactylis glomerata (unc)
Festuca nigrescens
Holcus lanatus
Poa annua
P. pratensis

SEDGES

Carex ovalis

RUSHES

Juncus articulatus
J. effusus

COMPOSITE HERBS

Hypochoeris radicata
Mycelis muralis
Taraxacum officinale s.l.

DICOT HERBS OTHER THAN COMPOSITES

Cerastium fontanum ssp. *vulgare*
Linum catharticum
Rumex acetosella
Sagina procumbens
Trifolium repens

No. of species: 18.

REFERENCES

Allan, H.H. 1961: "Flora of New Zealand." Government Printer, Wellington. Vol. 1. 1085 pp.

Cheeseman, T.F. 1906: "Manual of the New Zealand Flora." Government Printer, Wellington. 1199 pp.

Dawson, J.W.; Le Comte, J.R. 1978: Research on Acyphella - a progress report. Tuatara 23: 49-67.

Crookes, M. 1963: "New Zealand Ferns". Ed. 6. Whitcombe & Tombs. 408 pp.

Dawson, J.W. 1961: A revision of the genus Anisotome (Umbelliferae). Univ. Calif. Publics Bot. 33: 1-98.

Edgar, E. 1966: huzula in New Zealand. N.Z. Jl Bot. 4: 159-84.

— 1970: In "Flora of New Zealand." Government Printer, Wellington. Vol. 2. 1085 pp.

Franklin, D.A. 1962: The Ericaceae in New Zealand (Gaultheria and Pezouetya) Trans. R. Soc. N.Z. Bot. 1: 155-73.

Hamlin, B.G. 1959: A revision of the genus Uncinia (Cyperaceae - Caicorideae) in New Zealand. Dom. Mus. Bull. 19. 106 pp.

Lloyd, D.G.; Webb, C.J. 1987: The reinstatement of heptinella at generic rank, & the status of the 'Cotuleae' (Asteraceae, Anthemideae). N.Z. Jl Bot. 25: 99-105.

Moor, L.B. 1961: In "Flora of N.Z." Govt Printer, Wgtn. Vol. 1. 1085 pp.
— 1966: Australasian asteliads (Liliaceae). N.Z. Jl Bot. 4: 201-40.

Nordenstam, B. 1978: Opera Bot. 44: 1-84.

Nooteboom, H.P. 1978: A taxonomic revision of the Malesian and Australian species of Uncinia (Cyperaceae). Blumea 24: 511-20.

Oliver, W.R.B. 1935: The genus Coprosma. Bishop Mus. Bull. 132. 207 pp.

Petrie, D. 1912: Descriptions of new native species of phanerogams. Trans. N.Z. Inst. 44: 179-87.

Pritchard, G.G. 1957: Experimental taxonomic studies on species of Cardamine Linn. in New Zealand. Trans Roy. Soc. N.Z. 85: 75-89.

Raven, P.H.; Englehorn, T. 1971: New taxa and new combinations in Australasian Epilobium (Onagraceae). N.Z. Jl Bot. 9: 345-50.

Wardle, P. 1968: The taxonomy and distribution of the stipulate species of Pseudopanax in New Zealand. N.Z. Jl Bot. 6: 226-36.

Itoev, V.D. 1965: Grasses of the subantarctic islands of the NZ region. Rec. Dom. Mus. Wellington 5: 101-46.

