

INDIGENOUS HIGHER PLANTS (LYCOPODS, FERNS, GYMNOSPERM  
FLOWERING PLANTS) OF TERTIARY CALCAROUS ROCKS<sup>†</sup>, NEAR  
TABLELAND\* AND UPPER TAKAKA VALLEY, NORTH-WEST  
NELSON, 760-1220 M / 2500-4000 FT

*slg*

A.P. Druce, Wellington, May 1987  
(revised Oct. 1987)



Keep

(Based on visits Jan. 1975, April 1986, & May 1987)

\* Grid ref. S13 002469

+ For plants of non-calcareous rocks see

List No. 272



INDIGENOUS HIGHER PLANTS (LYCOPODS, FERNS, GYMNOSPERMS  
FLOWERING PLANTS) OF TERTIARY CALCAROUS ROCKS<sup>†</sup>, NEAR  
TABLELAND\* AND IN UPPER TAKAKA VALLEY, NORTH-WEST  
NELSON, 750-1220 M / 2500-4000 FT

A.P. Druce, Wellington, May 1987


(Revised Oct. 1987, June 1990, June 1991, August 1992)

(Based on visits Jan. 1975, April 1986, May 1987,  
April 1989, Dec. 1991)

\* Grid Ref. NZMS 260 M27 785023

+ For plants of non-calcareous rocks see

List No. 272



(unc) - uncommon or local

**GYMNOSPERM TREES AND SHRUBS**

- Halocarpus bidwillii* (unc)  
*H. biformis*  
*Lagarostrobos colensoi* (unc)  
*Libocedrus bidwillii*  
*Phyllocladus* sp. (unnamed; aff. *P. alpinus*) ("lowland" ecotype of Wardle 1969)  
*Podocarpus hallii* (unc)  
*P. nivalis* (unc)

**DICOT TREES**

- Aristotelia serrata*  
*Carpodetus serratus*  
*Dracophyllum traversii* (incl. *D. pyramidale*)  
*Elaeocarpus hookerianus*  
*Fuchsia excorticata*  
*Griselinia littoralis*  
*Hohoria lyallii* var. (*H. glabrata*)  
*Metrosideros umbellata*  
*Nothofagus fusca*  
*N. menziesii*  
*N. solandri* var. *cliffortioides*  
*Pittosporum tenuifolium*  
*Pseudopanax crassifolius*  
*P. linearis* (unc)  
*P. macintyreai*  
*P. simplex* (incl. *P. s. var. sinclairii*)  
*P. sp.* (unnamed; included in *P. colensoi*, as var. *fiordensis* and as var. *ternatus*, by Wardle 1968)  
*Schefflera digitata* (unc)  
*Waimeaia racemosa* var. *racemosa*

**DICOT SHRUBS**

- Archetia traversii* (incl. *A. t. var. australis*)  
*Aristotelia fruticosa* var. *A. f. X A. serrata* (unc)  
*Brachyglottis bidwillii* var. *viridis* (unc)  
*B. hectorii* (unc)  
*B. rotundifolia* var. (incl. *B. buchananii* and *B. cochaynai*) (unc)  
*Coprosma cheesemanii* ss.  
*C. ciliata* ssp. *ciliata*  
*C. colensoi* ss.  
*C. depressa*  
*C. foetidissima*  
*C. linariifolia*  
*C. lucida* ss. (unc)  
*C. propinqua* ssp. *propinqua*  
*C. pseudocuneata* ssp.  
*C. thammoides* (incl. *C. polymorpha*) (unc)  
*C. rugosa* ssp. *rugosa*  
*C. sp(i)* (unnamed; aff. *C. colensoi*, (low-growing shrubs with trailing branches) (unc)  
*C. sp.(ii)* (unnamed; included in *C. parviflora* by Oliver, 1935, and others) (lf undersurface not minutely hairy as is usual in *C. parviflora*; drupes vs. unpigmented, occ. purplish red) (unc)

<i>C. sp(iii)</i> (unnamed) ( <i>C. parviflora</i> var. <i>dumosa</i> sensu Allan 1961, non <i>C. p.</i> var. <i>dumosa</i> Cheeseman)	<i>Myrsine divaricata</i> s.s.
<i>C. propinquia</i> × <i>C. robusta</i> (unc)	<i>Neomyrtus pedunculata</i>
<i>Coraria sarmentosa</i>	<i>Olearia arborescens</i>
<i>Cyathodes ampetifolia</i>	<i>O. avicinifolia</i>
<i>C. sp.</i> ( <i>C. juniperina</i> agg.) (lvs vs. $L \text{ cm} \times L \text{ mm}$ )	<i>O. lacunosa</i> var. <i>lacunosa</i>
<i>Dracophyllum longifolium</i> var. (incl. <i>D. filifolium</i> and <i>D. oliveri</i> )	<i>O. nummulariifolia</i> var. (lvs distinctly longer than broad, rather than suborbicular as in var. <i>nummulariifolia</i> ) (unc)
<i>D. uniflorum</i>	<i>O. virgata</i> var. <i>implicata</i> (incl. <i>C. divaricata</i> and <i>O. laxiflora</i> )
<i>Gaultheria antipoda</i>	<i>O. capillaris</i> × <i>O. lacunosa</i> (unc)
<i>G. sp.</i> (unnamed; included in <i>G. depressa</i> , as var. <i>novae-zelandiae</i> , by Franklin 1962)	<i>Pentachondria pumila</i>
	<i>Pimelea suteri</i> var. ( <i>P. oreophila</i> )

<i>Habe canterburiana</i>	<i>Pittosporum rigidum</i> var. ( <i>P. crassicaule</i> )
<i>H. gracillima</i>	<i>Pseudopanax anomalus</i>
<i>H. odora</i> s.s.	<i>Pseudowintera colorata</i>
<i>H. tetragona</i> var. ( <i>H. coactata</i> )	<i>Traversia baccharoides</i>
<i>H. topialia</i>	<i>Urtica ferox</i> (unc)
<i>H. vernicosa</i>	
<i>H. sp.(i)</i> ( <i>heonoheba masoniae</i> *)	
<i>H. sp.(ii)</i> (unnamed) (lvs up to 4x1 cm, elliptic-oblong, bright green)	<b>DICOT LIANES AND RELATED/</b> <b>TRAILING PLANTS</b>

<i>H. sp.(iii)</i> (aff. <i>H. glaucophylla</i> )	<i>Muehlenbeckia australis</i>
→ <i>Helichrysum aggregatum</i> var. aggr.	<i>M. axillaris</i>
<i>H. intermedium</i> var. (unc)	<i>M. australis</i> × <i>M. axillaris</i> (unc)
<i>Leptospermum scoparium</i>	<i>Rubus cissoides</i> var. (leaflets broader than in var. <i>cissoides</i> )
<i>Leucopogon fasciculatus</i> var.	<i>R. schmidelioides</i> var. <i>schmidelioides</i>
<i>L. fraseri</i> s.s.	
<i>Molycytus alpinus</i> s.s.	
<i>M. obovatus</i> var. <i>obovatus</i> (unc)	
<i>H. sp(iv)</i> (undetermined) (unc)	* (incl. <i>L. m.</i> var. <i>rotundata</i> )

**LYCOPODS**

*Lycopodium australianum*  
*L. fastigiatum*  
*L. scandens* (unc)

**FERNS**

*Adiantum cunninghamii*  
*Asplenium bulbiferum* (unc)  
*A. flaccidum*  
*A. gracilimum*  
*A. hookerianum*  
*A. lyallii*  
*A. oblongifolium*  
*A. sp. (unnamed; aff. A. trichomanes)*  
*Blechnum chambersii*  
*B. discolor*  
*B. fluviatile agg.*  
*B. membranaceum*  
*B. penna-matina*  
*B. procerum*  
*B. vulcanicum*  
*B. sp. (unnamed) (common low-  
land sp.) (lower pinnae reduced  
in length)*  
*Cyathea colensoi*  
*Ctenopteris heterophylla*  
*Cystopteris tasmanica*  
*Grammitis billardierei*  
*G. magellanica* ssp. *nothofagi*  
*G. patagonica*  
*Histiopteris incisa*  
*Hymenophyllum demissum*  
 (unc)

*H. multifidum*

*H. rotundum* (unc)  
*H. villosum*

*Hypolepis millefolium*

*Leptopteris superba* (unc)

*Paesia scaberula* (unc)

*Phymatosorus postulatus*

*Pneumatopteris pennigera*

*Polystichum richardii* (unc)

*P. vestitum*

*Pyrosia eleagnifolia*

**ORCHIDS**

*Aporostylis bifolia*  
*Cotybas macranthus*  
*C. trilobus* ss.  
*Pterostylis oliveti* (unc)

**GRASSES**

*Chionochloa australis*  
*C. cheesemanii* (unc)  
*C. conspicua* ssp. *conspicua*  
*C. pallens* ssp. *pallens*  
*C. rubra* ssp. *occulta*  
*Deschampsia tenella*  
*Elymus* sp. (aff. *E. rectisetus*)  
 (vs glaucous)  
*Festuca* sp. (unnamed) (unc)  
*Lachnagrostis* sp. (unnamed;  
 included in *L. filiformis*, as var.  
*semiglabra*, by Zotov 1965)

*Mictolaena avenacea*  
*M. colensoi*  
*Poa breviglumi*s  
*P. cita* ssp. *cita*(unc)  
*P. colensoi*  
*P. imbecilla*  
*P. pusilla*  
*Rytidosperma buchananii*  
*R. gracile*  
*R. Setifolium*

*Simplicia buchananii*(unc)  
*Trisetum* sp.(i)(unnamed)(plants slender; lf sheath glabrous or nearly so; lf blade flat, margins smooth, us. with a few hairs near ligule)  
*T.* sp.(ii)(unnamed)(plants robust; lf sheath densely hairy in lower part; lf blade flat, margins scabrid, us. with some long hairs near ligule)

*C. sp.(ii)*(unnamed)(ls channelled, green to brownish green, us. recurved; spikes 3-5, approximate in small plants, slightly spaced in larger ones; stigmas 3)  
*C. sp.(iii)*(Sect. *Acutae*)

*Eleocharis gracilis*(unc)  
*Gahnia procera*  
*Isolepis aucklandica*  
*I. habra*  
*I. subtilissima*(unc)  
*Oreobolus strictus*  
*Schoenus pauciflorus* S.S.  
*Uncinia astonii*  
*U. caespitosa*  
*U. clavata*  
*U. divaricata*  
*U. filiformis*  
*U. fuscovaginata* var. (*glauca*)

### SEDGES

*Carex breviculmis*  
*C. comans*  
*C. dissita*  
*C. flavidiformis*  
*C. forsteri*  
*C. gaudichaudiana*  
*C. libera*  
*C. sinclairii*

*C. sp.(i)*(unnamed; aff. *C. testacea*)(ls upto 5mm wide, reddish, utricle margins not scabrid)

*U. rubra*  
*U. rupestris* (incl. *U. angustifolia*)  
*U. uncinata*  
*U. zotovii*  
*U. sp.*(undetermined)

**RUSHES AND ALLIED PLANTS**  
*Empodisma minus*  
*Juncus antarcticus*  
*J. gregiflorus*  
*J. novae-zelandiae*  
*J. pusillus*(unc)

*Luzula migiata*  
L.sp. (*L. picta* agg.)

*Brachyscome radicata*

MONOCOT HERBS OTHER THAN  
ORCHIDS, GRASSES, SEDGES,  
RUSHES AND ALLIED PLANTS

*Arthropodium candidum*

*Astelia nervosa* s.s.

*A. nivicola* s.s.

*A. petriei*

*A. sp.(i)* (unnamed; included in

*A. nivicola*, as var. *moreae*, by Moore 1966)

*A. sp.(ii)* (unnamed; aff. *A. nervosa*)  
(lvs broad, not conspicuously white adaxially as in *A. nervosa*)

*Bulbinella hookeri*

*Libertia ixioides* (unc)

*L. peregrinans*

*L. pulchella*

*Luzuriaga parviflora*

*Phormium cookianum*

*Potamogeton* sp. (unc)

COMPOSITE HERBS

*Abretanella caespitosa*

*A. lineatis* s.s.

*Anaphalis* sp. (*Helichrysum bellidioides* s.s.)

*Brachyglottis* sp. (unnamed; included in *B. bellidioides*, as var. *crassa*, by Nordenstam 1978)

*Calmisia dallii*

*C. incana* (incl. *C. discolor*)

*C. monroi* ssp. (*C. semicordata* ssp. *semicordata*)

*C. sp.* (unnamed; aff. *C. glacialis* (plants tufted; lvs linear to linear-lanceolate, up to 45 x 1.2 cm; lower surface white with cottony hairs)) (unc)

*Craspedia minor* var. *minor* (incl. *C. major*)

*C. sp. (i)* (unnamed)

*C. sp. (ii)* (unnamed)

*Gnaphalium avda*x

*G. delicatum* (unc)

*G. laterale*

*G. mackayi* s.s.

*G. polycephala*

*G. travatsii*

*Helichrysum filicaule*

<i>Lagenifera cuneata</i> (unc)	<i>Chacemania latisiliqua</i> (incl. <i>C. gibbsii</i> ) (unc)
<i>L. petiolata</i> (unc)	
<i>L. ppmila</i>	
<i>L. strangulata</i>	<i>Colobanthus affinis</i>
<i>Leptinella</i> sp. (unnamed; included in <i>L. squalida</i> , as ssp. <i>mediana</i> , by Lloyd and Webb, 1987)	<i>C. apetalus</i> (incl. <i>C. a.</i> var. <i>alpinus</i> )
<i>Microseris scapigera</i> var.	<i>C. canaliculatus</i>
<i>Pseudognaphalium</i> sp.(i) (unnamed; aff. <i>P. luteoalbum</i> ) (lvs mostly linear-spathulate to linear, acute) (unc)	<i>C. sp.</i> (unnamed)†
<i>P. sp.</i> (ii) (unnamed; aff. <i>P. luteoalbum</i> ) (lvs mostly oblong-spathulate to narrowly oblong, rounded)	<i>Drosera spathulata</i>
<i>Raoulia glabra</i>	<i>Epilobium brunnescens</i> ss.
<i>Senecio glaucophyllus</i> ssp. <i>glaucophyllus</i>	<i>E. chlorifolium</i>
<i>S. waitavensis</i>	<i>E. glabellum</i> s.s.
<i>Taraxacum magellanicum</i>	<i>E. gracilipes</i>
	<i>E. komarovianum</i>
	<i>E. linnaeoides</i>
	<i>E. heteroides</i>
	<i>E. nummulariifolium</i> (unc)
	<i>E. pubens</i>
	<i>E. rotundifolium</i>
	<i>E. vernicosum</i> (?) (unc)
	<i>E. sp.</i> (unnamed; included in <i>E. brunnescens</i> , as ssp. <i>minutiflorum</i> , by Raven and Engelhorn, 1971)

#### DICOT HERBS OTHER THAN

#### COMPOSITES

<i>Acaena anserinifolia</i>	<i>Galium parposillum</i> ss.
<i>A. profundeincisa</i>	<i>Forstera tanella</i>
<i>Aciphylla</i> sp. (unnamed; aff. <i>A. glaucescens</i> ) (pinnae rigid)	<i>Gentiana filipes</i>
<i>Ahisotome aromatica</i>	<i>G. montana</i> (incl. <i>G. m.</i> var. <i>stolonifera</i> , <i>G. patula</i> , and <i>G. bellidi-folia</i> var. <i>pulchella</i> )
<i>A. pilifera</i> (unc)	<i>Getanum</i> sp. (aff. <i>G. microphyllum</i> )
<i>Cardamine</i> sp.(i) ( <i>C. debilis</i> agg.) (cf. "Narrow Patal" of Pritchard 1957)	<i>Geum leiospermum</i>
<i>C. sp.</i> (ii) ( <i>C. debilis</i> agg.) ("Glossy Leaf" of Pritchard 1957)	<i>Gingidia decipiens</i>
<i>C. sp.</i> (iii) (unnamed) (lvs mostly simple; blade ± reniform)	<i>G. montana</i>
<i>C. sp.</i> (iv) (aff. <i>C. corymbosa</i> )	<i>Gonocarpus aggregatus</i>

† (lvs and sepals strongly acicular;  
lvs unbordered; sepals 5, broader  
than lvs, without a thickened midrib)

- H.sp(ii)(unnamed)(included in *H. moschata*, as var. *parvifolia*, by Carse 1929)
- Hydrocotyle elongata*
- H. heterometa* (unc)
- H. microphylla*
- H. moschata*
- H. sp(i)(unnamed; included in *H. novae-zelandiae*, as var. *montana*, by Kirk 1899)*
- Kelliera dieffenbachii* (unc)
- K. laxa*
- Mazus radicans* (unc)
- Mentha cunninghamii*
- Montia fontana* s.s.
- Myosotis brockiei*
- *M. petiolata* s.s. (unc)
- M. spathulata* (incl. *M. s. var. radicata*) (unc)
- M. sp.(i)(unnamed; included in *M. pygmaea*, as var. *drueei*, by Moore 1961)* (unc)
- M. forstari*
- M. sp.(ii) (M. australis agg.)* (fls small, white) (unc)
- M. sp.(iii)(undetermined)*
- Neopaxia sp(i)(incl. *Montia calycina* and *Claytonia australasica* var. *biflora*)* (unc)
- N. sp.(ii)(unnamed)* (lvs spathulate, up to 15mm long; fls 6mm diam; calyx lobes broad, obtuse; corolla lobes rounded, cucullate)

- Nertera depressa*
- N. sp. (unnamed; aff. *N. dichondrifolia*)* (hairs present on both surfaces of lf, straight, not hooked as in *N. dichondrifolia*)
- Oreamyrtis colensoi* s.s.
- O. sp. (unnamed; aff. *O. ramosa*)*
- Outisia lactea* ssp. *lactea*
- O. modesta* (unc)
- Oxalis magellanica*
- Plantago lanigera* s.l. (incl. *P. novae-zelandiae*)
- P. piaoulii*
- P. triandra* s.s.
- Pratia angulata* agg.
- Psychrophila novae-zelandiae* (unc)
- Ranunculus cheesemanii* var. *cheesemanii* (unc)
- R. foliosus*
- R. glabrifolius* (unc)
- R. insignis*
- R. membranifolius*
- R. reflexus*
- R. toyi*
- R. simulans* (unc)
- R. verticillatus*
- Schizolejeunea nitens*
- S. trifoliolatum*
- Stackhousia minima*

*Stellaria decipiens* (incl. *S.  
parviflora*)  
*Urtica incisa*  
*Viola cunninghamii*  
*V. filicaulis*  
*V. lyallii*  
*Wahlenbergia albomarginata* var. *albomarginata* (incl.  
*W. pygmaea* var. *laxa*, *W.*  
*brockiei*, and *W. simpsonii*)

No. of species: 319..

## REFERENCES

- Allan, H.H. 1961: "Flora of New Zealand." Government Printer, Wellington. Vol. I. 1085 pp.
- Carsc, H. 1929: Trans. N.Z. Inst. 59: 315-6.
- Franklin, D.A. 1962: The Ericaceae in New Zealand (Gaultheria and Pernettya). Trans. R. Soc. N.Z. Bot. 1: 155-73.
- Kirk, T. 1899: "The Student's Flora of New Zealand". Government Printer, Wellington. 408 pp.
- Lloyd, D.G.; Webb, C.J. 1987: The reinstatement of Heptaptera at generic rank, and the status of the 'Cotuleae' (Asteraceae, Anthemideae). N.Z. Jl Bot. 25: 99-105.
- Moore, L.B. 1961: In "Flora of New Zealand." Government Printer, Wellington. Vol. I. 1085 pp.
- 1966: Australasian asteliads (Liliaceae). N.Z. Jl Bot. 4: 201-40.
- Pitchard, G.G. 1957: Experimental taxonomic studies of Cardamine Linn. in New Zealand. Trans. R. Soc. N.Z. 85: 75-89.
- Raven, P.H.; Engelkorn, T. 1971: New taxa and new combinations in Australasian Epilobium (Onagraceae). N.Z. Jl Bot. 9: 345-50.
- Wardle, P. 1968: The taxonomy and distribution of the stipulate species of Pseudopanax in New Zealand. N.Z. Jl Bot. 6: 226-36.
- Zотов, В.Д. 1965: Grasses of the subantarctic islands of the New Zealand region. Rec. Dom. Mus. Wellington 5: 101-46.
- Nordenstam, B. 1978: Opera Bot. 44: 1-84.
- Oliver, W.R.B. 1935: The genus Coprosma. Bishop. Mus. Bull. 132. 207 pp.
- Wardle, P. 1969: N.Z. Jl Bot. 7: 76-95.
- . —

