


HIGHER PLANTS (LYCOPODS, FERNS, GYMNOSPERMS, FLOWERING PLANTS)
OF N BALD KNOB RIDGE[†], N.W. NELSON, 950-1450 M

A.P.Druce, Wellington; P.A.Williams, Botany Division, D.S.I.R.,
 Nelson; S.P.Courtney, Nelson; May 1990 (revised June 1991,
 July 1992)

(Based on visit Feb. 1989)

[†]Highest point: Trent Pk, 1446 m

Rocks: calcareous mudstone
 medium sandstone
 limestone lenses
 granitic sandstone } all Tertiary



(unc) - uncommon or local

(L) - seen only near lake at M28 647612

* - seen only on granitic sandstone

INDIGENOUS PLANTS

GYMNOSPERM TREES AND SHRUBS

**Halocarpus biformis*

Lepidothamnus laxifolius

**Libocedrus bidwillii*

Phyllocladus alpinus agg. (unc, exc.
on granitic sandstone)

Podocarpus nivalis

Brachyglottis adamsii (unc)

B. bidwillii var. *vitidis*

B. rotundifolia var. (incl. *B. buchan-*
anii and *B. cockaynei*) (unc)

Cassinia vauvilliersii

DICOT TREES

Dracophyllum traversii (incl. *D.*
pyramidalis)

Griselinia littoralis (seedlings
only)

Nothofagus fusca

N. menziesii

N. solandri var. *cliffortioides*

N. fusca × *N. solandri* (unc)

Pseudopanax linearis (unc)

P. simplex (incl. *P. s. var. sinclairii*)

P. sp. (unnamed; included in *P.*) (unc)

colensoi, as var. *fjordensis*

 and as var. *ternatus*, by Wardle

(1968)

Coprosma atropurpurea (unc)

C. cheesemanii ss

C. ciliata ssp. *ciliata*

C. colensoi s.s.

C. crenulata (unc except at
N end of Ridge)

C. depressa

C. foetidissima

C. perpusilla ssp. *perpusilla*

C. propinqua ssp. *propinqua*

C. pseudocuneata ssp.

C. thamnoidea (incl. *C. polymorpha*)

C. sp. (unc)

rugosa ssp. *rugosa*

C. serrulata (unc)

C. sp.(i) (unnamed; aff. *C. pseudo-*
cuneata) (low-growing shrubs with
small, thick, arcuate lvs)

DICOT SHRUBS

Archeria traversii (incl. *A. t.* var.
australis) (unc, except on gran-
itic sandstone)

C. sp.(ii) (unnamed) (*C. parviflora*

var. *dumosa* sensu Allan 1961, non
C. p. var. dumosa Cheeseman)

<i>C. sp.(iii)</i> (unnamed; included in <i>C. parviflora</i> by Oliver, 1935, & others)*	<i>Helichrysum intermedium</i> var. (unc) <i>Leucopogon fasciculatus</i> var. (unc) (If undersurface not minutely hairy as is usual in <i>C. parviflora</i> ; L. <i>fraseri</i> s.s. drupe vs. unpigmented, occ. purple red)(unc)
<i>Comaria plumosa</i>	<i>L. suaveolens</i> (incl. <i>Cyathodes colensoi</i>)
<i>C. sarmentosa</i>	<i>Melicytus alpinus</i> s.s.
<i>Cyathodes dealbata</i> (incl. <i>C. pomila</i>)	<i>Myrsine divaricata</i> s.s.
<i>C. cuneata</i>	<i>M. nummularia</i> (unc)
<i>Dracophyllum longifolium</i> var. (incl. <i>D. filifolium</i> and <i>D. oliveri</i>)	<i>Olearia arborescens</i> (unc)
<i>D. uniflorum</i>	<i>O. avicenniifolia</i>
* <i>Gaultheria antipoda</i> (unc)	<i>O. lacunosa</i> var. <i>lacunosa</i>
<i>G. crassa</i>	<i>O. nummulariifolia</i> var. (vs distinctly longer than broad, rather than suborbicular as in var. <i>nummulariifolia</i>)(unc)
<i>G. depressa</i> s.s.	<i>Pentachondra pumila</i>
<i>G. sp.</i> (unnamed; included in <i>G. depressa</i> , as var. <i>novae-zelandiae</i> , by Franklin 1962)	<i>Pimelea suteri</i> var. (<i>P. orzophila</i>)
<i>G. crassa</i> × <i>G. sp.</i> (unnamed)	<i>Pittosporum rigidum</i> var. (<i>P. crassicaule</i>)
<i>Hebe canterburiensis</i>	<i>Pseudopanax anomalus</i> (unc)
<i>H. ciliolata</i>	<i>Pseudowintera colchica</i> (unc)
<i>H. gracillima</i> (unc)	<i>Traversia baccharoides</i> (unc)
<i>H. macrantha</i> var. <i>brachiphylla</i>	

DICOT LIANES AND RELATED

TRAILING PLANTS

<i>H. paucitamosa</i>	<i>Clematis paniculata</i> (unc)
<i>H. salicifolia</i> var. <i>salicifolia</i> (unc)	<i>Muehlenbeckia axillaris</i>
<i>H. tetragona</i> var. (<i>H. coarctata</i>)	<i>Rubus cissoides</i> var. (leaflets broader than in var. <i>cissoides</i>)
<i>H. topiaria</i>	<i>R. schmidelioides</i> var. <i>schmidelioides</i> (unc)
<i>H. vernicosa</i>	
<i>H. sp.(i)</i> (<i>Leonotis masoniae</i> *)	
<i>H. sp.(ii)</i> (aff. <i>H. glaucophylla</i>) (unc)	
<i>H. sp.(iii)</i> (cf. <i>matthewsii</i>) (unc)	

*(incl. *L. m.* var. *rotundata*)

LYCOPODS

Lycopodium australianum (unc)
L. fastigiatum
L. scariosum

ORCHIDS

Adenochilus gracilis (unc)
Chiloglottis cornuta
Corybas macranthus (unc)
C. trilobus s.s.

Microtis unifolia (L)
Prasophyllum colensoi (unc)
Pterostylis humilis (unc)
P. lisoniana (unc)
Thelymitra hatchii (unc)

FERNS

Asplenium flaccidum
Blechnum discolor (unc)
B. fluvatile agg.
B. penna-martini
B. procetum
B. sp. (unnamed) (mountain sp.;
 lower pinnae not reduced in
 length) ("The Dunedin-Cass
 Variety" of Crookes 1963)
Cyathea colensoi
Cystopteris tasmanica (unc)
Grammitis billardierei
G. magellanica ssp. *nolhofageti*
G. patagonica (unc)
G. poeppigiana
Histiopteris incisa (unc)
Hymenophyllum multifidum
H. malingii (unc, on dead beech)
H. ratum (unc)
H. villosum
Hypolepis millefolium
Leptopteris superba (unc)
Ophioglossum coriaceum agg.
Polystichum vestitum

GRASSES

Agrostis dyeri ss.
A. personata
A. sp. (unnamed; aff. A. dyeri)
 (panicle spreading, not contracted
 as in *A. dyeri*)
Chionochloa australis
C. conspicua ssp. *conspicua*
C. pallens ssp. *pallens*
C. tubra ssp. *occulta*
C. sp. (unnamed) (*C. flavescens*
 auct. South Island, non *C. flav-
 escens* Zотов)
C. pallens X *C. sp. (unnamed)* (unc)
Deschampsia tenella
Deyeuxia aucklandica (unc)
Elymus enysii
*E. sp. (aff. *E. rectisetus*)* (lvs
 glaucous; awns 4-6 cm long)

<i>Festuca</i> sp.(i) (cf. <i>F.matthewsii</i>)	SEDGES
(non tussock-forming, glaucous)	<i>Carex comans</i> (L)
<i>F.</i> sp.(ii)(unnamed; aff. <i>F.tubra</i>)	<i>C.dipsacea</i> var. <i>dipsacea</i> (L)
(unc)	<i>C.flagellifera</i> (L)
	<i>C.libera</i> (unc)
	<i>C.secta</i> s.s. (L)
<i>Hierochloe novae-zelandiae</i>	<i>C.sinclairii</i> (unc)
(unc)	<i>C.trachycarpa</i> (unc)
<i>Lachnagrostis</i> sp. (unnamed; included in <i>L.filiformis</i> , as var. <i>semiglabra</i> , by Zотов 1965)	<i>C.wakatipu</i> s.s. <i>C.sp.</i> (Section <i>Acutae</i>)
<i>Microlaena avenacea</i>	
<i>M.colensoi</i>	<i>Carpha alpina</i> s.s.
<i>Poa breviglumis</i>	<i>Eleocharis acuta</i> (L)
<i>P.colensoi</i>	<i>Gahnia procera</i> (unc, except on granitic sandstone)
<i>P.kirkii</i>	<i>Isolepis habra</i> (unc)
<i>P.maia</i>	<i>Oreobolus pectinatus</i>
<i>Rytidosperma gracile</i>	<i>O.strictus</i>
<i>R.setifolium</i>	<i>Schoenus pauciflorus</i> s.s.
<i>Thysatum youngii</i>	
<i>T.</i> sp.(i)(unnamed; included in <i>T.</i> <i>antarcticum</i> , as ssp. <i>tenellum</i> , by Petrie 1912)	<i>Uncinia astonic</i>
	<i>U.caespitosa</i>
	<i>U.clavata</i>
<i>T.</i> sp.(ii)(unnamed) (plants slender; lf sheath glabrous or nearly so; lf blade flat, up to 2.5mm wide, margins smooth)	<i>U.divaricata</i>
	<i>U.filiformis</i>
	<i>U.fuscovaginata</i> var. (plants glaucous)
<i>T.</i> sp.(iii)(unnamed; aff. <i>T.antarc-</i> <i>ticum</i>) (plants robust; lf blade flat, up to 7mm wide; margins scabrid) (unc)	<i>U.gracilenta</i>
	<i>U.involuta</i>
	<i>U.nervosa</i> s.s.
	<i>U.rupestris</i> (incl. <i>U.augustifolia</i>)
	<i>U.silvestris</i> (incl. <i>U.affinis</i>)

RUSHES

- Juncus gregiflorus* (L)
J. novae-zelandiae
Luzula migrata
L. subclavata (unc)
L. sp. (unnamed; included in
L. picta, as ssp. *limosa*, by
Edgar 1966)

Brachyglottis lagopus var. (cottony
hairs on both surfaces of lf, dense
at first, but becoming sparse)

Brachyscome radicans (unc)

Calmisia armstrongii

MONOCOT HERBS OTHER THAN**ORCHIDS, GRASSES, SEDGES, RUSHES**

- Astelia graminea* (unc)
A. linearis var. *linearis*
A. nervosa s.s.
A. nivicola s.s.
A. petriei (unc)
A. skottsbergii
A. sp.(i) (unnamed; included in
A. nivicola, as var. *moticeae*,
by Moore 1966)
A. sp. (ii) (unnamed; aff. *A. nervosa*)
(lvs broad, not conspicuously
white adaxially as in *A. ner-
vosa*)
Libertia pulchella (unc, ex-
cept on granitic sandstone)
* *Luzuriaga parviflora*
Phormium cookianum

C. bellidioides

C. dallii (N end of Ridge only)

C. hieracifolia ssp.

C. incana (incl. *C. discolor*)

C. laricifolia

C. montoi ssp. (*C. cornacea* var.
semicordata)

C. petriei

C. similis (N end of Ridge only)

C. sinclairii (incl. *C. albuni* & *C. durietzii*)

C. spectabilis ssp. (unc, except
N end of Ridge)

C. sp. (i) (unnamed; aff. *C. graci-
lenta*) (lvs linear to linear-
lanceolate, up to 45 x 1.2 cm;
lower surface white with
cottony hairs) (N end of ridge
only)

C. sp. (ii) (unnamed; aff. *C. similis*)

(plants much larger; possibly
originated from *C. montoi* X *C.*
similis, but plants uniform
and abundant) (N end of
Ridge only)

COMPOSITE HERBS

- Abrotanella linearis*
Anaphalis sp. (*Holochlysum*
bellidioides s.s.)

<i>C. armstrongii</i> X <i>C. petriei</i> (unc)	<i>Senecio glaucophyllus</i> ssp. <i>glaucophyllus</i>
<i>C. a.</i> X <i>C. spectabilis</i> (unc)	<i>S. wairavensis</i>
<i>C. monroi</i> X <i>C. petriei</i> (unc)	<i>Taraxacum magellanicum</i>
<i>C. similis</i> X <i>C. spectabilis</i> (unc)	
<i>C. taspedia</i> sp. (i) (unnamed) (plants summer-green; lvs green, both sur- faces densely clad in short gland- ular hairs; margins ± white with cottony hairs; fls white)	DICOT HERBS OTHER THAN COMPOSITES
<i>C. sp. (ii)</i> (unnamed) (plants summer- green; lvs green, both surfaces densely clad in multicellular non-glandular hairs with cottony tips; margins ± white with cottony hairs; fls white)	<i>Acaena auschalinifolia</i>
<i>Dolichoglottis lyallii</i>	<i>A. profunda incisa</i>
	<i>Aciphylla anomala</i>
	<i>A. colensoi</i> var. <i>colensoi</i>
	<i>A. ferox</i>
	<i>Anemone tenuicaulis</i>
	<i>Anisotome aromatica</i>
	<i>A. filifolia</i> (unc)
	<i>A. haastii</i>
<i>Gnaphalium audax</i>	<i>A. pilifera</i> (unc)
<i>G. laterale</i> (L.)	<i>Cardamine</i> sp. (i) (<i>C. debilis</i> agg.) ("Long Style" of Pritchard 1957)
<i>G. mackayi</i> s.s.	<i>C. sp. (ii)</i> (<i>C. debilis</i> agg.) (cf. "Tussock Race" of Pritchard 1957) (unc)
<i>Lagenifera cuneata</i>	<i>C. sp. (iii)</i> (<i>C. debilis</i> agg.) (L)
<i>L. pinnatifida</i> (unc)	<i>C. sp. (iv)</i> (unnamed) (lvs mostly simple, occ. with up to 2 pairs of pinnae; blades ± teniform) (unc)
<i>L. pumila</i> (L.)	
<i>L. strangulata</i>	
<i>Leptinella</i> sp. (i) (<i>L. pyrethrifolia</i> agg.)	
<i>L. sp. (ii)</i> (unnamed; included in <i>L. squalida</i> , as ssp. <i>mediana</i> , by Lloyd and Webb, 1987)	<i>Cheesemania latisiliqua</i> (incl. <i>C. gibbsii</i>) (unc)
<i>Leucogenes grandiceps</i> (unc)	<i>Colobanthus affinis</i>
<i>Microseris scapigera</i> var.	<i>C. apetalus</i> (incl. <i>C. a. van alpinus</i>)
<i>Pseudognaphalium</i> sp. (L.)	<i>C. squarrosum</i>
<i>Raoulia glabra</i>	<i>Crassula sinclairii</i> (L.)
<i>R. grandiflora</i> var. <i>grandiflora</i> *	<i>Drosera arcturi</i>
<i>R. tenuicaulis</i> (incl. <i>R. t.</i> var. <i>dimerpha</i> and <i>R. t.</i> var. <i>pusilla</i>)	<i>D. spathulata</i>
	* <i>D. stenopetala</i>

<i>Epilobium alsinoides</i> s.s. (unc)	<i>Geranium</i> sp. (unnamed; aff. <i>G. microphyllum</i>)
<i>E. atriplicifolium</i> (unc)	<i>G. sessiliflorum</i> var. <i>novae-zelandiae</i>
<i>E. chlorifolium</i>	
<i>E. cockayneanum</i>	<i>Geum leiospermum</i> (unc)
<i>E. gracilipes</i>	<i>G.</i> sp. (<i>G. parviflorum</i> agg.) (stock short; terminal leaflet about as long as broad; fls up to 13 mm diam)
<i>E. komarovianum</i> (L)	
<i>E. parnitens</i>	<i>Gingidia decipiens</i>
<i>E. pubens</i> (unc)	<i>G. montana</i>
<i>E. vernicosum</i>	
<i>E.</i> sp. (unnamed; included in <i>E. brunnescens</i> , as ssp. <i>minutiflorum</i> , by Raven and Engelhorn, 1971)	<i>Gonocarpus aggregatus</i> (unc)
<i>Euphrasia cheesemanii</i> (unc)	<i>Hydrocotyle heteromeria</i> (unc)
<i>E. townsonii</i> s.s.	<i>H. moschata</i> (unc)
<i>E. zelandica</i> agg.	<i>H. sulcata</i> (L)
<i>Forstera mackayi</i>	<i>H.</i> sp. (unnamed; included in <i>H. novae-zelandiae</i> , as var. <i>montana</i> , by Kirk 1899)
<i>F. tenella</i>	<i>Hypenticum japonicum</i> agg. (L)
<i>Galium petrosilium</i> s.s.	<i>Kelleria laxa</i>
<i>Gentiana filipes</i>	<i>K. multiflora</i>
<i>G. montana</i> (incl. <i>G. m.</i> var. <i>stolonifera</i> , <i>G. patula</i> , and <i>G. bellidifolia</i> var. <i>pulchella</i>) (unc)	<i>Limosella lineata</i> (L)
<i>G.</i> sp. (i) (undetermined; aff. <i>G. montana</i> and <i>G. townsonii</i>)	<i>Myosotis macrantha</i> (unc)
<i>G.</i> sp. (ii) (unnamed; aff. <i>G. bellidifolia</i> (plants forming small to large patches; lvs very glossy; fls large, cream or white))	<i>M.</i> sp. (i) (<i>M. australis</i> agg.) (fls small, white) (unc)
<i>G.</i> sp. (iii) (unnamed; aff. <i>G. corymbifera</i>)	<i>M.</i> sp. (ii) (unnamed; included in <i>M. pygmaea</i> , as var. <i>divaricata</i> , by Moore 1961)
	<i>Neopaxia</i> sp. (incl. <i>Montia calycina</i> and <i>Claytonia australasica</i>)
	<i>Nartera ciliata</i>
	<i>N.</i> sp. (unnamed; aff. <i>N. dichondrifolia</i>) (hairs present on both surfaces of lf, straight, not hooked as in <i>N. dichondrifolia</i>)

<i>Oreomytthis colensoi</i> s.s. (incl. <i>O.c.</i> var. <i>hispida</i> and <i>O.c.</i> var. <i>multifida</i>)	<i>Urtica incisa</i>
<i>O. sp.</i> (unnamed; aff. <i>O. ramosa</i>)	<i>Viola cunninghamii</i>
<i>Osmia caespitosa</i> ssp. <i>caespitosa</i> (unc)	<i>V. filicaulis</i>
<i>O. lactea</i> ssp. <i>lactea</i>	<i>V. lyallii</i> (L)
<i>Oxalis magellanica</i>	<i>Wahlenbergia albomarginata</i>
<i>Patahebe lyallii</i>	var. <i>albomarginata</i> (incl. <i>W. pygmaea</i> var. <i>laxa</i> , <i>W. brockiei</i> , and <i>W. simpsonii</i>)
<i>Plantago raoulii</i>	

ADVENTIVE PLANTS

<i>P. triandra</i> s.s. (L)	
<i>P. uniflora</i>	
<i>Potentilla anserinoides</i> (L)	
<i>Pratia angulata</i> agg.	
<i>Psychotria novae-zelandiae</i>	
<i>Ranunculus altus</i>	
<i>R. foliosus</i> (unc, except L)	
<i>R. insignis</i>	
<i>R. membranifolius</i>	
<i>R. verticillatus</i>	
<i>R. sp.</i> (unnamed; included in <i>R. lappaceus</i> , as var. <i>macto-</i>	
<i>phyllus</i> , by Kirk 1899)	
<i>Schizolema haastii</i> s.s.	
<i>S. sp.</i> (unnamed; included in <i>S. haastii</i> , as ssp. <i>cyanopetalum</i> , by Domin 1908) (unc)	
<i>Scleranthus brockiei</i>	
<i>S. uniflorus</i>	
<i>Stellaria decipiens</i> (incl. <i>S.</i> <i>parviflora</i>)	

GRASSES

<i>Agrostis capillaris</i> (unc)
<i>Dactylis glomerata</i> (unc)
<i>Festuca nigrescens</i>
<i>Holcus lanatus</i>
<i>Poa annua</i>
<i>P. pratensis</i>

SEDGES

<i>Carex ovalis</i>

RUSHES

<i>Juncus articulatus</i> (L)
<i>J. bufonius</i> (L)
<i>J. effusus</i> (L)
<i>J. sp.</i> (<i>J. conglomeratus</i> ?) (unc)

COMPOSITE HERBS

<i>Cirsium arvense</i>
<i>C. vulgare</i> (unc)
<i>Crepis capillaris</i> (L)
<i>Gnaphalium coarctatum</i> (L)

Hieracium pilosella
H. sp. (undetermined) (unc)
Hypochoeris radicata
Leucanthemum vulgare (unc)
Mycelis muralis
Senecio jacobaea (L)
Sonchus asper (L)
Taraxacum officinale s.l.

Galium palustre ssp. palustre (L)
Linum catharticum (unc)
Lotus pedunculatus (L)
Mimulus moschatus (L)
Myosotis laxa ssp. caespitosa (L)
Plantago major (L)
Prunella vulgaris (L)
Ranunculus repens (L)
Rumex acetosella
R. crispus (L)
Sagina procumbens (L)
Trifolium dubium (L)
T. repens
Veronica serpyllifolia (L)

DECOT HERBS OTHER THAN
COMPOSITES

Anagallis arvensis (L)
Cerastium fontanum ssp. vulgare
Digitalis purpurea (unc)
Epilobium ciliatum (L)

No. of indigenous spp.: 311.
No. of adventive spp.: 41.

REFERENCES

- Allan, H.H. 1961: "Flora of New Zealand". Government Printer, Wellington.
 Vol. I. 1085 pp.
- Crookes, M. 1963: "New Zealand Ferns". Whitcombe & Tombs. 407 pp.
- Domin, K. 1908: Engl. Bot. Jk 40: 573-85.
- Edgar, E. 1966: Luzula in New Zealand. N.Z. Jl Bot. 4: 159-84.
- Franklin, D.A. 1962: The Ericaceae in N.Z. Trans R. Soc. NZ. Bot. 1: 155-73
- Kirk, T. 1899: "The Student's Flora of New Zealand", Government Printer, Wellington. 408 pp.
- Lloyd, D.G.; Webb C.J. 1987: The reinstatement of Leptinella at generic rank,
 & the status of the 'Cotyled' (Asteraceae, Anthemideae). N.Z. Jl Bot. 25: 99-105.
- Moore, L.B. 1961: In: "Flora of New Zealand". Government Printer,
 Wellington. Vol. 1085 pp.
- 1966: Australasian asteliads (liliaceae). N.Z. Jl Bot. 4: 201-40.
 (cont on p. 11)

- Oliver, W.R.B. 1935: The genus Coprosma. Bishop Mus. Bull. 132. 207 pp.
- Petrie, D. 1912: Trans. N.Z. Inst. 44: 179-87.
- Pritchard, G.G. 1957: Experimental taxonomic studies on species of Ceratamine Linn. in New Zealand. Trans. Roy. Soc. N.Z. 85: 75-89.
- Raven, P.H.; Engelhorn, T. 1971: New taxa and new combinations in Australasian Epilobium (Onagraceae). N.Z. Jl Bot. 9: 345-50.
- Wardle, P. 1968: Taxonomy and distribution of the stipulate species of Pseudopanax in New Zealand. N.Z. Jl Bot. 6: 226-36.
- Zotov, V.D. 1965: Grasses of the subantarctic islands of the New Zealand region. Rec. Dom. Mus. Wellington 5: 101-46.

