


HIGHER PLANTS (FERNS, FLOWERING PLANTS) OF LIMESTONE AREAS AT HEAD OF LOVERIDGE CK, S. ARTHUR RA., N.W. NELSON, 1300-1500 M

Grid ref. NZMS 260 M27 790950

A.P.Druce, Wellington; P.A.Williams, D.S.I.R. Land Resources, Nelson; S.P.Courtney, D.O.C., Nelson, June 1991 (revised July 1992)

(Based on visit Feb. 1991)

Rocks: Summit Limestone //



(unc) - uncommon

INDIGENOUS PLANTSDICOT SHRUBS*Aristotelia fruticosa* var.*Brachyglottis adamsii**B. bidwillii* var. *vitidis**Cassinia vauvilliersii**Coprosma atropurpurea* (unc)*C. cheesemanii* s.s.*C. depressa**C. rugosa* ssp. *rugosa**C. serrulata**Coriaria plumosa**C. p. X C. Sarmantosa**Gaultheria* sp. (unnamed; included in *G. depressa*, as var. *novae-zelandiae*, by Franklin 1962)*Hebe albicans* var. *albicans**H. ciliolata**H. cockayneana* var.*H. haastii* var. (aff. var. *humilis*; see Moore 1961, p. 940)*H. macrantha* var. *brachyphylla**H. teopatria**H. sp.* (unnamed) (lvs up to 4x1cm, elliptic-oblong, bright green; upper Lachnagrostis sp. (unnamed; surface very glossy; corolla tube very short, = or < calyx; corolla lobes obtusa; capsule pubescent) included in *L. filiformis*, as var. *semiglabra*, by Zofov 1965)*H. albicans* X *H. sp.* (unnamed) (unc)*Helichrysum intermedium* var.*Leucopogon fraseri* s.s.*Melicytus alpinus* s.s.*Pimelea suteri* var. (*P. oreophila*)FERNS*Asplenium lyallii**Blechnum penna-marina**Cystopteris tasmanica**Grammitis patagonica**G. poeppigiana**Ophioglossum eoriaceum* agg.*Polystichum vestitum*GRASSES*Chionochloa pallens* ssp. *pallens**C. sp.* (unnamed) (*C. flavesans* avct. South Island, non *C. flavesans* Zofov)*Elymus* sp. (aff. *E. rectisetus*) (plants glaucous; awns 4-6 cm long)*Festuca* sp. (cf. *F. matthewsii*) (plants tufted but not tussock-forming, glaucous)*Koeleria cheesemanii**Lachnagrostis* sp. (unnamed; included in *L. filiformis*, as var. *semiglabra*, by Zofov 1965)*Microlaena colensoi**Poa cita* ssp. *cita**P. colensoi*

P. subvestita

Rytidosperma setifolium

Trisetum sp. (unnamed; included *Gnaphalium audax* (unc)

in *T. antarcticum*, as ssp. *tanellum*,

by Petrie (1912)

ly clad in short glandular hairs, with or without longer multicellular, non-glandular hairs with cottony tips; margins conspicuously white with cottony hairs; fls white)

SEDGES

Carex wakatipu s.s.

Schoenus pauciflorus s.s.

Uncinia divaricata

U. fuscovaginata var. (plants glaucous)

Lagenifera concata

L. petiolata (unc)

Leptinella sp. (*L. pyrethrifolia* agg.)

Microseris scapigera var.

Raoulia apicinigra s.s. (unc)

Senecio glaucophyllus ssp. *glaucophyllus*

S. waitavensis

RUSHES AND ALLIED PLANTS

Luzula migrata

Marsippospermum gracile

DICOT HERBS OTHER THAN

COMPOSITES

Acaena fissistipula

A. profunda incisa

Aciphylla anomala

A. ferox

Anisotome pilifera

Cardamine sp.(i) (*C. debilis* agg.)

(cf. "Long Style" of Pritchard 1957)

C. sp.(ii) (unnamed) (lvs mostly simple; blade + reniform)

C. sp.(iii) (undetermined)

Cheesemania latisiliqua (incl. *C. gibbsii*)

Colobanthus affinis

C. canaliculatus

C. strictus

Epilobium chlorifolium

E. glabellum s.s.

E. gracilipes

COMPOSITE HERBS

Anaphalis sp. (*Helichrysum bellidoides* s.s.)

Calmsia dallii

C. monroi ssp.

C. sinclairii (incl. *C. allanii* and *C. duriotzii*)

Craspedia sp. (unnamed) (plants summer-green; lvs up to 18cm long green, with short broad petiole merging into lamina; both surfaces dense-

<i>E. matthewsii</i>	<i>Schizeilema</i> sp. (unnamed; included in <i>S. haastii</i> , as ssp. <i>cyanopetalum</i> , by Domin 1908)
<i>E. pubens</i>	
<i>E. sp.</i> (unnamed; included in <i>E. brunneocans</i> , as ssp. <i>minutiflorum</i> , by Raven and Engelhorn, 1971)	<i>Scleranthus brockiei</i>
	<i>S. uniflorus</i>
<i>Euphrasia cheesemanii</i>	<i>Stellaria decipiens</i> (incl. <i>S. parviflora</i>)
<i>Gentiana bellidifolia</i> var.	<i>Wahlenbergia albomarginata</i>
<i>G. filipes</i>	var. <i>albomarginata</i> (incl. <i>W. pygmaea</i> var. <i>alexae</i> , <i>W. brockiei</i> and <i>W. simpsonii</i>)
<i>G. sp.</i> (unnamed; aff. <i>G. corymbifera</i>)	
<i>Geranium</i> sp. (unnamed; aff. <i>G. microphyllum</i>)	

ADVENTIVE PLANTS

<u>GRASSES</u>	
<i>Grevillea</i> sp. (<i>G. parviflora</i> agg.)	<i>Anthoxanthum odoratum</i>
(stock short; terminal lfts about as long as broad; fls upto 13mm)	<i>Dactylis glomerata</i>
<i>Grevillea decipiens</i>	<i>Festuca nigrescens</i>
<i>G. montana</i>	<i>Poa pratensis</i>
<i>G. decipiens</i> X <i>G. montana</i> (unc)	

<u>COMPOSITE HERBS</u>	
<i>Myosotis</i> sp. (i) (<i>M. australis</i> agg.) (fls small, white) (unc)	<i>Hieracium lepidulum</i>
<i>M. sp.</i> (ii) (unnamed; included in <i>M. pygmaea</i> , as var. <i>drucei</i> , by Moore 1961)	<i>H. pilosella</i>
<i>Oreomyrrhis colensoi</i> ss. (incl. <i>O.c.</i> var. <i>hispida</i> and <i>O.c.</i> var. <i>multifida</i>)	<i>H. sp.</i>
<i>Oritzia lactea</i> ssp. <i>lactea</i>	<i>Hypochaeris radicata</i>
<i>O. simpsonii</i>	<i>Mycelis muralis</i>
<i>Parahelbe lyallii</i>	<i>Taraxacum officinale</i> s.l.

<u>DICOT HERBS OTHER THAN COMPOSITES</u>	
	<i>Ceratium fontanum</i> ssp. <i>vulgare</i>
	<i>Prunella vulgaris</i>
	<i>Trifolium repens</i>
	<i>Veronica</i> sp. (<i>V. arvensis</i> ?)

No. of indigenous spp: 101.

No. of adventive spp: 14.

REFERENCES

- Domin, K. 1908: Botanische Jahrbuch 40: 573-85.
Franklin, D.A. 1962: T.R.S.N.Z. Bot.J.: 155-73.
Moore, L.B. 1961: In: "Flora of N.Z." Vol. 1.
Petrie, D. 1912: T.N.Z. Inst. 44: 178-87.
Pritchard, G.C. 1957: T.R.S.N.Z. 85: 75-89.
Raven, P.H.; Engelhorn, T. 1971: N.Z. Jl Bot. 9: 345-50.
- . —

