

HIGHER PLANTS (LYCOPODS, FERNS, GYMNOSPERMS, FLOWERING PLANTS) OF SCHIST AREAS* AT HEAD OF IRONSTONE CK,
WEST OF DEVIL R. THRUST, S. ARTHUR RA, N.W. NELSON,
1100 - 1586 M



Highest point: Loveridge Peak, grid ref. NZMS 260 M27 790945

A.P.Druce, Wellington; P.A.Williams, D.S.I.R. Land Resources, Nelson; S.P.Courtney, D.O.C. Nelson.

June, 1991 (revised July 1992, Aug. 1993)

Based on visit Feb. 1991

Rocks: Waingaro Schist ==
 * usually calcareous



(unc) - uncommon or local
INDIGENOUS PLANTS

[GYMNOSPERM TREES AND SHRUBS]

Podocarpus nivalis

Gaultheria crassa

Hebe albicans var. *albicans*

H. ciliolata

H. cockayneana var.

H. haastii var. (aff. var. *humilis*;
 see Moore 1961, p. 940)

H. macrocarpa var. *brachyphylla*

H. tetragona var. (*H. coarctata*)

H. topiaria

Helichrysum intermedium var.

Leucopogon fraseri s.s.

Melicytus sp. (unnamed)

Myrsine divaricata s.s.

M. nummularia

Olearia arborescens (unc)

O. avicenniifolia

O. colensoi var. *edentata*

O. nummulariifolia var. (lvs distinctly longer than broad, rather than suborbicular as in var. *nummulariifolia*)

Pentachondra pumila

Traversia baccharoides

[DICOT SHRUBS]

Aristotelia fruticosa var.

Brachyglottis adamsii

B. bidwillii var. *ritidis*

Cassiniaaustralis

Coprosma cheesemanii s.s.

C. ciliata ssp. *ciliata*

C. depressa

C. perpusilla ssp. *perpusilla*

C. rugosa ssp. *rugosa*

C. serrulata

C. sp. (unnamed; aff. *C. colensoi*)

(low-growing shrubs with trailing branches)

Coriaria angustissima

C. plumosa

Cyatodes dealbata (incl. *C. pumila*)

Dracophyllum uniflorum

[DICOT LIANES AND RELATED]

[TRAILING PLANTS]

Muehlenbeckia axillaris

[LYCOPODS]

Lycopodium fastigiatum

L. scariosum

FERNS

- Asplenium flaccidum*
A. richardii
Blechnum sp. (unnamed) (mount-
 ain sp.; lower pinnae not reduc-
 ed in length) ("The Dunedin-
 Cass Variety" of Crookes 1963)
Cystopteris tasmanica
Grammitis patagonica
Hypolepis millefolium
Ophioglossum coriaceum agg. (unc)
Polystichum vestitum

Lachnagrostis sp. (unnamed;
 included in *L. filiformis*, as var.
semiglabra, by Zotov 1965)

- Microlepia colensoi*
Poa breviglumis
P. celsa
P. cita, ssp. *cita*

P. colensoi
P. imbecilla (unc)
P. subvestita

P. sp. (unnamed; aff. *P. novae-zelandiae*) (scree plants with narrow lvs)
 (unc)

GRASSES

- Agrostis dyeri* s.s.
A. sp. (unnamed; aff. *A. dyeri*)
 (panicle broad, not contracted
 as in *A. dyeri*) (unc)
Chionochloa australis
C. conspicua ssp. *conspicua*
C. pallens ssp. *pallens*

C. rubra ssp. *occulta*
C. sp. (unnamed) (*C. flavescens*
 avct. South Island, non *C. flav-
 escens* Zotov)

Elymus sp. (aff. *E. rectisetus*)
 (plants glaucous; awns 4-6 cm long)
Festuca sp. (i) (cf. *F. matthewsii*)
 (tufted but not tussock-forming,
 glaucous)

F. sp. (ii) (unnamed; aff. *F. rubra*)
Microchloa novae-zelandiae

Rytidosperma gracile

R. setifolium

Trisetum youngii

T. sp. (unnamed) (plants slender;
 If sheath glabrous or nearly so)

SEDGES

Carex sinclairii

C. wakatipu s.s.

C. sp. (unnamed; aff. *C. coriacea*)
 (lvs slender, bright green, not
 becoming stiff and yellow-green
 in exposed sites)

Carpha alpina s.s.

Isolepis caligenis

Uncinia divaricata

U. fuscovaginata var. (plants glaucous)

U. involuta

U. sp. (unnamed) (*U. affinis* sensu)

Hamlin 1959, non *U. riparia* var.
affinis Col. ex Clarke; see Nooter-
boom 1978)

Schoenus pauciflorus s.s. (= "slender
var." of Edgar 1970, p. 194)

RUSHES AND ALLIED PLANTS

Luzula migrata

**MONOCOT HERBS OTHER GRASSES,
SEDGES, RUSHES + ALLIED PLANTS**

Astelia nervosa s.s.

A. nivicola s.s.

A. petriei

Libertia peregrinans

Phormium cookianum

C. sp.(ii) (unnamed; aff. *C. gracilenta*)
(plants tufted; lvs linear to linear-
lanceolate, up to 4.5 x 1.2 cm; lower
surface white with cottony hairs)

Craspedia sp.(i) (unnamed) (plants sum-
mergreen; lvs up to 18 cm long, green,
with short broad petiole merging
into lamina; both surfaces densely
clad in short glandular hairs, with
or without longer multicellular non-
glandular hairs with cottony tips;
margins conspicuously white with
cottony hairs; fls white)

C. sp.(ii) (plants summer-green; lvs
green; both surfaces sparsely clad in
in multicellular non-glandular hairs;
midrib on undersurface glabrous; fls
white)

Gnaphalium mackayi s.s.

G. ruahinicum

Lagenifera cuneata

Leptinella pyrethrifolia agg.

B. bellidioides s.s. (unnamed; aff. *B. bellidioides*, as var. *B. sp.* (unnamed; aff. *L. squalida*)
(*pinnae entire or nearly so*)

Brachyscome sinclairii s.s.

Calochortus dallii

C. incana (incl. *C. discolor*)

C. laticifolia

C. monroi ssp.

C. sessiliflora

C. sinclairii (incl. *C. callanii* and
C. durietzii)

C. spectabilis ssp.

C. sp.(i) (unnamed; aff. *C. graci-
lenta*) (plants strongly rhizo-
matous)

Microrhynchus scapigerus var.

Pseudognaphalium sp. (unnamed;
aff. *P. luteoalbum*) (mountain sp.;
lvs mostly oblong-spathulate to
narrowly oblong, rounded)

Raoulia glabra

Senecio glaucocephalus ssp. *glauco-
cephalus*

DICOT HERBS OTHER THAN COMPOSITES

- Acacia anserinifolia*
- A. profundaincisa*
- Aciphylla anomala*
- A. ferox*
- A. glaucescens* s.s.
- Anisotome deltoides*
- A. haastii*
- Cardamine* sp.(i) (*C. debilis* agg.)
(cf. "Long Style" of Pitchard.
1957)
- C. sp.(ii)* (unnamed) (lvs mostly
simple; blade ± raniform)
- Cheesemania latisiliqua*
(incl. *C. gibbsii*)
- Colobanthus canaliculatus*
- C. strictus* (unc)
- Epilobium chlorifolium*
- E. elegans* (unc)
- E. matthewsii*
- E. pubens*
- E. sp.(i)* (unnamed; aff. *E. glab-*
illum) (unc)
- E. sp.(ii)* (unnamed; included in
E. brunneescens, as ssp. *minuti-*
florum, by Raven and Engel-
horn, 1971)
- Euphrasia cheesemanii*
- E. zealandica* agg. (corolla 4-5 mm
wide)
- E. sp.(i)* (unnamed; aff. *E. laingii*)
- E. sp.(ii)* (unnamed; aff. *E. pefrici*)
- Forstera mackayi*
- F. tanella*
- Gentiana bellidifolia*
- G. filipes*
- Geranium* sp. (unnamed; aff. *G.*
microphyllum)
- Gum* sp. (*G. parviflorum* agg.)
(stock short; terminal leaflet
about as long as broad; fls up
to 13 mm diam.)
- Gingidia montana*
- Gunnera monoica*
- Hydrocotyle* sp. (unnamed;
included in *H. novae-zealandiae*,
as var. *montana*, by Kirk 1899)
- Kellaria laxa*
- K. multiflora*
- Montia fontana* s.s.
- Myosotis* sp.(i) (*M. australis* agg.)
(fls small, white)
- M. sp.(ii)* (unnamed; included in
M. pygmaea, as var. *drucei*, by
Moore 1961)
- Nertera ciliata*
- Ostasia caespitosa* ssp. *caespit-*
osa
- O. lactea* ssp. *lactea*
- O. simpsonii*
- Oxalis magellanica*
- Parahelbe linifolia* ssp. *linifolia*
- P. lyallii*
- Plantago lanigera* s.l. (incl. *P. n.z.*)
- P. raoulii*

P. uniflora
Pratia angulata agg.
Ranunculus altus
R. foliosus (unc)
R. insignis
R. mitis (unc)
R. verticillatus
R. sp. (undetermined)
Schizolema haastii s.s.
S. sp. (unnamed; included in
S. haastii, as ssp. *cyanopetal-*
um, by Domin 1908)
Scleranthus uniflorus
Stellaria decipiens (incl. *S.*
paniflora)
S. gracilenta
Urtica incisa
Viola cunninghamii
V. lyallii
Wahlenbergia albomarginata
 var. *albomarginata* (incl. *W.*
pygmaea var. *laxa* and
W. brockiei)

ADVENTIVE PLANTS

GRASSES

Agrostis capillaris
Anthoxanthum odoratum
Dactylis glomerata
Festuca nigrescens

RUSHES

Juncus articulatus
J. effusus

COMPOSITE HERBS

Cirsium vulgare
Hieracium lepidulum
H. pilosella
H. sp.
Hypochoeris radicata
Mycelis muralis
Taraxacum officinale s.l.

DICOT HERBS OTHER THAN COMPOSITES

Ceratostium fontanum ssp. *vulgare*
Digitalis purpurea
Rumex acetosella
Sagina procumbens
Trifolium repens

No. of indigenous spp. : 174.

No. of adventive spp. : 18.

REFERENCES

- Crookes, M. 1963. "Flora of N.Z." Vol. 1.
- *Domin, K. 1908. *Bot. Jahr.* 40: 573-85.
- Edgar, E. 1970. In "Flora of N.Z." Vol. 1.
- Kirk, T. 1899. "The Student's Flora of N.Z."
- Moore, L.B. 1961. In "Flora of N.Z." Vol. 1.
- Nootboom, H.P. 1978. *Blumea* 24: 511-20.
- Nordenstam, B. 1978. *Opera Bot.* 44: 1-84.
- Pritchard, G.G. 1957. *T.R.S. N.Z.* 85: 75-89.
- Raven, P.H.; Engelhorn, T.: (1971). *N.Z. Jl. Bot. Q.* 345-50.
- Wardle, P. 1968. *N.Z. Jl. Bot.* 6: 226-36.
- Zotov, V.D. 1965. *Rec. Dom. Mus. Leningrad.* 5: 1044.

* Hamlin, B.G. 1959. *Dom. Mus. Bull.* 19.