

HIGHER PLANTS (LYCOPODS, FERNS, GYMNOSPERMS, FLOWERING PLANTS) IN VICINITY OF KAWEKA LAKES<sup>†</sup>, EASTERN FOOT OF KURIPAPANGO HILL, KAWEKA RANGE, 600-750 M.

<sup>†</sup>Grid ref. N123 81-55-.

A.P.Druce, Botany Division, D.S.I.R., 1972  
 (Revised: Apr. 1974, Dec. 1974, July 1975,  
 June 1976, May 1977, Oct. 1977, June 1980,  
 Oct. 1981, Oct. 1982, May 1984, June 1984, July 1985,  
 March 1986, April 1986, June 1990, Judy 1992)

(Based on visits 1957, 1960, 1964, 1974, 1984, 1986)  
 (unc - uncommon)

INDIGENOUS SPECIES

GYMNOSPERM TREES AND SHRUBS

*Phyllocladus alpinus* ss.(unc)

*Pseudopanax arboreus* var.  
 arboreus (unc)  
*P.colensoi* s.s. (unc)

*Podocarpus hallii* X

*P.nivalis* (unc)

MONOCOT TREES

*Cordyline australis*

DICOT SHRUBS

*Brachyglossis repanda* ss. (unc)  
*Carmichaelia* sp. (subgenus  
*Carmichaelia*) (unc)

DICOT TREES

*Carpodetus serratus*

*Cassinia vauvilliersii*

*Griselinia littoralis* (unc)

*Coprosma acerosa* var. (unc)

*Nothofagus fusca* (unc)

*C. cheesemanii* ss

*N.solandri* (unc)

*C. propinqua* ssp. *propinqua*

*N.fusca* X *N.solandri* (unc)

*Pittosporum ralphii*

*P.tenuifolium* ssp. *tenuifolium* (unc)

*P.ralphii* X *P.tenuifolium* (unc)

<i>C. thamnoides</i> (incl. <i>C. polymorpha</i> )	<i>Hebe stricta</i> var. <i>stricta</i>
	<i>Helichrysum aggregatum</i> var. <i>aggregatum</i> (unc)
<i>C. robusta</i> (unc)	<i>Kunzea ericoides</i> agg.
<i>C. rugosa</i> ssp. <i>rugosa</i>	
<i>C. sp. (i)</i> (unnamed; aff. <i>C. cheesemanii</i> ) (lvs broader than in <i>C. cheesemanii</i> ) (unc)	<i>Leptospermum scoparium</i> <i>Leucopogon fasciculatus</i> var. <i>L. fraseri</i> s.s.
<i>C. sp. (ii)</i> (unnamed; included in <i>C. parviflora</i> by Oliver, 1935, and others) (lf undersurface not minutely hairy as is usual in <i>C. parviflora</i> ; drupes vs. un- pigmented, occ. purplish red)	<i>Olearia arborescens</i> (unc) <i>O. furfuracea</i> var.
<i>C. propinqua</i> X <i>C. robusta</i>	<i>O. nummularifolia</i> var. <i>nummularifolia</i>
<i>Coriaria arborea</i>	
<i>C. kingiana</i>	<i>Pimelea longifolia</i> var. <i>longifolia</i> (unc)
<i>C. arborea</i> X <i>C. kingiana</i> (unc)	<i>P. prostrata</i>
<i>C. a.</i> X <i>C. pteridoides</i> (unc)	
<i>Cotoneaster</i> cotoneaster var. cotoneaster (unc)	<b>DICOT LIANES AND RELATED PLANTS</b> <i>Clematis forsteri</i>
<i>Cyathodes empetrifolia</i>	<i>Muehlenbeckia australis</i>
<i>C. sp. (Juniperina agg.)</i> (lvs vs. L 1 cm. X L 1 mm)	<i>M. axillaris</i>
<i>Dracophyllum subulatum</i>	<i>M. australis</i> X <i>M. axillaris</i> (unc)
<i>Epacris alpina</i>	<i>Rubus cissoides</i> var. (leaflets narrower than in var. <i>cissoides</i> ) (unc)
<i>Gaultheria antipoda</i>	
<i>G. macrostigma</i>	
<i>G. sp.</i> (unnamed; included in <i>G.</i> depressa, as var. <i>novae-zelandiae</i> , by Franklin 1962)	
<i>G. antipoda</i> X <i>G. sp.</i> (unnamed)	<b>LYCOPODS</b> <i>Lycopodium australe</i> (unc)
<i>G. a.</i> X <i>G. macrostigma</i>	<i>L. fastigiatum</i>

*L. scariosum*

*L. volubile*

### FERNS

*Adiantum cunninghamii* (unc)

*Asplenium flabellifolium* agg (unc)

*A. flaccidum*

*A. hookerianum*

*A. terrestre* ssp. *terrestre*

*A. flaccidum* X *A. hookerianum*  
(unc)

*Blechnum chambersii*

*B. colensoi* (unc)

*B. fluviatile* agg.

*B. penna-martini*

*B. ptocerum*

*B. minus*

*B. vulcanicum*

*B. sp.* (unnamed) (common  
sp.; lower pinnae reduced in  
length)

*Bottleychium austiale* (unc)

*Ctenopteris heterophylla* (unc)

*Dicksonia fibrosa* (unc)

*Gleichenia dicarpa*

*Histiopteris incisa*

*Hymenophyllum cypresiforme*  
(unc)

*H. multifidum*

*Hypolepis ambigua*

*Leptopteris hymenophylloides*  
(unc)

*Ophioglossum coriaceum* agg.  
*Paeonia scaberula*

*Pellaea rotundifolia* (unc)

*Phymatosorus postulatus* (unc)

*Polystichum richardii*

*P. vestitum* (unc)

*Pteridium esculentum*

*Pyrrhisia elatagnifolia* (unc)

### ORCHIDS

*Aporostylis bifolia* (unc)

*Caladenia carneola* agg.

*Chiloglottis cornuta*

*Corybas macranthus*

*C. oblongus* (unc)

*C. rivularis* s.s. (unc)

*C. thilobus* s.s.

*Gastrolechia cunninghamii*

*G. minor* (unc)

*Gnaphelioides nudum* (unc)

*Microtis unifolia*

*Orthoceras novae-zealandiae*

*Prasophyllum colensoi*

*Pterostylis foliata* (unc)

*P. micromega* (unc)

*P. patens*

*P. sp.* (unnamed; aff. *P. montana*)

*Spiranthes novae-zealandiae*

*Thelymitra cyanea*  
*T. decora*  
*T. hatchii* (unc)  
*T. longifolia*  
*T. pauciflora*

*P. imbecilla*

*Rytidosperma clavatum*  
*R. gracile*  
*R. setifolium*  
*R. viride*

### GRASSES

*Amphibromus fluitans* (unc)  
*Chionochloa rubra* ssp. *rubra*  
*Cortaderia fulvida*  
*Deschampsia caespitosa* var.  
*macrantha*  
*Deyeuxia avenoides* (incl. *D. a.*  
*var. brachyantha*)  
*D. sp.* (*D. quadriseta* agg.)  
*Dichelachne crinita*  
*Echinopogon ovatus*

*Elymus* sp. (aff. *E. rectisetos*)  
 (Group Waicuru of Löve and  
 Connor, 1982)

*Hierochloe redolens*  
*Lachnagrostis filiformis* s.s.  
*L. striata*  
*L. sp.* (unnamed; included in *L.*  
*filiformis*, as var. *semiglabra*,  
 by Zofov 1965)  
*L. filiformis* × *L. striata*  
*Mirolaena stipoides*  
*Poa anceps* ssp. *anceps*  
*P. cita* ssp.  
*P. colensoi*

*Trisetum* sp. ( ) (unnamed; aff.  
*T. antarcticum*) (lf sheath  
 densely hairy)  
*T. sp. ( )* (unnamed; aff. *T. ant-*  
*arcticum*) (plants robust; lf  
 sheath glabrous at least in upper  
 part; lf blade margins scabrid)  
*T. sp. (c)* (unnamed) (plants slender;  
 lf sheath glabrous or nearly so)

### SEDGES

*Baumea rubiginosa*  
*B. tenax* (unc)  
*Carex breviculmis*  
*C. cirtosa* (unc)  
*C. coriacea*  
*C. dipsacea* var. (*C. tahoata*)  
*C. dissita* s.s.  
*C. echinata* var. (*C. stellulata*  
 var. *australis*)  
*C. fascicularis* (unc)  
*C. flagellifera*  
*C. gaudichaudiana*  
*C. invertsa*  
*C. maorica*  
*C. secta* s.s.  
*C. sinclairii*

*C.solandri*  
*C.testacea* s.s. (unc)  
*C.virgata*

*Carpha alpina* ss.  
*Eleocharis acuta*  
*E.gracilis*  
*E.sphacelata*  
*Isolepis distigmatosa*  
*I.fluitans*  
*I.pottsii*  
*I.tetricularis*  
*I.subtilissima*  
*Lepidosperma australe*  
*Schoenus maschalinus*  
*Uncinia rubra*  
*U.tupectris* (incl. *U.angustifolia*)  
*U.scabra*  
*U.uncinata*

### RUSHES

*Juncus graciliflorus*  
*J.planifolius*  
*J.novae-zelandiae*  
*Luzula decipiens* (unc)  
*L.migrata*  
*L.picta* s.s.  
*L.subclavata* (unc)

### MONOCOT HERBS OTHER THAN ORCHIDS, GRASSES, SEDGES, RUSHES

*Arthropodium candidum* (unc)  
*Astelia fragrans* (unc)  
*Herpolitha novae-zelandiae* (unc)  
*Lemna* sp. (*L.minor* auct.)  
*Phormium cookianum*  
*P.fenax*  
*Potamogeton cheesemanii*  
*P.suboblongus*  
*Triglochin striata*  
*Typha orientalis*

### COMPOSITE HERBS

*Anaphalis* spp. (see p. 6)  
*Brachyglossis lagopus* var. *lagopus* (unc)  
*Celmisia glandulosa* ssp.  
*glandulosa* (unc)  
*C.gracilenta* ssp.  
*C.spectabilis* ssp. *spectabilis*  
*C.* sp. (unnamed; aff. *C.gracilenta*  
 (plants strongly rhizomatous))  
*Craspedia minor* var. *minor*  
 (incl. *C.major*)

*C.viscosa*  
*C.* sp. (unnamed) (summer-green  
 bog plant; capitula 1.0-1.5 cm  
 diam.)  
*Gnaphalium avicinum*  
*G.delicatulum*  
*G.gymnocephalum*  
*G.involucratum*

<i>G. linosum</i> s.s.	<i>Anisotome aromatica</i> (unc)
<i>G. sphaericum</i>	
<i>Helichrysum filicaule</i>	
<i>Anaphalis</i> sp.(i) (unnamed; aff. <i>A. kerriensis</i> ) (lvs shorter, and capitula smaller, than in <i>A. kerriensis</i> )	
<i>A. sp.(ii)</i> ( <i>Helichrysum bellidoides</i> ss)	<i>Cardamine</i> sp.(i) ( <i>C. debilis</i> agg.) ("Long Style" of Pritchard 1957)
<i>A. sp.(iii)</i> (unnamed) (" <i>Helichrysum alpinum</i> " of Cockayne 1928) (lvs and capitula larger than in <i>A. sp.ii</i> )	<i>C. sp.(ii)</i> ( <i>C. debilis</i> agg.) ("Narrow Petal" of Pritchard 1957)
<i>Lagenifera pumila</i>	<i>C. sp.(iii)</i> ( <i>C. debilis</i> agg.) ("Glossy Leaf" of Pritchard 1957)
<i>Microseris scapigera</i> var.	<i>Centella uniflora</i>
<i>Pseudognaphalium</i> sp. ( <i>P. luteo-album</i> agg.) (lvs mostly linear-spathulate to linear, acute)	<i>Dichondra repens</i> agg.
<i>Raoulia albosericea</i> (unc)	<i>Drosota binata</i>
<i>R. tenerrima</i> (incl. <i>R.t. var.dimorpha</i> and <i>R.t. var. pusilla</i> )	<i>D. peltata</i> ssp. <i>auriculata</i>
<i>R. glabra</i> X <i>Helichrysum filicaule</i> (unc)	<i>Epilobium alsinoides</i> ss.
<i>Seneio glaucophyllus</i> ssp. <i>toa</i>	<i>E. atriplicifolium</i> s.s.
<i>S. minimus</i>	<i>E. brunneascens</i> s.s.
<i>S. quadridentatus</i>	<i>E. billardiereanum</i> s.s. (unc)
<i>S. sp.</i> (unnamed; included in <i>S. glaucophyllus</i> , as ssp. <i>discoideus</i> , by Onduff 1960)	<i>E. chionanthum</i>
<i>Vittadinia australis</i> agg.	<i>E. cinereum</i>
	<i>E. insulare</i>
	<i>E. komarovianum</i>
	<i>E. glabellum</i> S.S. (unc)
	<i>E. melanocaulon</i> (unc)

#### DICOT HERBS OTHER THAN COMPOSITES

<i>Acaena anserinifolia</i>	<i>E. microphyllum</i> (unc)
<i>A. sp.</i> ( <i>A. emittans</i> or <i>A. juvenca</i> )	<i>E. heteroides</i>

*Aciphylla* sp. (unnamed; aff. *A. squarrosa*) (unc)

<i>E.nummularifolium</i>	<i>H.sp.(unnamed)</i> (much smaller than <i>H.japonicum</i> )
<i>E.pedunculare</i> s.s. (unc)	<i>Hypselia rivalis</i>
<i>E.pubens</i>	<i>Lilacopsis ruthiana</i>
<i>E.refundifolium</i>	
<i>E.tenuipes</i> s.s.	<i>Linum monogynum</i>
<i>E.sp. (unnamed; included in <i>E.brunn-Mentha cunninghamii</i> escens, as ssp. <i>minutiflorum</i>, by <i>Montia fontana</i> s.s. Raven and Engelhorn, 1971)</i>	<i>Myriophyllum pedunculatum</i> ssp. <i>novae-zealandiae</i>
<i>Euphrasia cuneata</i>	
<i>Galium perpusillum</i> s.s.	<i>M.propinquum</i>
<i>G.propinquum</i>	<i>M.triphyllum</i>
<i>G.trilobum</i> (unc)	<i>Nertera ciliata</i>
<i>G.sp. (unnamed; aff. <i>G.perpusillum</i>)</i> (lvs wider than stipules, narrowly lanceolate; apex rounded to subacute, bristles 0-2)	<i>N.depressa</i> (incl. <i>N.cunninghamii</i> )
<i>Gentiana grisebachii</i> s.s. (unc)	<i>N.setulosa</i>
<i>Geranium microphyllum</i>	<i>Oreomyrsis</i> sp.(i) (unnamed; aff. <i>O.ramosa</i> )
<i>G.potentilloides</i> var. <i>potentilloides</i>	<i>O.sp.(ii)</i> (unnamed; included in <i>O.colensoi</i> , as var. <i>delicatula</i> , by Allan 1961) (unc)
<i>G.sessiliflorum</i> var. <i>novae-zealandiae</i>	<i>Durisia macrophylla</i> ssp. <i>robusta</i> (unc)
<i>Glossostigma elatinoides</i>	<i>Oxalis exilis</i>
<i>Gonocarpus aggregatus</i>	
<i>G.micranthus</i> ssp. <i>micranthus</i>	<i>Plantago raoulii</i>
<i>Gratiola sexdentata</i>	
<i>Gunnera protegens</i>	<i>Potentilla anserinoides</i>
<i>Hydrocotyle elongata</i>	<i>Pratia angulata</i> agg.
<i>H.hydrophila</i>	<i>P.perpusilla</i>
<i>H.microphylla</i>	<i>Ranunculus amphitrichus</i>
<i>H.moschata</i>	<i>R.glabrifolius</i>
<i>H.novae-zealandiae</i> s.s.	<i>R.limosella</i> (unc)
<i>Hypericum japonicum</i> agg.	<i>R.mactopus</i> (unc)
	<i>R.reflexus</i>

*Rorippa palustris*  
*Rumex flexuosus* (unc)  
*Scleranthus uniflorus* (unc)  
*Stackheuscia minima*  
*Urtica incisa*  
*Viola cunninghamii*  
*V. lyallii*  
*Wahlenbergia albomarginata*  
 var. *pygmaea*  
*W. colensoi*  
*W. rupestris*

No. of indigenous species: 290.

### ADVENTIVE SPECIES

#### GRASSES

*Agrostis capillaris*  
*Aira caryophyllea*  
*Anthoxanthum odoratum*  
*Bromus hordeaceus*  
*Dactylis glomerata*  
*Festuca arundinacea* sl. (unc)  
*F. nigrescens*  
*Holcus lanatus*  
*Poa pratensis*  
*Rytidosperma racemosum*  
*Vulpia bromoides*

#### RUSHES

*Juncus articulatus*  
*J. bufonius*  
*J. effusus*  
*J. tenuis*

#### MONOCOT HERBS OTHER THAN GRASSES, SEDGES, RUSHES

*Ottelia ovalifolia*  
*Potamogeton crispus*

#### COMPOSITE HERBS

*Centipeda cunninghamii*  
*Cirsium arvense*  
*C. vulgare*  
*Conyza sp.*  
*Crepis capillaris*  
*Gnaphalium spicatum*

#### SEDGES

*Carex ovalis*

*Hieracium pilosella*  
*Hypochoeris radicata*  
*Leontodon taraxacoides*  
*Mycelis muralis*

*Myosotis laxa* ssp. *caespitosa*  
*Pelargonium inodorum*  
*Plantago major*  
*Polygonum hydropiper*  
*P. prostratum*

*Senecio jacobaea*  
*S. sylvatica*  
*Sonchus asper*  
*Taraxacum officinale* s.l.

*Prunella vulgaris*  
*Ranunculus repens*  
*R. sp. (R. fluitans avct. N.Z.)*  
*Rumex acetosella*

**DICOT HERBS OTHER THAN/  
 COMPOSITES**

*Acaena agnipila*  
*A. novae-zelandiae*  
*Anagallis arvensis*  
*Actium sp. (unc)*  
*Arenaria serpyllifolia*  
*Centaurium erythraea*  
*Cerastium fontanum* ssp. *vulgare*  
*Epilobium ciliatum*  
*Hyperticum humifusum*  
*Linum catharticum*  
*Lotus pedunculatus*

*Sagina procumbens*  
*Solanum americanum*  
*Trifolium dubium*  
*T. repens*  
*Verbascum creticum*  
*V. thapsis*  
*Veronica sp.*  
*Wahlenbergia marginata*

No. of adventive species: 62.

**REFERENCES**

- Ornduff, R. 1960: An interpretation of the *Senecio lautus* complex in N.Z. *T.R.S.N.Z.* 80: 63-77.
- Allan, H.H. 1961: "Flora of New Zealand." Government Printer, Wellington. Vol. I. 1085 pp.
- Cockayne, L. 1928: "The Vegetation of New Zealand." 2nd ed. Englemann. 456 pp.
- Franklin, D.A. 1962: The *Ericaceae* in New Zealand (*Gaultheria* and *Perpetua*). *Trans. Roy. Soc. N.Z. Bot.* 1: 155-73.
- Oliver, W.R.B. 1935: The genus *Coprosma*. *Bishop Mus. Bull.* 132 + 207 pp.
- Pitchard, G.G. 1957: Experimental taxonomic studies on species of *Cardamine* Linn. in New Zealand. *Trans. Roy. Soc. N.Z. Bot.* 85: 75-89.
- Raven, P.H.; Englehorn, T. 1971: New taxa and new combinations in Australasian *Epilobium* (Onagraceae). *N.Z.J. Bot.* 9: 345-50.
- Zotov, V.D. 1965: Grasses of the subantarctic islands of the New Zealand region. *Rec. Dom. Mus. Wellington* 5: 101-46.
- Löve, A.; Connor, H.E. 1982: Relationships & taxonomy of N.Z. wheatgrasses. *N.Z.J. Bot.* 20: 169-86.

