


INDIGENOUS HIGHER PLANTS (PSILOPSIDS, LYCOPODS, FERNS, GYMNO-SPERMS, FLOWERING PLANTS) OF MT PUREORA†, HAUHUNGAROA RANGE, WEST TAUPO, 2300-3823 FT (700 -1170 M)

†N93 123543, alt. 3823 ft (c.1170m)

A.P. Druce, Botany Division, D.S.I.R., 1972
(Revised April 1974, July 1975, June 1976, April 1977,
May 1977, November 1977, November 1978, June 1979,
March 1980, June 1980, May 1983*, Aug. 1983*, June 1984,
Nov. 1984, April 1985, May 1985, July 1985, Aug. 1990)
*Incorporating observations by B.R. Clarkson 1982-3

Based on visits 1959 (A.P.D.), 1960 (A.P.D.),
1982-3 (B.R.C.), and 1985 (A.P.D.)



GYMNOSPERMS

Dacrycarpus dactydioides (seedlings)
Dacrydium cupressinum
Halocarpus bidwillii
Phyllocladus alpinus

P. glaucus
Podocarpus hallii
Prumnopitys ferruginea
P. taxifolia

MONOCOT TREES

Cordyline australis
C. indivisa

DICOT TREES

Aristotelia serrata
Beilschmiedia tawa
Carpodetus serratus
Elaeocarpus dentatus (incl. *E. d.*
var. obovatus)
E. hookerianus
Fuchsia excorticata
Griselinia littoralis
Hedycarya arborea
Melicytus lanceolatus (incl. *M. l.* var. *C. sp. (C. juniperina agg.)* (lvs us. $< 1 \text{ cm}$
latior)
M. ramiflorus ssp. *ramiflorus*
Myrsine salicina
Nestegis cunninghamii
N. lanceolata
Olearia rani var. *colorata*
Pennantia corymbosa
Pseudopanax arboreus var. *arboreus*
P. colensoi ss.
P. crassifolius
P. edgerleyi
P. simplex (incl. *P. s.* var. *sinclairii*)
Quintinia serrata (incl. *Q. acuti-*
folia and *Q. elliptica*)
Schefflera digitata
Weinmannia racemosa var. *racemosa*

DICOT SHRUBS

Alseuosmia pusilla
Aristotelia fruticosa var. *microphylla*
Brachyglottis repanda ss.
B. rotundifolia var. (*B. elaeagnifolia*)
B. kirkii ss.
Cassinia leptophylla var. (*C. vauvilliersii*)
Coprosma foetidissima
C. grandifolia
C. lucida ss.
C. rhamnoides ssp. *rhamnoides* (incl. *C. polymorpha*)
C. rigida
C. robusta
C. rugosa ssp. *rugosa*
C. tenuifolia
C. sp. (i) (unnamed; aff. *C. cheesem-*
manii) (lvs broader than in *C.*
cheesemania)
C. sp. (ii) (unnamed; included in
C. parviflora by Oliver, 1935,
 and others (lf undersurface
 not densely hairy as in *C. parvi-*
flora; drupes us. unpigmented,
 occ. purplish red)
Cotokia buddleioides (unc)
Cyathodes empetrifolia
C. sp. (C. juniperina agg.) (lvs us. $< 1 \text{ cm}$
 $\times < 1 \text{ mm}$)
Dracophyllum subulatum
Gaultheria antipoda
G. sp. (unnamed; included in *G.*
depressa, as var. *novae-zeeland-*
iae, by Franklin 1962)
G. sp. (unnamed) \times *Pernettya*
macrostigma
Hebe macrocarpa var. (*H.*
corriganii)
H. stricta var. *stricta*
Kunzea ericoides var.
Leptospermum scoparium var.
Leucopogon fasciculatus

Melicope simplex
Myrsine divaricata s.s.
Neomyrtus pedunculata
Olea arborescens
O. ilicifolia
O. arborescens x *O. ilicifolia* (incl.
O. macrodonta)
Pentachondra pumila
Pseudopanax anomalus
P. a. x *P. simplex*
Pseudowintera axillaris
P. colorata

DICOT LIANES

Clematis foetida
C. paniculata
Metrosideros diffusa
Muehlenbeckia australis
Parsonia capsularis var.
Rubus cissoides
R. schmidelioides var. *schmideli-*
oides

PSILOPSIDS AND LYCOPODS

Tmesipteris elongata (incl. *T. a.* ssp.
robusta)
T. tannensis
Lycopodium australianum
L. fastigiatum
L. laterale (incl. *L. ramulosum*?)
L. scariosum
L. varium (incl. *L. billardierei* and
L. novae-zelandicum)
L. volubile

FERNS

Asplenium bulbiferum
A. flaccidum
A. gracillimum
A. hookerianum

A. oblongifolium
A. polyodon
Blechnum chambersii
B. colensoi
B. discolor
B. fluviatile agg.
B. membranaceum
B. minus
B. nigrum
B. penna-marina
B. procerum
B. vulcanicum
B. sp. (unnamed) (common lowland
 sp.; lower pinnae reduced in
 length)

Ctenopteris heterophylla
Dicksonia fibrosa
D. lanata s.s.
D. squarrosa
Gleichenia dicarpa
Grammitis billardierei
G. magellanica ssp. *nottofageti*
Histopteris incisa
Hymenophyllum armstrongii
H. bivalve
H. demissum
H. dilatatum
H. ferrugineum
H. flabellatum
H. multifidum
H. pulcherrimum
H. rarum
H. revolutum
H. sanguinolentum agg.
H. scabrum
Hypolepis rufobarbata
H. millefolium
Lastreopsis glabella
Leptolepia novae-zelandiae
Leptopteris hymenophylloides
L. superba
L. hymenophylloides x *L. superba*

- Lindsaea trichomanoides*
- Ophioglossum coriaceum* agg.
- Paesia scaberula*
- Phymatosorus diversifolius*
- P. novae-zelandiae*
- Polystichum silvaticum*
- P. vestitum*
- Pteridium esculentum*
- Pyrosia cleagnifolia*
- Rumohra adiantiformis*
- Sticherus cunninghamii*
- Trichomanes teniforme*
- T. venosum*

ORCHIDS

- Aporostylis bifolia*
- Chiloglottis cornuta*
- Corybas trilobus* s.s.
- Eanna autumnalis*
- E. mucronata* s.s.
- Microtis oligantha*
- Prasophyllum colensoi*
- P. sp. (aff. P. patens)*
- Pterostylis banksii*
- P. graminea*
- P. sp. (aff. P. montana)*
- Thelymitra cyanea*
- T. sp.*

GRASSES

- Cortaderia fulvida*
- Deyeuxia aucklandica*
- D. avenoides* (incl. *D. a. var. brachyantha*)
- D. sp. (D. quadrisetata agg.)*
- Heterochloa redolens*
- Microlaena avenacea*
- Poa breviglumis*
- Rytidosperma gracile*

SEDGES

- Caumea rubiginosa*
- B. tenax*
- Carex dissita* s.s.
- C. echinata* var. (*C. stellulata* var. *australis*)
- C. lessoniana*
- C. testacea* s.s.
- Carpha alpina* s.s.
- Fleocharis gracilis*
- Gahnia procera*
- Lepidosperma australe*
- Oreobolus pectinatus*
- Schoenus maschalinus*
- Isolopis crassiuscula*
- I. distigmata*
- I. habra*
- I. pottsii*
- I. subtilissima*
- Uncinia ferruginea*
- U. filiformis*
- U. gracilentata*
- U. rupestris* (incl. *U. angustifolia*)
- U. uncinata*
- U. zotovii*
- U. sp. (U. affinis sensu Hamlin 1959, non U. hipatia var. affinis Col. ex Clarke)*

RUSHES AND ALLIED PLANTS

- Centrolepis ciliata*
- Juncus antarcticus*
- J. gregiflorus*
- J. novae-zelandiae*
- J. planifolius*
- Luzula picta* s.s.
- L. subclavata*
- L. sp. (unnamed; included in L. picta, as var. limosa, by Edgar 1966)*

MONOCOT HERBS OTHER THAN ORCHIDSGRASSES, SEDGES, RUSHES

Astelia fragrans
A. grandis
A. sp. (unnamed; aff. *A. nervosa*) (lvs not conspicuously white adaxially as in *A. nervosa*)
Collospermum microspermum
Dianella nigra
Herpolition novae-zelandiae
Luzuriaga parviflora
Potamogeton suboblongus
Triglochin striata

COMPOSITE HERBS

Celmisia gracilentata var.
C. sp. (unnamed; aff. *C. gracilentata*)
Gnaphalium audax
G. delicatum
G. sphaericum unnamed;
Pseudognaphalium sp. (aff. *P. luteocal-
 bom*) (lvs mostly linear-spathulate to linear, acute)
Helichthysum filicaule
Anaphalis sp. (i) (cf. *A. katiensis*)
A. sp. (ii) (unnamed) ("Helichthysum alpinum" of Cockayne 1928)
Lagenifera pumila
Raoulia glabra
Senecio minimus
S. rufiglandulosus var.

DICOT HERBS OTHER THAN COMPOSITES

Acaena ansetinifolia
Cardamine sp. (*C. debilis*) ("Long Style" of Pritchard 1957)

Drosera binata
D. spatulata
Epilobium alsinoides s.s.
Gentiana grisebachii s.s.
Geranium microphyllum
G. potentilloides var. *potentilloides*
E. cockayneanum

Gonocarpus aggregatus
G. micranthus ssp. *micranthus*

Hydrocotyle heteromeia
H. microphylla
H. moschata
Jovellana repens
Kelleria dieffenbachii
Nertera ciliata
N. scapanioides
Oreomyrrhis sp. (*O. colensoi* or *O. tamosa*)
Ourtisia macrophylla var. *robusta*
Pratia angulata agg.
Ranunculus amphitrichus
R. reflexus
Stellaria decipiens (incl. *S. parviflora*)
Urtica incisa
Utricularia novae-zelandiae

Viola cunninghamii
V. filicaulis

No. of species: 244.

REFERENCES

- Hamlin, B.G. 1959: A revision of the genus *Uncinia* in N.Z. Dom. Mus. Bull. 19: 106 pp.
 Cockayne, L. 1928: The Vegetation of N.Z. 2nd ed. Englemann, Leipzig. 456 pp.
 Franklin, D.A. 1962: The Ericaceae in N.Z. (*Gaultheria* & *Pernettya*). Trans. R. Soc. N.Z. Bot. 1: 155-73
 McKelvey, P.J. 1963: N.Z. Forest Service Bull. 14: 127 pp.
 Oliver, W.R.B. 1935: The genus *Ceprosoma*. Bishop Mus. Bull. 132: 207 pp.
 Pritchard, G.G. 1957: Experimental taxonomic studies on species of *Cardamine* Linn. in N.Z. Trans. R. Soc. N.Z. 85: 75-89.
 Edgar, E. 1966: *huzula* in New Zealand. N.Z. J. Bot. 4: 159-84.
 Record not accepted: *Polystichum cystostegia* (McKelvey 1963, p. 122)

