

VASCULAR PLANTS OF MT RAUKUMARA<sup>†</sup>, RAUKUMARA RA.,  
3000-4638 FT (900-1400 M)

A.P.Druce, Botany Division, D.S.I.R., and M. Heginbotham, Opotiki  
 (1972; revised Apr. 1974, June 1976, May 1977, June 1979, May 1980,  
 Aug. 1981, April 1983, Feb 1984, July 1985, Aug 1990)  
 Based on visits 1950 (A.P.D.), 1981 (M.H.), and 1983 (A.P.D.)

<sup>†</sup> Grid ref. N71 530465 ✓

INDIGENOUS PLANTS

GYMNOSPERM TREES AND SHRUBS

<i>Halocarpus biformis</i>	<i>Ixerba brexioides</i>
<i>Lepidothamnus laxifolius</i> (Creswell 1961, as <i>Dactylium laxifolium</i> )	<i>Melicytus lanceolatus</i> (incl. M.l. var. <i>laticor</i> )
<i>Libocedrus bidwillii</i>	
<i>Phyllocladus hallii</i>	<i>M. ramiflorus</i> ssp. <i>ramiflorus</i>

<i>Podocarpus hallii</i>	<i>Myrsine salicina</i>
<i>P. nivalis</i> (Creswell 1961)	
<i>Prumnopitys ferruginea</i>	

MONOCOT TREES

<i>Cordyline indivisa</i>	
---------------------------	--

DICOT TREES

<i>Aristotelia serrata</i>	
<i>Carpodetus serratus</i>	
<i>Dracophyllum traversii</i> (incl. <i>D. pyramidalis</i> )	<i>folia</i> and <i>Q. elliptica</i>
<i>Fuchsia excorticata</i>	<i>Schefflera digitata</i>
<i>Griselinia littoralis</i>	<i>Weinmannia racemosa</i>
<i>Hochstetia populnea</i> var. <i>lanceolata</i>	

### DICOT SHRUBS

- Alseuosmia pusilla*  
Brachyglottis repanda s.s.  
*B. rotundifolia* var. (*B. elaeagnifolia*)  
*B. Kirkii* ss.  
  
*Coprosma colensoi* s.s.  
*C. foetidissima*  
*C. grandifolia*  
*C. pseudocuneata* ssp. *pseudocuneata*  
*C. tenuifolia*  
*C. sp.* (unnamed; included in *C. parviflora* by Oliver, 1935, and others)  
(fl surface not minutely hairy as  
is usual in *C. parviflora*; drupes vs. un pigmented, occ. purplish red)
- Olearia arborescens*  
*O. colensoi* var. *colensoi*  
*O. ilicifolia*  
*O. nummularifolia* (Cruswell 1961)  
*O. arborescens* × *O. ilicifolia*  
*Pentachondra pumila*  
*Pimelea buxifolia*  
*Pittosporum rigidum* var. *rigidum*  
  
*Pseudopanax anomalous*  
*P. a. × P. simplex*  
*Pseudowintera colorata*

*Cyathodes ampetrifolia*  
*Diacophyllum sinclairii* (incl. *D. adamsii* and *D. vitide*)

*Gaultheria antipoda*

*G. paniculata*

*G. sp.* (unnamed; included in *G. depressa*, as var. *novae-zelandiae*, by Franklin 1962)

*G. paniculata* × *G. sp.* (unnamed)

*Hebe macrocarpa* var. (*H. cortiganii*)

*Myrsine divaricata* s.s.

*Olearia arborescens*

*O. colensoi* var. *colensoi*

*O. ilicifolia*

*O. nummularifolia* (Cruswell 1961)

*O. arborescens* × *O. ilicifolia*

*Pentachondra pumila*

*Pimelea buxifolia*

*Pittosporum rigidum* var. *rigidum*

### DICOT LIANES

*Clematis paniculata*

*Metrosideros diffusa*

*M. fulgens*

*Muehlenbeckia australis*

*Rubus cissoides* var. *cissoides*

### LYCOPODS

*Lycopodium australianum*

*L. fastigiatum*

*L. scariosum*

*L. variuum* (incl *L. billardierei* and

*L. novae-zelandicum*)

### FERNS

*Asplenium bulbiferum*

*A. flaccidum*

*A. bulbiferum* × *A. flaccidum*

*Blechnum chambersii*

*B. discolor*

*B. fluviatile* agg.

*B. procerum*

*B. sp. (unnamed)* (common lowland sp.; lower pinnae reduced in length) *Thelymitra* sp.

*B. procerum* X *B. sp. (unnamed)*

*Cyathea colensoi*

*C. smithii*

*Dicksonia squarrosa*

*Grammitis billardierei*

*G. magellanica* ssp. *nothofageti*

*Histoiopteris incisa*

*Hymenophyllum demissum*

*H. flabellatum*

*H. multifidum*

*H. peltatum*

*H. tarum*

*H. revolutum*

*H. sanguinolentum* agg.

*Hypolepis rufobarbata*

*Leptopteris superba*

*Paxisia scaberrula*

*Phymatosorus diversifolius*

*P. novae-zelandiae*

*Pneumatopteris pennigera*

*Polystichum vestitum*

## ORCHIDS

*Aporostylis bifolia*

*Chiloglottis cornuta*

*Cotybas trilobus* s.s.

*Dendrobium cunninghamii*

*Eatinà autumnalis*

*E. mucronata* ss.

*Prasophyllum colensoi*

*Pterostylis banksii*

*P. humilis*

*P. venosa*

*Thelymitra* sp.

## GRASSES

*Chionochloa pallens* var. (lvs broader than in var. *pallens*)

*Deyeuxia aucklandica*

*Lachnagrostis* sp. (unnamed; included in *L. filiformis*, as var. *semi-glabra*, by Zofov 1965)

*Microseris avenacea*

*M. stipoides*

*Poa breviglumis*

*P. colensoi*

*Rytidosperma gracile*

*Trisetum* sp. (unnamed; aff. *T. antarcticum*) (lf sheath distinctly hairy)

## SEDGES

*Carex dissita* ss.

*C. forsterti*

*C. solandri*

*C. sp. (Sect. Acutae)*

*Carpha alpina* s.s.

*Gahnia procera*

*G. sp. (G. pauciflora or G. setifolia)*

*Isolepis habra*

I. pottsii

Oreobolus pectinatus

Schoenus maschalinus

Uncinia caespitosa

U. clavata

U. ferruginea

U. filiformis

U. gracilenta

U. rupestris (incl. U. angustifolia)

U. silvestris (incl. U. affinis)

U. uncinata

U. zotovic

### RUSHES

Juncus grægiflorus

J. planifolius

Luzula migrata

L. subclavata

L. sp. (unnamed; included in L. picta, as var. limosa; by Edgar 1966)

### MONOCOT HERBS (OTHER THAN ORCHIDS, GRASSES, SEDGES, RUSHES)

Astelia fragrans

A. solandri

A. sp. (unnamed; aff. A. nervosa) (lvs not conspicuously white adaxially as in A. nervosa)

Colospermum microspermum

Libertia pulchella

Luzuriaga parviflora

### COMPOSITE HERBS

Anaphalis sp. (unnamed; aff. Helichrysum bellidioides) ("H. alpinum" of Cockayne 1928) (lvs

and capitula larger than in H. bellidioides)

Gnaphalium avolax

G. gymnocephalum

G. tuahinicum

G. sp. (unnamed; aff. G. limosum) (lvs linear)

Helichrysum filicaule

Pseudognaphalium sp. (unnamed; aff. P. luteoalbum)

Raoulia tenuicaulis (incl. R.t. var. dimorpha and R.t. var. pusilla)

Senecio minimus

### DICOT HERBS (OTHER THAN COMPOSITES)

Acaena anserinifolia

Anisotome aromatica

Cardamine sp. (C. debilis agg.) ("Glossy Leaf" of Pritchard 1957)

Epilobium alsinoides s.s.

E. brunneascens s.s.

E. cockayneana

E. heteroides

E. nummularifolium

E. linnaeoides

Euphrasia cuneata

Gentiana sp. (unnamed; aff. G. grisebachii)

*Gonocarpus micranthus* ssp.  
*micranthus*

*Hydrocotyle heteromeria*  
*H. microphylla*

*Nertera ciliata*

*N. sp. (unnamed; aff. N. dichondri-* *T. effusus*  
*folia) (hairs present on both surfaces of* *T. tenuis*  
*If straight, not hooked as in N. dichondr.)*

*Oreomythris colensoi* (Creswell 1961) *Achillea millefolium*  
*Ourisia lactea* ssp. *drucei*

*Oxalis magellanica* agg.

*Plantago taoulii* var. (lvs narrow-  
elliptic)

*Pratia angulata* ass.

*Ranunculus membranifolius*

*R. reflexus*

*Stellaria decipiens* (incl. *S. parviflora*)

*Urtica incisa*

*Viola filicaulis*

*Dactylis glomerata*  
*Festuca nigrescens*  
*Holcus lanatus*  
*Poa annua*  
*Vulpia bromoides*

**RUSHES**

*Juncus articulatus*

*J. effusus*

*J. tenuis*

**COMPOSITE HERBS**

*Cirsium arvense*

*C. palustre*

*C. vulgare*

*Conyza floribunda*

*Crepis capillaris*

*Hypochoeris radicata*

*Leontodon taraxacoides*

*Senecio jacobaea*

*S. sylvaticus*

*S. vulgaris*

*Sonchus asper*

**DICOT HERBS (OTHER THAN COMPOSITES)**

*Callitricha stagnalis*

*Cerastium fontanum* ssp. *vulgare*

*Digitalis purpurea*

*Linum catharticum*

*Lotus pedunculatus*

*Prunella vulgaris*

*Ranunculus repens*

*Rumex acetosella*

*Sagina procumbens*

*Veronica serpyllifolia*

No. of species: 177.

ADVENTIVE PLANTS

**DICOT SHRUBS**

*Eruca* sp.

**GRASSES**

*Agrostis capillaris*

*A. stolonifera*

*Aira caryophylla*

*Anthoxanthum odoratum*

No. of species: 35.

Record not accepted:

*Alseuosmia quercifolia* (Druce 1954) [= *A. pusilla*]

REFERENCES

- Cockayne, L. 1928: "The Vegetation of New Zealand". 2nd ed. Engleman, Leipzig. 456 pp.
- Creswell, R.D. 1961: An ascent of Mt Rauhukura. Bull. Wellington Bot. Soc. 32: 8-9.
- Druce, A.P. 1954: Plant distribution records(2). Bull. Wellington Bot. Soc. 27: 14-9.
- Druce, D.G. 1972: The cluster and solitary-headed cudweeds native to New Zealand: (Gnaphalium Section Euclatia, - Compositae). N.Z. Jl Bot. 10: 112-79.
- Edgar, E. 1966: Luzula in New Zealand. N.Z. Jl Bot. 4: 159-84.
- Franklin, D.A. 1962: The Ericaceae in New Zealand (Gaultheria and Petitia). Trans. R. Soc. N.Z. Bot. 1: 155-73.
- Kirk, T. 1899: "The Student's Flora of New Zealand.". Government Printer, Wellington. 408 pp.
- Moore, L.B. 1961: Hebe (except Whipple spp.). Pp. 885-952 in "Flora of New Zealand". Government Printer, Wellington. Vol. I. 1085 pp.
- Oliver, W.R.B. 1935: The genus Coprosma. Bishop Mus. Bull. 132. 207 pp
- Zotov, V.D. 1965: Grasses of the subantarctic islands of the New Zealand region. Rec. Dom. Mus. Wellington 5: 101-46.