


VASCULAR PLANTS OF MAUNGAHARURU RANGE¹, TE WAKA RANGE², AND
MT HUKANUI³

A.P.Druce, Botany Division, D.S.I.R., 1971 (revised Sept. 1974 May 1979
(Based on observations 1960, 1969-73, 1980, 1985, 1989) Oct. 1974 July 1980
July 1975 May 1981
July 1976 June 1981
May 1977 May 1982
Oct. 1977 July 1982
Nov. 1987
June 1988
March 1989
Nov. 1989
May 1990
Nov. 1990

¹ Highest point: Mt Taraponui, grid ref. N 114 171828, alt. 4290 ft ✓

² Highest point: Mt Te Waka, grid ref. N 114 092713, alt. 3351 ft

³ Grid ref. N 124 985692, alt. 3244 ft



INDIGENOUS PLANTS

GYMNOSPERM TREES

<i>Dacrycarpus dacrydioides</i>	W M	<i>Mastigodiosma excelsum</i> var. <i>excelsum</i>	W
<i>Dacrydium cupressinum</i>	W M	<i>Melicytus lanceolatus</i> (incl. M.l. var. <i>laticor</i>)	W M
<i>Phyllocladus alpinus</i> s.s.	W M	<i>M. ramiflorus</i> ssp. <i>ramiflorus</i>	W M
<i>Podocarpus hallii</i>	W M	<i>Mida salicifolia</i> (incl. M.s. var. <i>myrtifolia</i>)	W
<i>P. totara</i> s.s.	W M	<i>Myoporum laetum</i> var. <i>laetum</i>	W
<i>P. hallii</i> × <i>P. nivalis</i>	M	<i>Myrsine australis</i>	W M
<i>Prumnopitys ferruginea</i>	W M	<i>M. salicina</i>	W M
<i>P. taxifolia</i>	W M	<i>Nestegis cunninghamii</i>	W M

MONOCOT TREES

<i>Cordyline australis</i>	W M	<i>N. lanceolata</i>	W M
<i>C. banksii</i>	W M	<i>N. montana</i>	M
<i>C. indivisa</i>	W M	<i>Nothofagus fusca</i>	W M
<i>C. australis</i> × <i>C. banksii</i>	W M	<i>N. solandri</i> var. <i>solandri</i>	W M
		<i>N. fusca</i> × <i>N. solandri</i>	W M

DICOT TREES

<i>Alectryon excelsus</i> var. <i>excelsus</i>	W M	<i>Pennantia corymbosa</i>	W M
<i>Aristotelia serrata</i>	W M	<i>Pittosporum eugenioides</i>	W M
<i>Beilschmiedia tawa</i>	W M	<i>P. ralphii</i>	W M
<i>Carpodetus setatus</i>	W M	<i>P. tenuifolium</i> ssp. <i>tenuifolium</i>	W M
<i>Cotylocarpus laevigatus</i>	M	<i>P. t. ssp. t. → P. t. ssp. colensoi</i>	W
<i>Dodonaea viscosa</i> var.	M	<i>P. ralphii</i> × <i>P. tenuifolium</i>	W
<i>Elaeocarpus dentatus</i> (incl E.d. var. <i>obovatus</i>)	M	<i>Plagianthus regius</i> var. <i>regius</i>	W M
<i>E. hookerianus</i>	W M	<i>Pseudopanax arboreus</i> var. <i>arborescens</i>	W M
<i>Fuchsia excorticata</i>	W M	<i>P. colensoi</i> s.s.	W M
<i>Griselinia littoralis</i>	W M	<i>P. crassifolius</i>	W M
<i>Hedycarya arborea</i>	W	<i>P. edgerleyi</i>	W M
<i>Hoheria populnea</i> var. <i>lanceolata</i>	W M	<i>P. simplex</i> (incl. <i>P. s. var. sinclairii</i>)	W M
<i>Ixerba brexioides</i> (Elder 1959, p. 15)	W		
<i>Knightia excelsa</i>	W M		
<i>Lophomyrtus obcordata</i>	W M		

<i>Quintinia serrata</i> (incl. <i>Q.acutifolia</i> and <i>Q. elliptica</i>)	M	<i>C. perpusilla</i> ssp. <i>perpusilla</i>	M
<i>Schefflera digitata</i>	W M	<i>C.petriei</i>	W
<i>Sophora microphylla</i>	W M	<i>C.propinqua</i> ssp. <i>propinqua</i>	W
<i>S.tetraptera</i>	W M	<i>C.rhamnoides</i> (incl. <i>C.poly-morpha</i>)	W
<i>S.microphylla</i> X <i>S.tetraptera</i>	W	<i>C.rigida</i>	W
<i>Streblus heterophyllus</i>	W	<i>C.robusta</i>	W M
<i>Wainmannia racemosa</i> var. <i>racemosa</i>	W M	<i>C.rotundifolia</i>	W
DICOT SHRUBS			
<i>Alseuosmia pusilla</i>	W	<i>C.rubra</i>	W
<i>Aristotelia fruticosa</i> var. <i>microphylla</i>	W M	<i>C.rugosa</i> ssp. <i>rugosa</i>	W M
<i>Brachyglottis repanda</i> s.s.	W M	<i>C.tenuifolia</i>	W M
<i>B.rotundifolia</i> var. (<i>B.elaeagnifolia</i>)	M	<i>C.sp.(i)</i> (unnamed; included in <i>C.parviflora</i> by Oliver, 1935, and others) (If undersurface not minutely hairy as is usual in <i>C.parviflora</i> ; drupes vs unpigmented, occ. purplish red)	W M
<i>Carmichaelia arborea</i> var. (<i>C. odorata</i>)	W M	<i>C.sp.(ii)</i> (unnamed; aff. <i>C. cheesemanii</i>) (lvs broader than in <i>C. cheesemanii</i>)	W M
<i>C. Sp. (subgenus Carmichaelia)</i>	W M	<i>C.propinqua</i> X <i>C.robusta</i>	W M
<i>Cassinia vauvilliersii</i>	W M	<i>C.p.</i> X <i>C.tenuifolia</i>	W
<i>Clinanthus puniceus</i> (formerly on "Hiru-o-Tureia" and "Awa-o-Totara" — Guthrie-Smith 1953)	M	<i>C.robusta</i> X <i>C.rugosa</i>	M
<i>Coprosma acerosa</i> ssp.	W M	<i>Cotoneaster arborea</i>	W M
<i>C. areolata</i>	W M	<i>C.kingiana</i>	W M
<i>C. cheesemanii</i> ss.	W M	<i>C.arborea</i> X <i>C.kingiana</i>	W M
<i>C. crassifolia</i>	W	<i>Corokia cotoneaster</i> var. <i>cotoneaster</i>	W
<i>C. depressa</i>	M	<i>Cyathodes empetrifolia</i>	W M
<i>C. foetidissima</i>	M	<i>C.juniperina</i> agg.	W M
<i>C. grandifolia</i>	M		
<i>C.linariifolia</i> (Oliver 1935, Elder 1950)	W		
<i>C.lucida</i> s.s.	W M		

<i>Dracophyllum strictum</i>	W	<i>Myrsine divaricata</i> s.s.	W M
<i>D. subulatum</i>	W M	<i>Neomyrtus pedunculata</i>	W M
<i>Epacris alpina</i>	W M	<i>Olearia arborescens</i>	W M
<i>Fuchsia excorticata</i> X <i>F. perscandens</i>	W M	<i>O. furfuracea</i> var.	W M
<i>Gaultheria antipoda</i>	W M	<i>O. hectorii</i> var. (lvs oval, smaller than in var. <i>hectorii</i> , c. 1.5 x 1 cm)	W
<i>G. macrostigma</i>	W M	<i>O. ilicifolia</i>	W M
<i>G. sp.</i> (unnamed; included in <i>G.</i> <i>depressa</i> , as var. <i>novae-zelandiae</i> , by Franklin 1962)	W M	<i>O. nummulariifolia</i> var. <i>nummul-</i> <i>aria</i> <i>folia</i>	W M
<i>G. antipoda</i> X <i>G. sp.</i> (unnamed)	M	<i>O. paniculata</i>	M
<i>G. a.</i> X <i>G. macrostigma</i>	W	<i>O. rani</i> var. <i>colectata</i>	W M
<i>G. macrostigma</i> X <i>G. sp.</i> (unnamed)	M	<i>O. virgata</i> var. <i>virgata</i> (incl. O.v. var. <i>tamuliflora</i>)	M
<i>Genicetoma rupestre</i> var. <i>lig-</i> <i>ustifolium</i>	W M	<i>O. v.</i> var. (lvs smaller than in O.v. var. <i>virgata</i>)	W M
<i>Hebe colensoi</i> (incl. <i>H.c.</i> var. <i>hillii</i>)	W	<i>O. furfuracea</i> X <i>O. ilicifolia</i>	M
<i>H. odora</i> s.s.	M	<i>Pentachondra pumila</i>	M
<i>H. stricta</i> var. <i>stricta</i>	W M	<i>Pernettya macrostigma</i>	W M
<i>H. sp. (i)</i> (<i>H. stricta</i> var. <i>agmenitiana</i>)	M	<i>Pimelea longifolia</i> var. <i>longifolia</i>	M
<i>H. sp. (ii)</i> (<i>Veronica squallida</i>)	M	<i>P. prostrata</i> s.s.	W M
<i>H. sp. (iii)</i> (<i>Veronica arborea</i>)	M	<i>P. atridula</i> var. (prostrate shrubs; lvs up to 1cm long, densely hairy on both surfaces; fl red)	W M
<i>Helichrysum aggregatum</i> var. agg.	W M	<i>P. atridula</i> X <i>P. prostrata</i>	M
<i>Ileostylus micranthus</i>	W	<i>Pittosporum divaricatum</i> (incl. <i>P. linicarum</i>)	IV
<i>Korthalsella lindsayi</i> s.s.	W	<i>Pomaderris ericifolia</i>	M
<i>Kunzea ericoides</i> agg.	IV M	<i>Pseudopanax anomalus</i>	W M
<i>Laptespermum scoparium</i>	W M	<i>Pseudowintera axillaris</i>	W M
<i>Leucopogon fasciculatus</i> var.	W M	<i>P. colorata</i>	W M
<i>L. fraseri</i> s.s.	W M		
<i>Malicope simplex</i>	W	<i>Solanum aviculare</i> var. <i>aviculare</i>	M
<i>Malicytus alpinus</i> s.s.	W M		
<i>M. micranthus</i> (incl. <i>M.m.</i> var. <i>micro-</i> <i>phyllus</i>)	W	<i>Tupaia antarctica</i>	W
		<i>Urtica ferox</i>	W M

MONOCOT LIANES

<i>Ripogonum scandens</i>	W M
---------------------------	-----

DICOT LIANES AND RELATED PLANTS

<i>Calystegia tugutiorum</i>	W
<i>Clematis foetida</i>	W M
<i>C. forsteri</i>	W M

<i>C. paniculata</i>	W M
<i>C. quadibracteolata</i>	W M
<i>Metrosideros colensoi</i> (incl. <i>M.c.</i> var. <i>pendens</i>)	W M

<i>M. diffusa</i>	W M
<i>M. perfotata</i>	W M
<i>Muchlenbeckia australis</i>	W M
<i>M. axillaris</i>	W M
<i>M. ephedroides</i>	W M
<i>M. australis</i> X <i>M. axillaris</i>	W M
<i>Parsonsia capsularis</i> var. <i>P. heterophylla</i>	W M
<i>P. capsularis</i> X <i>P. heterophylla</i>	W
<i>Passiflora quadrangularis</i>	M
<i>Rubus australis</i>	W
<i>R. cissoides</i> var. <i>cissoides</i>	W M
<i>R. c. var.</i> (leaflets narrower than in <i>R. c. var. cissoides</i>)	W M

<i>R. schmideloides</i> var. <i>schmideloides</i> (incl. <i>R.s. var. subpacipentus</i>)	W M
---	-----

<i>R. squarrosus</i>	M
<i>R. cissoides</i> var. <i>cissoides</i> ⇔ var. (leaflets narrower)	M

PSILOPSIDS AND LYCOPODS

<i>Tmesipteris elongata</i> (incl. <i>T.e. ssp. robusta</i>)	W M
<i>T. tannensis</i>	M
<i>Lycopodium fastigiatum</i>	W M
<i>L. scariosum</i>	W M
<i>L. varium</i> (incl. <i>L. billardierei</i> and <i>L. novae-zelandicum</i>)	W M

<i>L. volubile</i>	W M
--------------------	-----

FERNS

<i>Adiantum cunninghamii</i>	W M
<i>Asplenium bulbiferum</i>	W M
<i>A. flabellifolium</i> agg.	W
<i>A. flaccidum</i>	WM
<i>A. gracillimum</i>	WM
<i>A. hookerianum</i>	W M
<i>A. lyallii</i>	W M
<i>A. oblongifolium</i>	W M
<i>A. polyodon</i>	W M
<i>A. terrestre</i>	W M
<i>A. sp. (A. trichomanes agg.)</i>	M
<i>A. bulbiferum</i> X <i>A. flaccidum</i>	W
<i>A. b. X A. hookerianum</i>	W M
<i>A. flaccidum</i> X <i>A. hookerianum</i>	W
<i>A. f. X A. lyallii</i>	W M
<i>A. hookerianum</i> X <i>A. lyallii</i>	W
<i>A. h. X A. terrestre</i> (Brownsey 1977)	W
<i>Azolla filiculoides</i>	W M

<i>Blechnum chambersii</i>	W M	<i>Hymenophyllum bivalve</i>	W
<i>B. colensoi</i>	W M	<i>H. cupressiforme</i>	M
<i>B. discolor</i>	M	<i>H. demissum</i>	W M
<i>B. fluviatile agg.</i>	W M	<i>H. dilatatum</i>	W M
<i>B. membranaceum</i>	W M	<i>H. flabellatum</i>	M
<i>B. minus</i>	W M	<i>H. minimum</i>	M
<i>B. pennatifida</i>	W M	<i>H. multifidum</i>	W M
<i>B. procerum</i>	W M	<i>H. pulchellum</i>	W
<i>B. Vulcanicum</i>	W M	<i>H. ratum</i>	W M
<i>B. sp(i) (unnamed) (common lowland sp.; lower pinnae reduced in length)</i>	W M	<i>H. revolutum</i>	M
<i>B. sp(ii) (unnamed) (cliff sp.; lower pinnae not reduced)</i>	W M	<i>H. sanguineolentum agg.</i>	W M
<i>Bettychium australe</i>	W	<i>H. scabrum</i>	W
<i>B. bifforme</i>	W M	<i>H. villosum</i>	M
<i>Ctenopteris heterophylla</i>	W M	<i>Hypolepis ambigua</i>	W M
<i>Cyathea dealbata</i>	W M	<i>H. millefolium</i>	W M
<i>C. medullaris</i>	W M	<i>H. tofobarbata</i>	W M
<i>C. smithii</i>	W M	<i>Lastreopsis glabella</i>	W M
<i>Cystopteris tasmanica</i>	W M	<i>L. hispida</i>	M
<i>Dicksonia fibrosa</i>	W M	<i>L. microsora</i> ssp. <i>pentangularis</i>	M
<i>D. lanata</i> s.s.	W M	<i>L. velutina</i>	M
<i>D. squarrosa</i>	W M	<i>Leptolepia novae-zelandiae</i>	W M
<i>Gleichenia dicarpa</i>	W M	<i>Leptopteris hymenophylloides</i>	W M
<i>Grammitis billardierei</i>	W M	<i>L. superba</i>	W M
<i>G. patagonica</i>	W M	<i>L. hymenophylloides</i> X <i>L. superba</i>	W
<i>G. magellanica</i> ssp. <i>nothofagi</i>	M	<i>Ophioglossum coriaceum</i> agg.	W M
<i>G. poeppigiana</i>	M	<i>Paesia scaberula</i>	W M
<i>G. patagonica</i> X <i>G. poeppigiana</i>	M	<i>Pallaea rotundifolia</i>	W M
<i>Histeropteris incisa</i>	W M	<i>Phymatosorus novae-zelandiae</i>	W
		<i>P. pustulatus</i>	W M
		<i>Pneumatopteris pennigera</i>	W M
		<i>Polystichum richardii</i>	W M
		<i>P. silvaticum</i>	W M
		<i>P. vestitum</i>	W M
		<i>P. richardii</i> X <i>P. vestitum</i>	W
		<i>Pteridium esculentum</i>	W M
		<i>Pteris pendula</i>	W
		<i>P. tremula</i>	W M
		<i>Pyrrhia cleagnifolia</i>	W M
		<i>Rumohra adiantiformis</i>	W

GRASSES

7

<i>Sticherus cunninghamii</i>	W M
<i>Trichomanes venosum</i>	W M
ORCHIDS	
<i>Acianthus sinclairii</i>	W
<i>Cyrtostylis teniformis</i> ss.	W
<i>Aporostylis bifolia</i>	W M
<i>Caladenia carnea</i> agg.	W M
<i>Chiloglottis cornuta</i>	M
<i>Cotybas mactanthus</i>	W M
<i>C. rivularis</i> s.s.	W M
<i>C. trilobus</i> s.s.	M
→ <i>Dactylobium cunninghamii</i>	M
<i>Eatina autumnalis</i>	M
<i>E. mucronata</i> ss.	W M
<i>Gastroleia cunninghamii</i>	M
<i>G. minor</i>	M
→ <i>Microlis cligantha</i>	W M
<i>M. unifolia</i>	W M
<i>Ottoceras novae-zelandiae</i>	M
<i>Prasophyllum colensoi</i>	W M
<i>Genoplesium nudum</i>	M
<i>Pterostylis albula</i>	W
<i>P. foliata</i>	M
<i>P. patens</i>	W M
<i>P. fimbriata</i>	W M
<i>P. sp. (unnamed; aff. P. montana)</i>	W M
<i>Spiranthes novae-zelandiae</i>	W
<i>Thelymitra decora</i>	W
<i>T. hatchii</i>	M
<i>T. longifolia</i>	W M
<i>T. pauciflora</i>	M

<i>Echinopogon ovatus</i>	M
<i>Elymus harduroides</i>	W M
<i>E. multiflorus</i>	M
<i>E. sp. (E. rectisetus agg.; Group</i>	
Waioatu of Löve & Connor, (1982)	W M
<i>Agrostis personata</i>	M
<i>Chionochloa flavicans</i>	W M
<i>C. pallens</i> ssp. <i>pallens</i>	W
<i>C. rubra</i> ssp. <i>rubra</i>	W M
<i>Cortaderia fulvida</i>	W M
<i>Deschampsia novae-zelandiae</i>	W M
<i>Deyeuxia aucklandica</i>	M
<i>D. avenoides</i> (incl. <i>D. a. var.</i> <i>brachyantha</i>)	W M
<i>Dichelachne erinacea</i>	W M
<i>Festuca multinetidis</i>	W M
<i>F. novae-zelandiae</i>	W M
<i>Hierochloe recutita</i>	M
<i>H. redolens</i>	W M
<i>Isachne globosa</i>	M
<i>Lachnagrostis striata</i>	W
<i>L. sp.(i) (unnamed; included in <i>L. filiformis</i>, as var. <i>semi-</i> <i>glabra</i>, by Zofov 1965)</i>	W M
<i>L. sp.(ii) (unnamed)</i>	W
<i>Microlaena avenacea</i>	W M
<i>M. stipoides</i>	W M

<i>Poa anceps</i> ssp <i>aneaps</i>	W M	<i>C. dipsacea</i> var. (<i>C. tahoata</i>)	W M
<i>P. bteriglumis</i>	W M	<i>C. dissita</i> s.s.	W M
<i>P. cita</i> ssp.	W M	<i>C. echinata</i> var. (<i>C. stellulata</i> var. <i>australis</i>)	W M
<i>P. colensoi</i>	M	<i>C. enysii</i>	
<i>P. imbecilla</i>	W M	<i>C. flagellifera</i>	W M
<i>P. matthewsii</i>	W	<i>C. forsteri</i>	W M
<i>P. pusilla</i>	WM	<i>C. inversa</i>	W M
		<i>C. lessoniana</i>	M
<i>Rytidosperma buchananii</i>	W M	<i>C. maotica</i>	W M
<i>R. clavatum</i>	W M	<i>C. rubicunda</i>	W M
<i>R. gracile</i>	W M	<i>C. secta</i> s.s.	W M
<i>Thisetum</i> sp.(i) (unnamed) (?)	M	<i>C. sinclairii</i>	M
		<i>C. solandri</i>	W M
		<i>C. spinostachys</i> (Hamlin 1958, as <i>C. vacillans</i>)	M
		<i>C. testacea</i> s.s.	W M
<i>T. sp.(ii)</i> (unnamed; aff. <i>T. antarcticum</i>) (If sheath distinctly hairy)	W M	<i>C. virgata</i>	W M
<i>T. sp.(ii)</i> (unnamed; aff. <i>T. antarcticum</i>) (If sheath glabrous, tubular but seen splitting)	W M	<i>C. sp.</i> (unnamed; aff. <i>C. geminata</i>) (much larger than <i>C. geminata</i> , lvs up to 2.5cm wide)	M
<i>T. sp.(iv)</i> (unnamed) (plant slender; lf sheath glabrous, or nearly so)	W M	<i>C. comans</i> X <i>C. rubicunda</i>	M
<i>T. sp.(i)</i> X <i>T. sp.(ii)</i>	M	<i>C. c.</i> X <i>C. dipsacea</i>	W
<i>T. sp.(ii)</i> X <i>T. sp.(iii)</i>	W M	<i>C. diandra</i> X <i>C. secta</i>	M
		<i>C. d.</i> X <i>C. virgata</i> (?)	M
<i>Zoysia minima</i>	W	<i>C. dipsacea</i> X <i>C. rubicunda</i>	M
		<i>Catpha alpina</i> s.s.	W
		<i>Cyperus ustulatus</i>	M
		<i>Eleocharis acuta</i>	W M
		<i>E. gracilis</i>	W M
		<i>Gahnia pauciflora</i>	W M
		<i>Lepidosperma australe</i>	W M
		<i>Machaerina sinclairii</i>	W M
		<i>Moratotia affinis</i>	M
		<i>Oreobolus pectinatus</i>	W M
		[Isolopis - from p.9]	

[top p.8]			
<i>Schoenus apogon</i>	M	<i>J. planifolius</i>	W M
<i>S. maschalinus</i>	W M	<i>J. pusillus</i>	M
<i>S. pauciflorus</i> s.s.	W	<i>J. sp.</i>	W
<i>Isolepis caliginis</i>	M	<i>Luzula decipiens</i>	W M
<i>I. carnua</i>	W	<i>L. migrata</i>	W M
<i>I. distigmatosa</i>	W	<i>L. picta</i> s.s.	W M
<i>I. habra</i>	W M	<i>L. subclavata</i>	W M
<i>I. nodosa</i>	M		
<i>I. pottsii</i>	W M		
<i>I. subtilissima</i>	W		

MONOCOT HERBS (CTHER THAN)

[ORCHIDS, GRASSES, SEDGES, RUSHES]

<i>Uncinia banksii</i>	W	<i>Arthropodium candidum</i>	W M
<i>U. caespitosa</i>	M	<i>Astelia fragrans</i>	W M
<i>U. clavata</i>	M	<i>Colospermum microspermum</i>	W M
<i>U. distans</i>	W	<i>Dianella nigra</i>	W
<i>U. divaricata</i>	W M	<i>Herpolition novae-zelandiae</i>	W M
<i>U. egmontiana</i>	M	<i>Iphigenia novae-zelandiae</i>	W
<i>U. fettuginea</i>	VI M	<i>Lemna</i> sp. (<i>L. minor</i> auct.)	W M
<i>U. gracilenta</i>	W	<i>Libertia grandiflora</i>	W M
<i>U. laxiflora</i>	W M	<i>L. ixoides</i>	W M
<i>U. tubra</i>	W M	<i>Luzuriaga parviflora</i>	M
<i>U. rupestris</i> (incl. <i>U. angustifolia</i>)	W M	<i>Phormium cookianum</i>	W M
<i>U. scabra</i>	W M	<i>Potamogeton cheesemanii</i>	W M
<i>U. uncinata</i>	W M	<i>P. suboblongus</i>	W
<i>U. vitidis</i>	M	<i>Typha orientalis</i>	W M
<i>U. zotovii</i>	W M		
<i>U. sp. (U. affinis Hamlin 1959, non U. bipartita var. affinis Col ex Clarke)</i>	W M		
<i>U. rupestris</i> X <i>U. zotovii</i> (?)	M		

RUSHES

<i>Juncus caespiticius</i>	M	<i>Brachyscome radicata</i>	M
<i>T. gregiflorus</i>	W M		
<i>T. novae-zelandiae</i>	W M		
<i>T. pallidus</i>	M		
		<i>Calmisia glandulosa</i> ssp. <i>glandulosa</i> (incl. <i>C. g. var. latifolia</i>)	W M
		<i>C. gracilenta</i> var.	W M
		<i>C. spectabilis</i> ssp. <i>spectabilis</i>	W M
		<i>C. sp. (unnamed; aff. C. gracilenta)</i> (plants strongly rhizomatous)	W
		<i>C. gracilenta</i> X <i>C. spectabilis</i>	W M

[to p. 9]

<i>Cotula australis</i>	W M	<i>Pseudognaphalium</i> sp. (unnamed; aff. <i>P. luteoalbum</i>) (Ivs narrow, acute)	WM
<i>Craspedia minor</i> var. <i>minor</i> (incl. <i>C. major</i>)	W M	<i>Raoulia albosericea</i>	M
<i>C. viscosa</i>	W M	<i>R. glabra</i>	M
<i>Gnaphalium audax</i>	W M	<i>R. tenuicaulis</i> (incl. <i>R. t.</i> var. <i>dimorpha</i> and <i>R. t.</i> var. <i>pusilla</i>)	WM
<i>G. delicatum</i>	W M	<i>Senecio banksii</i> var	WM
<i>G. gymnocephalum</i>	W M	<i>S. glaucophyllum</i> ssp. <i>toa</i>	W
<i>G. involucratum</i>	W M	<i>S. sp.</i>	WM
<i>G. limosum</i> s.s.	W M	<i>S. minimus</i>	WM
<i>G. polylepis</i>	M	<i>S. quadridentatus</i>	WM
<i>G. ruahinicum</i>	M	<i>S. rufiglandulosus</i> var. <i>rufiglandulosus</i>	WM
<i>G. sphaeticum</i>	M	<i>S. waitauensis</i>	WM
<i>G. traversii</i>	W M	<i>Vittadinia australis</i> agg.	WM
<i>Anaphalis kerriensis</i> s.s.	M		
<i>A. subrigida</i>	W M		
<i>A. sp.(i) ("Helichrysum bellidioides")</i>	WM	DICOTHERBS (OTHER THAN COMPOSITES)	
<i>A. sp.(ii) (unnamed) ("Helichrysum alpinum" of Cockayne 1928)</i>	WM	<i>Acaena anserinifolia</i>	WM
<i>A. kerriensis</i> X <i>A. sp.(i)</i>	M	<i>A. inermis</i>	W
<i>Helichrysum filicaule</i>	W M	<i>A. microphylla</i> var. <i>microphylla</i>	WM
		<i>Aciphylla colensoi</i> var. <i>colensoi</i>	W
		<i>A. sp. (unnamed; aff. <i>A. squarrosa</i>)</i>	WM
		<i>Anisotome aromatica</i>	WM
		<i>Cardamine</i> sp.(i) (<i>C. debilis</i> agg.) ("Narrow Petal" of Pritchard 1957)	WM
		<i>C. sp.(iii) (unnamed; aff. <i>C. corymbosa</i>)</i>	WM
<i>Lagenifera montana</i>	W M	<i>C. sp.(iii) (<i>C. debilis</i> agg.) (capsules pubescent)</i>	WM
<i>L. pinnatifida</i>	W	<i>C. sp.(iv) (<i>C. debilis</i> agg.) ("Long Style" of Pritchard 1957)</i>	WM
<i>L. pumila</i>	W M	<i>C. sp.(v) ("Scree Race" of Pritchard 1957)</i>	M
<i>L. strangulata</i>	W	<i>Centella uniflora</i>	WM
<i>Microseris scapigera</i> var.	W M		
<i>Leptinella pusilla</i>	W		
<i>L. squalida</i> agg.	W M		

<i>Colobanthus apetalus</i> (incl. <i>C.a.</i> var. <i>alpinus</i>)	W M	
<i>C. strictus</i>	W M	
<i>Crassula sieberiana</i>	W M	
<i>Dactylanthus taylorii</i> (Cheeseman 1925, W Elder 1950)		
<i>Daviesia glochidiatus</i>	W	
<i>Dichondra repens</i> var.	W M	
<i>D. sp.(i)</i> (unnamed; aff. <i>D. brevifolia</i>)	W M	
<i>D. sp.(ii)</i> (unnamed) (Slender)	M	
<i>Drosera peltata</i> ssp. <i>articulata</i>	M	
<i>D. binata</i>	W M	<i>E. billardiereanum</i> × <i>E. pallidiflorum</i> (?) M
<i>Epilobium alsinoides</i> s.s.	W M	<i>E. brunnescens</i> × <i>E. sp</i> (unnamed) W M
<i>E. atriplicifolium</i> s.s.		<i>Euphrasia cuneata</i> W M
<i>E. billardiereanum</i> s.s.	W M	<i>E. zelandica</i> agg. M
<i>E. brunnescens</i> s.s.	W M	<i>Galium parvifolium</i> W M
<i>E. chickanthum</i>	W M	<i>G. propinquum</i> W M
<i>E. chlorifolium</i>	M	<i>G. trilobum</i> W M
<i>E. cinereum</i>	W M	<i>Gentiana bellidifolia</i> var. <i>bell.</i> M
<i>E. cockayneanum</i>	W M	<i>G. griselinachii</i> W M
<i>E. gracilipes</i>	W M	<i>G. sp.</i> (unnamed) aff. <i>G. micropylum</i> W M
<i>E. insulare</i>	W M	<i>G. potentilloides</i> var. <i>potentilloides</i> W M
<i>E. komarovianum</i>	W M	<i>G. sessiliflorum</i> var. <i>leuczel-</i>
<i>E. linnaeoides</i>	W M	<i>zlandiae</i> W M
<i>E. melanocaulon</i>	W	<i>Glossostigma elatinoides</i> W
<i>E. microphyllum</i>	W	<i>Gnophocarpus aggregatus</i> W M
<i>E. netteroides</i>	W M	<i>G. micranthus</i> ssp. <i>micranthus</i> W M
<i>E. nummulariifolium</i>	W M	<i>Gratiola sexdentata</i> W M
<i>E. pallidiflorum</i>	W M	<i>Gunnera monoeica</i> W M
<i>E. pubens</i>	W M	<i>Halcragis erecta</i> ssp. <i>erecta</i> W M
<i>E. rotundifolium</i>	W M	<i>Hydrocotyle elongata</i> W M
<i>E. teretipes</i> s.s.	W M	<i>H. heteromeria</i> M
		<i>H. hydrophila</i> W
		<i>H. mictophylla</i> W M
		<i>H. moschata</i> W M

<i>H. novae-zelandiae</i> s.s.	W M		
<i>H. pterocarpa</i>	M		
<i>H. sulcata</i>	W M		
<i>H. sp. (unnamed)</i> (= <i>H. novae-zelandiae</i> var. <i>montana</i>)	W M		
<i>Hypenticum japonicum</i> agg.	W M		
<i>Jovellana repens</i>	W		
<i>J. sinclairii</i>	W M		
 <i>Kalsteria laxa</i>	 M		
<i>Lilaeopsis ruthiana</i>	W		
<i>Limosella lineata</i>	W M		
<i>Linum monogynum</i>	W M		
<i>Mazus radicans</i>	W M		
<i>Mentha cunninghamii</i>	W M		
<i>Montia fontana</i> s.s.	W		
<i>Myosotis forstieri</i>	W M		
<i>M. petiolata</i> s.s.	W		
<i>M. pygmaea</i> s.s.	W M		
<i>M. saxosa</i> (incl. <i>M. amabilis</i>)	W M		
<i>M. spathulata</i> (incl. <i>M. s.</i> var. <i>radicata</i>)	W M		
	W M		
 <i>M. sp. (unnamed; included in <i>M. pygmaea</i>, as var. <i>drucei</i>, by Moore 1961)</i>	 W		
<i>Myriophyllum propinquum</i>	M		
<i>Nertera ciliata</i>	W M		
<i>N. depressa</i> (incl. <i>N. cunninghamii</i>)	W M		
<i>N. setulosa</i>	W M		
 <i>N. sp. (unnamed; aff. <i>N. dichondrifolia</i> (hairs present on both surfaces of lf, straight not hooked as in <i>N. dichondrifolia</i>))</i>	 W M		
 <i>Oreomytthis colensoi</i> s.s. (incl. <i>O. c.</i> var. <i>hispida</i> and <i>O. c.</i> var. <i>multifida</i>)	 W M		
<i>O. sp. (unnamed; aff. <i>O. tenuissima</i>)</i>	W M		
<i>Ourisia macrophylla</i> ssp. <i>robusta</i>	W M		
<i>Oxalis exilis</i>	W M		
<i>O. magellanica</i>	W M		
 <i>Parahebe diffusa</i> ssp. <i>diffusa</i>	 W M		
<i>P. olsonii</i>	M		
 <i>Plantago raoulii</i>	 W M		
<i>P. triandra</i> s.s.	W M		
<i>Polygonum salicifolium</i> agg.	M		
 <i>Potentilla anserinaoides</i>	 W M		
<i>Pratia angulata</i> agg.	W M		
<i>P. perpusilla</i>	W		
<i>Ranunculus amphitrichus</i>	W M		
<i>R. glabratifolius</i>	W M		
<i>R. insignis</i>	M		
<i>R. reflexus</i>	W M		
<i>R. toyi</i>	M		
 <i>R. sp. (unnamed; included in <i>R. lappaceus</i>, as var. <i>mactophyllum</i>, by Kirk 1899)</i>	 W M		
 <i>Rumex flexuosus</i>	 W		
 <i>Scandia</i> sp. (<i>S. rosifolia</i> agg.)	 W M		
 (leaflets ovate-lanceolate)	 W M		

<i>Schizolema colensoi</i>	W M	<i>Rosa rubiginosa</i>
<i>S. trifoliolatum</i>	W M	<i>Rubus fruticosus</i> agg.
<i>Scleranthus biflorus</i>	W M	<i>Salix fragilis</i>
<i>S. blockiei</i>	W M	
<i>Stackhousia minima</i>	W M	<i>Ulex europaeus</i>
<i>Stellaria decipiens</i> (incl <i>Spatiflora</i>)	W M	
<i>Urtica incisa</i>	W M	LIANES
<i>Viola cunninghamii</i>	W M	<i>Calystegia sylvatica</i>
<i>V. filicaulis</i>	W M	<i>Solanum dulcamara</i>
<i>Wahlenbergia albomarginata</i> var. pygmaea	W M	

DOUBTFUL AND ERRONEOUS RECORDS

<i>Astelia nervosa</i> (Elder 1950)	
<i>Brachyscome sinclairii</i> (Guthrie-Smith 1953)	
<i>Celmisia incana</i> (Guthrie-Smith 1953)	
<i>Coprosma parviflora</i> (Clivat 1935, Elder 1950)	
<i>Deyeuxia quadriseta</i> (Guthrie-Smith 1953)	
<i>Pittosporum australum</i> (Allan 1961)	
<i>Raoulia australis</i> (Guthrie-Smith 1953)	
<i>Trisetum antarcticum</i> (Guthrie-Smith 1953)	
<i>Lagenifera cuneata</i> (Elder 1950)	

ADVENTIVE PLANTS

(excluding vicinity of homesteads)

TREES AND SHRUBS

<i>Acacia decurrens</i>	
<i>Crataegus monogyna</i>	
<i>Lupinus arboreus</i>	<i>Cytisus scoparius</i>
<i>Pinus radiata</i>	
<i>Ribes uva-crispa</i>	

<i>Rosa rubiginosa</i>	
<i>Rubus fruticosus</i> agg.	
<i>Salix fragilis</i>	

LIANES

<i>Calystegia sylvatica</i>	
<i>Solanum dulcamara</i>	

GRASSES

<i>Agrostis capillaris</i>	
<i>Aira caryophyllea</i>	
<i>Alopecurus geniculatus</i>	
<i>Anthoxanthum odoratum</i>	
<i>Bromus hordeaceus</i>	
<i>B. sterilis</i>	
<i>Cynodon dactylon</i>	
<i>Cynosurus cristatus</i>	
<i>Dactylis glomerata</i>	
<i>Elymus rectisetus</i> s.s.	
<i>Festuca arundinacea</i> s.l.	

<i>F. nigrescens</i>	
<i>Holcus lanatus</i>	
<i>Hordeum murinum</i>	
<i>Lolium perenne</i>	
<i>Poa annua</i>	

<i>P. pratensis</i>	
<i>P. trivialis</i>	
<i>Rytidosperma penicillatum</i>	
<i>R. racemosum</i>	

SEDGES

<i>Carex ovalis</i>	
<i>Isolepis setacea</i>	

RUSHES

Juncus articulatus
J. bufonius
J. effusus
J. inflexus
J. tanus

COMPOSITE HERBS

Achillea millefolium
Anthemis cotula
Arctium sp.
Bellis perennis
Carduus nutans
Cirsium arvense
C. vulgare
Conyza sp.
Crepis capillaris
Gnaphalium coarctatum
G. purpureum var. ustulatum
Hieracium pilosella
Hypochoeris radicata
Lactuca serriola
Lapsana communis
Matricaria matricarioides
Mycelis muralis
Senecio jacobaea
S. sylvaticus
Soliva valdiviana

MONOCOT HERBS (OTHER THAN GRASSES, SEDGES, RUSHES)

Elodea canadensis

DICOT HERBS (OTHER THAN COMPOSITES)

Amaranthus hybridus
Aphanes microcarpa
Acaena agnipila
Arenaria serpyllifolia
Brassica sp.
Callitrichie stagnalis
Cardamine hirsuta
Cerastium fontanum ssp. vulgare
Chenopodium murale
Dianthus armeria
Digitalis purpurea
Dipsacus fullonum
Epilobium ciliatum
E. obscurum
E. parviflorum
Erodium cicutarium
Galium aparine
Gentianella molle
Hyperticum humifusum
Linum bienne
L. catharticum
Lotus pedunculatus
Marrubium vulgare
Medicago lupulina
M. polymorpha
Melilotus alba
Mimulus moschatus
Myosotis sp.
Nasturtium officinale
Navarretia squarrosa
Nymphaea alba
Otobauchie minor
Potentilla viscosa
Pastinaca sativa

Plantago hirsutella

P. lanceolata

P. major

Polycarpon tetraphyllum

Polygonum aviculare agg.

P. hydropiper

Prunella vulgaris

Ranunculus acris

R. repens

R. sceleratus

Rumex acetosella

R. crispus

R. obtusifolius

Sagina apetala

S. procumbens

Sisymbrium officinale

Solanum nodiflorum

Stachys sylvatica

Stellaria media

Trifolium arvense

T. dubium

T. pratense

T. repens

T. sp.

Urtica urens

Verbascum thapsus

Verbena officinalis

Veronica arvensis

V. serpyllifolia

Wahlenbergia marginata

No. cf indigenous spp. : 547.

No. cf adventive spp. : 125.

REFERENCES

- Allan, H.H. 1961: "Flora of New Zealand". Government Printer, Wellington. Vol. I. 1085 pp.
- Brownsay, P.J. 1977: Asplenium hybrids in the New Zealand flora. N.Z. Jl Bot. 15: 601-37.
- Cheeseman, T.F. 1925: "Manual of the New Zealand Flora". Ed. 2, Edited by W.R.B. Oliver. Government Printer, Wellington. 1163 pp.
- Cockayne, L. 1928: "The Vegetation of New Zealand". 2nd ed. Eulemann, Leipzig. 456 pp.
- Edgar, E. 1970: Cyperaceae. Pp. 167-285 in "Flora of New Zealand". Government Printer, Wellington - Vol. 2. 354 pp.
- Elder, N.L. 1950: Balls Clearing and Hutchinson Bush. Bull. Wellington bot. Soc. 23: 1-8.
- 1959: Vegetation of the Kaweka Range. Trans. R. Soc. N.Z. 87: 9-26.
- Franklin, D.A. 1962: The Ericaceae in New Zealand (Gaultheria and Pernettya). Trans. R. Soc. N.Z. 1: 155-73.
- Guthrie-Smith, H. 1953: "Tutira: The Story of a New Zealand Sheep Station". Ed. 3. A.H. and A.W. Reed.
- Hamlin, B.G. 1958: Studies in New Zealand Carices, 4 - The Section Elatae Kükenthal. Trans. R. Soc. N.Z. 85: 387-96.
- 1959: A revision of the genus Uncinia (Cyperaceae) in New Zealand. Dom. Mus. Bull. 19. 106 pp.
- Kirk, T. 1899: "The Students' Flora of New Zealand." Government Printer, Wellington. 408 pp.
- Moorc, L.B. 1961: Boraginaceae. Pp. 806-33 in "Flora of New Zealand". Government Printer, Wellington. Vol. I. 1085 pp.
- Oliver, W.R.B. 1935: The genus Coprosma. Bishop Mus. Bull. 132. 207 pp.
- Löve, A.; Connor, H.E. 1982: Relationships and taxonomy of New Zealand wheat grasses. N.Z. Jl Bot. 20: 169-86.

- Petrie, D. 1902: Descriptions of new native plants, and notes. Trans. N.Z. Inst.
34: 390-6.
- Pritchard, G.G. 1957: Experimental taxonomic studies on species of Cordamine
Linn. in New Zealand. Trans. R. Soc. N.Z. 85: 75-89.
- Raven, P.H.; Englehorn, T. 1971: New taxa and new combinations in
Australasian Eriogonum (Onagraceae). N.Z. Jd Bot. 9: 345-50.
- Zotov, V.D. 1965: Grasses of the subantarctic islands of the New Zealand
region. Rec. Dom. Mus. Wellington 5: 101-46.

