

MOSSES of KAITAWA SCENIC RESERVE, MANGAONE

Prepared 4 August 1990 by the Wellington Botanical Society including Patrick Brownsey, Barbara Polly, Darea Sherratt and Rodney Lewington.

Acrocladium cuspidatum  
Acrophyllum dentatum  
Acrophyllum quadrifarium  
Atrichum androgynum  
Breutelia pendula  
Bryum billardieri  
Calyptopogon mnioides  
Campylopus clavatus  
Campylopus introflexus  
Catagonium nitens ssp nitens  
Catharomnion ciliatum  
Cladomnion ericoides  
Cryphaea dilatata  
Cyathophorum bulbosum  
Cyrtopus setosus  
Dawsonia suberba var superba  
Dicnemon calycinum  
Dicranoloma billardierei  
Dicranoloma menziesii  
Dicranoloma robustum var robustum  
Distichophyllum microcarpum  
Ditrichum difficile  
Echinodium hispidum  
Fissidens asplenioides  
Fissidens rigidulus  
Fissidens sp  
Homalia falcifolia  
Hypnodendron arcuatum  
Hypopterygium commutatum  
Hypopterygium filiculaeforme  
Leucobryum candidum  
Lopidium concinnum  
Macromitrium helmsii  
Macromitrium sp  
Mittenia plumula  
Neckera pennata  
Papillaria flavo-limbata  
Polytrichadelphus magellanicus  
Polytrichum juniperinum  
Psilopilum crispulum  
Ptychomnion aciculare  
Pyrrhobryum bifarium  
Racopilum convolutaceum  
Rhizogonium distichum  
Sematophyllum amoenum  
Tetraphidopsis pusilla  
Thamnobryum pandum  
Thuidium furfurosum  
Thuidium laeviusculum  
Trachyloma diversinerve  
Weymouthia cochlearifolia  
Weymouthia mollis  
Wjikia extenuata