

file: kortmar2.doc(060697)

On base ✓
Q. b. b. c.

Re. *Korthalsella lindsayi* at Riverside Road, Martinborough, S27.188.008 etc.

Korthalsella lindsayi was found at a number of sites (all within 1 km of each other) at Riverside Road, Riverside Station, Martinborough, on 26.05.1996.

Site 1 - S27.188.008, gully, grazed forest remnant:

<i>Korthalsella lindsayi</i>	mistletoe
<i>Podocarpus totara</i>	totara
<i>Prumnopitys taxifolia</i>	matai
<i>Alectryon excelsus</i>	titoki
<i>Coprosma areolata</i>	
<i>C. crassifolia</i>	
<i>C. propinqua</i>	
<i>C. rhamnoides</i>	
<i>Melicytus micranthus</i>	
<i>M. ramiflorus</i>	
<i>Myrsine divaricata</i>	weeping matipo
<i>Sophora microphylla</i>	kowhai
<i>Streblus heterophyllus</i>	small-leaved milk tree
<i>Ripogonum scandens</i>	supplejack
<i>Muehlenbeckia australis</i>	
<i>M. complexa</i>	

Site 2 - S27.188.010, scattered trees on a west facing hill slope, grazed.

<i>Korthalsella lindsayi</i> *	mistletoe
<i>Podocarpus totara</i>	totara
<i>Alectryon excelsus</i>	titoki
<i>C. crassifolia</i>	
<i>C. rhamnoides</i>	
<i>Cordyline australis</i>	cabbage tree
<i>Kunzea ericoides</i>	kanuka
<i>Lophomyrtus obcordata</i>	rohutu
<i>Melicytus micranthus</i>	
<i>Myoporum laetum</i>	ngaio
<i>Myrsine divaricata</i>	weeping matipo
<i>Nestegis cunninghamii</i>	black maire
<i>Sophora microphylla</i>	kowhai
<i>Streblus heterophyllus</i>	small-leaved milk tree
<i>Muehlenbeckia australis</i>	
<i>M. complexa</i>	
<i>Ripogonum scandens</i>	supplejack

* *Korthalsella lindsayi* on *Lophomyrtus obcordata*, *Coprosma crassifolia* and *Myrsine divaricata*.

Site 3 - forest remnant along a small winding stream, in a valley, grazed, S27.189.012, approx. 200 x 50-100 m, potentially very nice area. In places closed canopy and regeneration. Should be fenced off.

<i>Korthalsella lindsayi</i> *	mistletoe
<i>Dacrycarpus dacrydoides</i>	kahikatea
<i>Podocarpus totara</i>	totara
<i>Prumnopitys taxifolia</i>	matai
<i>Alectryon excelsus</i>	titoki
<i>Coprosma areolata</i>	
<i>C. crassifolia</i>	
<i>C. propinqua</i>	
<i>C. rhamnoides</i>	
<i>C. rigida</i>	
<i>C. rubra</i>	
<i>C. spp.</i>	
<i>Cordyline australis</i>	cabbage tree
<i>Hoheria sextylosa</i>	houhere
<i>Kunzea ericoides</i>	kanuka
<i>Lophomyrtus obcordata</i>	rohutu
<i>Melicope simplex</i>	
<i>Melicytus micranthus</i>	
<i>M. ramiflorus</i>	
<i>Myoporum laetum</i>	ngaio
<i>Myrsine divaricata</i>	weeping matipo
<i>Nestegis lanceolata</i>	white maire
<i>Pseudopanax anomalus</i>	
<i>Sophora microphylla</i>	kowhai
<i>Streblus heterophyllus</i>	small-leaved milk tree
<i>Muehlenbeckia australis</i>	
<i>M. complexa</i>	
<i>Parsonsia</i> spp.	
<i>Rubus schmideloides</i>	bush lawyer
<i>Ripogonum scandens</i>	supplejack

(*Carex secta* grows in the stream)

* *Korthalsella lindsayi* on *C. crassifolia*, *C. rigida*, *Lophomyrtus obcordata*, *Myrsine divaricata* and *Muehlenbeckia australis*.

Aalbert Rebergen
Conservation Officer
DOC Masterton

