

Native vascular plants of swamp forest, Waingawa, Masterton

Location : between Hughes Lane and Perrys Road south of

Waingawa River ; centre NZMS 260 Map S26/282203. 106m

Han's  
Bush

C. C. Cyc

Dept. of Conservation  
Waingawa

Based on visit with W. Baxter, 13.10.88

a = abundant ; o = occasional ; u = uncommon (or actual no. of plants seen)

### Gymnosperms

<i>Dacrycarpus dacrydioides</i>	(kahikatea)	a
<i>Dacrydium cupressinum</i>	(rimu)	l.(adult)
<i>Podocarpus totara</i>	(totara)	o
<i>Prumnopitys taxifolia</i>	(matai)	u

### DICOT TREES and SHRUBS

<i>Alectryon excelsum</i>	(titoki)	o
<i>Aristotelia serrata</i>	(winederry)	u
<i>Beilschmiedia tawa</i>	(tawa)	o
<i>Carpodetus serratus</i>	(putaputaweta)	o

*Coptosma crassifolia* u

*C. grandifolia* (kanono) u

*C. propinqua* o

*C. rhamnoides* o

*C. rigidula* u

*C. robusta* (karamu) o

*C. tenuicaulis* a

*C. propinqua* x *C. robusta* u

*Elaeocarpus hookerianus* (pokaka) l.(adult)

*Fuchsia excorticata* (kotukutuku) l

*Geniostoma rupestre* var. *ligustrifolium* (largehange) o

*Hedycarya arborea* (pigeonwood) l.

*Hoheria populnea* var. [*H. sexstylosa*]

(lacebark) u

*Kunzea ericoides* 2

*Laurelia novae-zelandiae* o

*Leptospermum scoparium* o

*Melicytus ramiflorus* o

<i>Nestegis lanceolata</i>	(white maire)	1.
<i>Grevillea virgata</i> var. <i>virgata</i>		a
<i>Pittosporum eugenioides</i> (lemonwood)		u
<i>Prumnopitys ferruginea</i> var. <i>tenuifolium</i> (Kohuhu)		u
<i>Pseudopanax arboreus</i> (fivefinger)		2.
<i>P. crassifolius</i>	(lancewood)	3.
<i>Schefflera digitata</i>	(pate)	u
<i>Solanum laciniatum</i>	(poroporo)	o
<i>Streblus heterophyllus</i> [ <i>Paratrophis microphylla</i> ]	(small-leaved milktree)	1.
<i>Syzygium maire</i>		a
<i>Weinmannia racemosa</i> var. <i>racemosa</i> (komati)		o

#### DICOT LIANES

<i>Metrosideros diffusa</i>		u
<i>M. fulgens</i>		1.
<i>M. perforata</i>		u
<i>Muehlenbeckia australis</i> (pohuehue)		o
<i>M. complexa</i> (pohuehue)		u
<i>Parsonsia heterophylla</i> (NZ jasmine)		u
<i>Passiflora tetrandra</i> (NZ passionflower)		u
<i>Rubus australis</i> (lawyer)		o
<i>R. australis</i> X <i>R. complexa</i>		u

#### MONOCOT TREE

<i>Cordyline australis</i> (cabbage tree)	o
---	---

#### MONOCOT LIANES

<i>Freylinetia baueriana</i> ssp. <i>banksii</i> (kiekie)	1.
<i>Ripogonum scandens</i>	a.

#### MONOCOT HERBS

<i>Astelia fragrans</i>	o
<i>Baumea tenax</i>	o
<i>Carex forsteri</i>	1.
<i>C. lessoniana</i>	la
<i>C. secta</i> s.s.	la
<i>C. virgata</i>	o

<i>Cotyledon toetoe</i>	(toetoe)	o
<i>Corybas rivularis</i>	(spider orchid)	l
<i>Elegia acuta</i>	(sharp spike sedge)	l ?
<i>E. gracilis</i>	(slender " " )	l
<i>Juncus planifolius</i>		u
<i>J. sorophorus</i>		u
<i>Lemna sp.</i> [L minor and NI] (duckweed)		l.
<i>Microseris avenacea</i> (bush ricegrass)		u.
<i>Phormium tenax</i> (NZ flax)		u.
<i>Pterostylis banksii</i> (green hood orchid)		o
<i>Schoenus maschalinus</i>		u
<i>Typha orientalis</i> (raupo)		a
<i>Uncinia uncinata</i> (hookgrass)		o

### DICOT HERBS

<i>Cardamine sp.</i> [E. debilis agg.]	
? C. "narrow petal" of Pritchard 1957	[bittercress]
<i>Callitrichia muelleri</i>	o
<i>Senecio minimus</i>	u
<i>Urtica incisa</i>	l.

### FERNS

<i>Asplenium bulbiferum</i> (hen and chicken fern)	o
<i>A. flaccidum</i> ssp. <i>flaccidum</i>	u
<i>A. gracillimum</i>	u
<i>A. oblongifolium</i> (shining spleenwort)	u
<i>A. polyodon</i>	u
<i>Azolla filiculoides</i> ssp. <i>rubra</i>	l
<i>Blechnum filiforme</i> (climbing blechnum)	o
<i>B. minus</i> (swamp kio kio)	o
<i>B. sp.</i> [B. capense agg., common lowland sp. with proximal pinnae reduced in size] (kio kio)	l.
<i>Dicksonia squarrosa</i> (wheki)	a.
<i>Histiopteris incisa</i>	u
<i>Hymenophyllum demissum</i> (filmy fern)	l.
<i>Hypolepis ambigua</i>	o
<i>Lastreopsis glabella</i>	u
<i>Paesia scaberula</i>	u
<i>Phymatosorus diversifolius</i> (hounds tongue)	u

<i>Pneumatopteris pennigera</i>	4	o
<i>Ferns</i>		1.
<i>Pyrrhosia elaeagnifolia</i>		u
<i>Trichomanes venosum</i> (filmy fern)		1.

### PLANTED NATIVE SPECIES

In forest close to farm buildings/dwellings, the following native plants were seen but are regarded as not natural to this area

<i>Corynocarpus laevigatus</i>	(karaka)
<i>Marattia salicina</i>	(king fern)
<i>Sophora microphylla</i>	(kowhai)

### MAJOR WEEDS

*Cobaea scandens* - a smothering vine in western end of forest.

*Lonicera japonica* (Japanese honeysuckle); a few in western end of forest, abundant in raupo patches.

*Rubus fruticosus* (blackberry) : abundant in raupo patches

*Selaginella krausiana* : common over c. 200m of forest floor at western end of block.

### MAIN PLANT COMMUNITIES

mature kahikatea / (mahoe) / (hangahange) : swamp forest  
 semi-mature kahikatea - [pukatea] - (swamp maire) / wheki - (supplejack),  
 swamp forest.  
 (cabbage tree) / *Olearia virgata* - *Cop. tenuicaulis* / raupo / (*Carex secta*) ; swamp  
 [*Olearia virgata*] / blackberry - Japanese honeysuckle - raupo : vine-reedland