

Indigenous plants of Greystoke Reserve, Jefferson Line, Marton.

Wanganui Botanical Group, 24.3.91; (Grid ref NZMS 260/sheet S23/173297: 230 m a.s.l); compiled by Colin Ogle and Wanganui Botanical Group, 24.03.91.

Wanganui Plant List 26

Department of Conservation, Wanganui

a = abundant; c = common; o = occasional; u = uncommon

* denotes adventive species including indigenous species not natural to this area (planted specimens are not listed)

Gymnosperms

<i>Dacrycarpus dacrydioides</i>	(kahikatea)	u (1 adult)
<i>Dacrydium cupressinum</i>	(rimu)	o
<i>Podocarpus totara</i>	(totara)	(planted? 2 dead)

Dicot trees and shrubs

<i>Alectryon excelsus</i>	(titoki)	o
<i>Beilschmiedia tawa</i>	(tawa)	c
<i>Coprosma areolata</i>		c
<i>C. grandifolia</i>	(kanono, raurekou)	o
<i>C. rigida</i>		u
<i>Corynocarpus laevigatus</i> only?)	(karaka)	u (seedlings)
<i>Elaeocarpus dentatus</i>	(hinau)	o
<i>Geniostoma rupestre</i> var. <i>ligustrifolium</i>	(hangehange)	o
<i>Hedycarya arborea</i>	(pigeonwood)	o
<i>Hoheria populnea</i> var. [= <i>H.</i> <i>sexstylosa</i>]	(lacebark)	u
* <i>Hydrangea macrophylla</i>	(hydrangea)	u
* <i>Ilex aquifolium</i>	(holly)	u
<i>Knightia excelsa</i>	(rewarewa)	u
<i>Laurelia novae-zelandiae</i>	(pukatea)	u
* <i>Ligustrum sinense</i>	(privet)	u
<i>Lophomyrtus bullata</i>	(ramarama)	u
<i>Macropiper excelsum</i>	(kawakawa)	c
<i>Melicytus ramiflorus</i>	(mahoe)	a
<i>Myoporum laetum</i>	(ngaio)	u
<i>Myrsine australis</i>	(mapou)	o
<i>Pennantia corymbosa</i>	(kaikomako)	u
<i>Pittosporum tenuifolium</i> var. <i>tenuifolium</i>	(kohuhu)	o
<i>Pseudopanax crassifolius</i>	(lancewood)	u
* <i>Sambucus nigra</i>	(elderberry)	c
<i>Schefflera digitata</i>	(pate)	u

*Solanum pseudocapsicum	(Jerusalem cherry)	u
S. sp.	(poroporo)	u
Sophora microphylla	(kowhai)	u (planted only ?)
Streblus heterophyllus	(small-leaved milktree)	o
* Ulex europaeus	(gorse)	u
* Vitex lucens	(puriri)	(u, 2 seedlings)
Monocot Tree		
Cordyline australis	(cabbage tree)	o
Dicot lianes		
Clematis foetida		u
C. forsteri (?)		u
*C. vitalba	(old man's beard)	u
*Jasminum polyanthum	(jasmine)	u
Metrosideros colensoi	(white climbing rata)	o
M. diffusa	(white climbing rata)	u
M. perforata	(white climbing rata)	u
Muehlenbeckia australis	(pohuehue)	u
Parsonia heterophylla	(NZ jasmine)	o
*Passiflora mollissima	(banana passionfruit)	a
P. tetrandra	(NZ passionflower)	o
*Tropaeolum speciosum	(Chilean flame creeper)	u
Monocot liane		
*Asparagus scandens	(climbing asparagus)	o
*Bomarea multiflora		u
Ripogonum scandens	(supplejack)	o
Dicot herbs		
*Cirsium vulgare	(Scotch thistle)	o
*Senecio bipinnatisectus	(fireweed)	u
*Solanum nigrum	(black night shade)	u
Monocot herbs		
Astelia solandri		u
Carex solandri		u
Colospermum hastatum	(perching lily)	o
Corybas trilobus	(spider orchid)	u
*Dactylis glomerata	(cock'sfoot)	o
Earina mucronata		u
Uncinia uncinata	(hook grass)	c

Ferns

<i>Asplenium bulbiferum</i>	(hen and chicken fern)	o
<i>A. flaccidum</i>	(hanging spleenwort)	u
<i>A. gracillimum</i>		c
<i>A. oblongifolium</i>	(shining spleenwort)	o
<i>A. polyodon</i>		u
<i>Blechnum filiforme</i>	(climbing blechnum)	c
<i>B. sp. [B. capense agg.</i>		
"Black spot"]	(kio kio)	u
<i>Cyathea dealbata</i>	(ponga)	o
<i>C. medullaris</i>	(mamaku)	u
<i>Dicksonia squarrosa</i>	(wheki)	u
<i>Histiopteris incisa</i>	water fern	u
<i>Hymenophyllum demissum</i>	(filmy fern)	u
<i>Hypolepis ambigua</i>		o
<i>Lastreopsis glabella</i>		u
<i>L. hispida</i>		u
<i>Paesia scaberula</i>	(ring fern)	u
<i>Pellaea rotundifolia</i>	(button fern)	c
<i>Phymatosorus diversifolius</i>	(hounds tongue)	u
<i>P. scandens</i>		o
<i>Polystichum richardii</i>		u
<i>Pteridium esculentum</i>	(bracken)	u
<i>Pteris pendula</i> [P. macilenta of NZ authors]		u
<i>P. tremula</i>		o
<i>Pyrrosia eleagnifolia</i>		u
* <i>Selaginella kraussiana</i>		u

Total = 67 indigenous species

Vegetation: tawa-titoki forest with occasional large pukatea and rewarewa, and rarely rimu and kahikatea. Understorey chiefly mahoe, hangehange, Coprosma areolata, rarely tree ferns, and an often sparse groundcover but with locally common climbing blechnum and button fern.

The main weeds throughout the forest are banana passionfruit, elderberry and climbing asparagus - the first of these needs urgent control. Most other weeds were on the margins, including a solitary plant of Old Man's Beard on the western side, where privet, holly, Bomarea, and Chilean flame creeper also occur. Other weeds are encroaching from the house garden on the eastern side, including jasmine and hydrangea; several other species have been planted into the forest edge here, and may spread in the future e.g. Crinum.